

ANNIHILATING FIELDS OF STANDARD MODULES OF $\mathfrak{sl}(2, \mathbb{C})$ AND COMBINATORIAL IDENTITIES

ARNE MEURMAN AND MIRKO PRIMC

ABSTRACT. We show that a set of local admissible fields generates a vertex algebra. For an affine Lie algebra $\tilde{\mathfrak{g}}$ we construct the corresponding level k vertex operator algebra and we show that level k highest weight $\tilde{\mathfrak{g}}$ -modules are modules for this vertex operator algebra. We determine the set of annihilating fields of level k standard modules and we study the corresponding loop $\tilde{\mathfrak{g}}$ module—the set of relations that defines standard modules. In the case when $\tilde{\mathfrak{g}}$ is of type $A_1^{(1)}$, we construct bases of standard modules parameterized by colored partitions and, as a consequence, we obtain a series of Rogers-Ramanujan type combinatorial identities.

CONTENTS

- Introduction
- 1. Formal Laurent series and rational functions
- 2. Generating fields
- 3. The vertex operator algebra $N(k\Lambda_0)$
- 4. Modules over $N(k\Lambda_0)$
- 5. Relations on standard modules
- 6. Colored partitions, leading terms and the main results
- 7. Colored partitions allowing at least two embeddings
- 8. Relations among relations
- 9. Relations among relations for two embeddings
- 10. Linear independence of bases of standard modules
- 11. Some combinatorial identities of Rogers-Ramanujan type
- References

INTRODUCTION

In this paper we give an explicit construction of standard (i.e. integrable highest weight) representations of affine Lie algebra $\tilde{\mathfrak{g}}$ of the type $A_1^{(1)}$, and as a consequence we get a series of Rogers-Ramanujan type identities.

1991 *Mathematics Subject Classification.* Primary 17B67; Secondary 05A19.

Key words and phrases. affine Lie algebras, vertex operator algebras, vertex operator formula, standard modules, loop modules, Rogers-Ramanujan identities, colored partitions, partition ideals.

Partially supported by a grant from the Göran Gustafsson Foundation for Research in Natural Sciences and Medicine and by the Ministry of Science and Technology of the Republic of Croatia

In order to describe our main result, let $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{C})$ and let as usual $\tilde{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c \oplus \mathbb{C}d$ be the corresponding affine Lie algebra. We fix the usual $\mathfrak{sl}_2$ -basis $\{x, h, y\}$ in $\mathfrak{g}$. For an element $X \in \mathfrak{g}$ we write $X(n) = X \otimes t^n$. Set

$$\bar{B} = \{b(n) \mid b \in \{x, h, y\}, n \in \mathbb{Z}\}, \quad \bar{B}_- = \{b(n) \in \bar{B} \mid n \leq 0, n < 0 \text{ if } b \neq y\}.$$

Then $\bar{B}_-$ is a basis of subalgebra $\tilde{\mathfrak{n}}_-$ of affine Lie algebra $\tilde{\mathfrak{g}}$. We define a linear order $\preccurlyeq$ on $\bar{B}$ by $\cdots \prec x(j-1) \prec y(j) \prec h(j) \prec x(j) \prec y(j+1) \prec \cdots$ and we write

$$(1) \quad u(\pi) = \prod_{a \in \bar{B}} a^{\pi(a)} = a_1 \cdots a_s$$

to denote a (finite) product in $U(\tilde{\mathfrak{g}})$ of elements $a_j \in \bar{B}$, $a_1 \preccurlyeq \cdots \preccurlyeq a_s$, where each factor $a \in \bar{B}$ of $u(\pi)$ appears precisely $\pi(a)$ times. We may think of such product $u(\pi)$ in the universal enveloping algebra simply as a colored partition $\pi: \bar{B} \rightarrow \mathbb{N}$ consisting of $\pi(a)$ parts a , each part $a = b(n)$ having a degree n and color $b \in \{x, h, y\}$. We denote the set of all colored partitions by $\mathcal{P}(\bar{B})$, and by $\mathcal{P}(\bar{B}_-)$ we denote all colored partitions with parts in $\bar{B}_-$.

The set $\{u(\pi) \mid \pi \in \mathcal{P}(\bar{B}_-)\}$ is a Poincaré-Birkhoff-Witt basis of $U(\tilde{\mathfrak{n}}_-)$, so for a highest weight $\tilde{\mathfrak{g}}$ -module V with a highest weight vector v_0 the set of vectors

$$(2) \quad u(\pi) \cdot v_0, \quad \pi \in \mathcal{P}(\bar{B}_-),$$

is a spanning set of V . In the case that $V = M(\Lambda)$ is a Verma module, the set (2) is a basis. In the case that $V = L(\Lambda)$ is a standard $\tilde{\mathfrak{g}}$ -module, the spanning set (2) is not a basis, and our main result is that for $\Lambda = k_0\Lambda_0 + k_1\Lambda_1$, $k_0, k_1 \in \mathbb{N}$, the set of vectors of the form (2) subject to conditions

$$(3) \quad \begin{aligned} \pi(y(j-1)) + \pi(h(j-1)) + \pi(y(j)) &\leq k, \\ \pi(h(j-1)) + \pi(x(j-1)) + \pi(y(j)) &\leq k, \\ \pi(x(j-1)) + \pi(y(j)) + \pi(h(j)) &\leq k, \\ \pi(x(j-1)) + \pi(h(j)) + \pi(x(j)) &\leq k, \\ \pi(x(-1)) &\leq k_0, \quad \pi(y(0)) \leq k_1 \end{aligned}$$

is a basis of standard module $L(\Lambda)$ of level $k = k_0 + k_1$.

As a consequence of this result we get a series of combinatorial identities of Rogers-Ramanujan type. For example, if we write a partition of a nonnegative integer n in the form $n = \sum_{j \geq 0} j f_j$, then the number of partitions of n such that each part appears at most once (i.e. $f_j \leq 1$ for all j) equals the number of partitions of n such that each part appears at most twice (i.e. $f_j \leq 2$ for all j), but subject to the conditions

$$(4) \quad \begin{aligned} f_{3j+2} + f_{3j+1} + f_{3j} &\leq 2, \\ f_{3j+2} + f_{3j} + f_{3j-1} &\leq 2, \\ f_{3j+1} + f_{3j} + f_{3j-2} &\leq 2, \\ f_{3j} + f_{3j-1} + f_{3j-2} &\leq 2, \\ f_1 &\leq 1, \quad f_2 \leq 1. \end{aligned}$$

Before we say something about the proof itself, let us say a few words about some general ideas involved.

If $\tilde{\mathfrak{g}}$ is an affine Lie algebra, then a standard $\tilde{\mathfrak{g}}$ -module $L(\Lambda)$ is a quotient of the Verma module $M(\Lambda)$ by its maximal submodule $M^1(\Lambda)$, and this construction works for any Kac-Moody Lie algebra. On the other hand, for affine Lie algebras we also have various vertex operator constructions of basic (or level 1) representations: The representation space is explicitly given, usually the bosonic or fermionic Fock space, and the action of the affine Lie algebra is given by the vertex operator formula, roughly of the form $X(z) = E^-(z)E^+(z)E^0(z)$, where $E^\pm(z)$ are some exponentials of Heisenberg Lie algebra elements, and $E^0(z)$ is an exponential involving “group elements” (in principle defining operators only on integrable representations).

J.Lepowsky and R.Wilson gave a Lie-theoretic interpretation and proof of the Rogers-Ramanujan identities by explicitly (explicitly in a broader sense) constructing level 3 standard representations of $\mathfrak{sl}(2, \mathbb{C})^\sim$: The vertex operator formula in the principal picture for level 1 standard modules implies (very roughly speaking) a formula of the form $X(z)^2 = E^-(z)X(z)E^+(z)E^0(z)$ relating the action of various Lie algebra elements on level 3 standard modules. In this case one cannot write the representation space “explicitly” nor an “explicit” action for the Lie algebra in terms of some formula for $X(z)$. But one can start with a Poincaré-Birkhoff-Witt basis of the Verma module $M(\Lambda)$, see the corresponding spanning set in $L(\Lambda)$, and by using the above mentioned formula, reduce this spanning set to a spanning set of $L(\Lambda)$ parameterized combinatorially by partitions satisfying the difference two conditions. At this point one could either recall the Rogers-Ramanujan identities (and a specialization of the Weyl-Kac character formula) and conclude that this is a basis, or somehow prove directly linear independence of this set and, as a consequence, prove the Rogers-Ramanujan identities. Later on came a third idea for proving linear independence, somewhere in between the first two: The correct character formula for $L(\Lambda)$ in terms of partitions satisfying difference two conditions is replaced by finding a basis of $M^1(\Lambda)$ parameterized by partitions. It should be noticed that in the principal picture “group elements” $E^0(z)$ in the vertex operator formula happen to be ± 1 , and hence are well defined operators on the Verma module.

The Lepowsky-Wilson ideas of constructing representations by using vertex operator formulas work as well in the case of the homogeneous picture, except that there the presence of $E^0(z)$ prevents one to use Verma modules: one has to choose among the two.

The starting point of our construction was the first consequence of Frenkel-Kac vertex operator formula (in the homogeneous picture) which does not contain terms $E^0(z)$: the relation

$$(5) \quad x_\theta(z)^{k+1} = 0$$

which holds on every standard $\tilde{\mathfrak{g}}$ -module $L(\Lambda)$ of level k . Here θ is the maximal root of the corresponding finite dimensional Lie algebra $\mathfrak{g}$, x_θ is a corresponding root vector and $x_\theta(z) = \sum_{n \in \mathbb{Z}} x_\theta(n)z^{-n-1}$. Of course, $x_\theta(z)^{k+1}$ is a formal Laurent series in z ; the coefficients of this series are sums of elements of the universal enveloping algebra $U(\tilde{\mathfrak{g}})$, and relation (5) means that all these coefficients are zero on $L(\Lambda)$.

Relation (5) exhibits several “universal” properties: The adjoint action of $\mathfrak{g}$ on $x_\theta(z)^{k+1}$ produces new series which vanish on $L(\Lambda)$, and these series form an irreducible representation R of $\mathfrak{g}$ with highest weight $(k+1)\theta$. Moreover, all the coefficients of the series from R form a loop $\tilde{\mathfrak{g}}$ -module $\bar{R}$ for the adjoint action of $\tilde{\mathfrak{g}}$. We call the elements of $\bar{R}$ the relations (for the standard module) because a highest weight $\tilde{\mathfrak{g}}$ -module V of level k is standard if and only if relation (5) holds on V . Besides that, $\bar{R}$ seems to be “the smallest possible” loop module that annihilates $L(\Lambda)$, certainly “the smallest possible” that can be obtained from vertex operator formula in any picture.

With regard to the Verma modules of level k we have the property that $\bar{R}M(\Lambda) = M^1(\Lambda)$ if and only if Λ is dominant integral, otherwise $\bar{R}M(\Lambda) = M(\Lambda)$. Hence in the case of the standard module $L(\Lambda)$ we have

$$(6) \quad M^1(\Lambda) = U(\tilde{\mathfrak{g}})\bar{R}v_\Lambda,$$

where v_Λ denotes a highest weight vector in $M(\Lambda)$. This result is quite similar to the one that holds in the principal picture, and allows one to try to prove “linear independence” by finding a basis in $M^1(\Lambda)$ of the form $ur \cdot v_\Lambda$, $u \in U(\tilde{\mathfrak{g}})$, $r \in \bar{R}$. Of course, as in the principal picture, one should expect that many of these elements are linearly dependent, and one should construct relations among them in order to reduce this set to a basis. This kind of relations we call relations among relations.

By an analogy with the principal picture, for $\mathfrak{sl}(2, \mathbb{C})^\sim$ we could guess correctly a space of all relations among relations that are needed: For level 1 modules it was some $\tilde{\mathfrak{g}}$ -module having a “7-dimensional” loop submodule and a “5-dimensional” loop quotient—it looked like another ad hoc argument.

The elements $x_\theta(z)^{k+1}$ are zero on $L(\Lambda)$, so they should be seen somewhere as nonzero elements. The emerging theory of vertex operator algebras provided a suitable setting where this could be achieved. It was known that the irreducible module $L(k\Lambda_0)$ could be given the structure of a vertex operator algebra in a natural way, but here the vector $f_0^{\Lambda(h_0)+1}v_\Lambda$ that should parameterize $x_\theta(z)^{k+1}$ vanishes. We thus guessed and proved that a generalized Verma module extension $N(k\Lambda_0)$ of $L(k\Lambda_0)$ also had the structure of vertex operator algebra. In $N(k\Lambda_0)$, $f_0^{\Lambda(h_0)+1}v_\Lambda$ is nonzero and this provided the required setting to explain the “universal” properties of (5). Later it was also possible to prove “all necessary” relations among relations by calculations in $N(k\Lambda_0)$: the relations among relations arise by writing the vector $x_\theta(-2)x_\theta(-1)^{k+1}\mathbf{1}$ in two different ways, thus obtaining two different expressions for the same field.

The construction of the vertex operator algebra $N(k\Lambda_0)$ in Section 3 is based on a general “generators and relations” construction of vertex operator algebras presented in Theorem 2.6. The proof of Theorem 2.6 is inspired by the ideas of P. Goddard on locality. In our case the vertex operator algebra $V = N(k\Lambda_0)$ is generated by a set of fields $x(z)$, $x \in \mathfrak{g} \cong \mathfrak{g}(-1)\mathbf{1} \subset V$. In the general construction we start with a set of series $u(z)$, $u \in U \subset V$, which are admissible, local and generate V . By using normal order products we extend this generating set to a set of mutually local fields $v(z)$ parameterized by all vectors $v \in V$, and then a general duality argument implies that V is a vertex algebra. In Section 4 we prove that a large class of level k $\tilde{\mathfrak{g}}$ -modules (including all highest weight modules) can be given the structure of modules over the vertex operator algebra $N(k\Lambda_0)$, a result that has been announced by B. Feigin and E. Frenkel and also deduced by I. B.

Frenkel and Y. Zhu by another method. Our proof uses an inductive construction of intertwining operators local with respect to the generating fields $x(z)$, $x \in \mathfrak{g}$, acting on a highest weight $\tilde{\mathfrak{g}}$ -module, and then again a general duality argument applies.

The results described above are the content of Sections 1–5, phrased and proved in terms of vertex operator algebra $N(k\Lambda_0)$. In Section 8 we construct (some) relations among relations in a manner mentioned above.

At this point we should say that by now there are several proofs of the fact that $N(k\Lambda_0)$ is a vertex operator algebra and that highest weight $\tilde{\mathfrak{g}}$ -modules are its modules. Moreover, several proofs about properties of annihilating field (5) have appeared as well (see remarks at the end of Sections 4 and 5).

In a sharp contrast to the general results on algebraic properties of relations $\bar{R}$, we can successfully apply them only to the case of $\tilde{\mathfrak{g}} = \mathfrak{sl}(2, \mathbb{C})^\sim$. The reader will find Section 6 a sort of compromise between general wishes and particular results that we can prove, at least allowing one to speculate what may be going on for higher ranks. In order to describe these results, we return to our set $\bar{B}$ defined above and the corresponding set $\mathcal{P}(\bar{B})$ of colored partitions. It is a monoid, and we often use a multiplicative notation suggested by (1). We construct a particular order $\preceq$ on $\mathcal{P}(\bar{B})$ that allows inductive arguments, behaves well with respect to the multiplicative structure and allows every nonzero element $r \in \bar{R}$ to be written in the form

$$(7) \quad r = c_\rho u(\rho) + \sum_{\pi \succ \rho} c_\pi u(\pi),$$

where $c_\rho, c_\pi \in \mathbb{C}$, $c_\rho \neq 0$. We call ρ the leading term of r and write $\rho = \ell(r)$. Moreover, the set $\ell(\bar{R})$ of leading terms of all nonzero elements in $\bar{R}$ parameterize a nice basis $\{r(\rho) \mid \rho \in \ell(\bar{R})\}$ of $\bar{R}$, where $\rho = \ell(r(\rho))$. Since $r = 0$ on $L(\Lambda)$, the product $u(\rho)$ can be expressed in terms of $u(\pi)$, $\pi \succ \rho$, and by induction one easily gets that

$$(8) \quad u(\pi) \cdot v_{k\Lambda_0}, \quad \pi \in \mathcal{P}(\bar{B}_-), \quad \pi \notin (\mathcal{P}(\bar{B}_-) \cdot \ell(\bar{R})) \cup (\mathcal{P}(\bar{B}_-) \cdot y(0)),$$

is a spanning set of $L(k\Lambda_0)$. Moreover, this is a basis, and conditions (3) are just explicitly spelled out conditions in (8).

One way of proving that the set of vectors (8) is a basis of $L(k\Lambda_0)$ would be by invoking the correct character formula, and this is what we do: The “correct character formula” would be a basis of $M^1(k\Lambda_0)$ parameterized by partitions $\mathcal{P}(\bar{B}_-) \cap ((\mathcal{P}(\bar{B}_-) \cdot \ell(\bar{R})) \cup (\mathcal{P}(\bar{B}_-) \cdot y(0)))$. So we start, as it was indicated earlier, with a spanning set of $M^1(\Lambda)$, $\Lambda = k\Lambda_0$, of the form

$$(9) \quad u(\pi)r(\rho) \cdot v_\Lambda, \quad \pi \in \mathcal{P}(\bar{B}_-), \quad \rho \in \ell(\bar{R}).$$

Each element $u(\pi)r(\rho)$ is a sum of the form

$$(10) \quad u(\pi)r(\rho) = u(\pi \cdot \rho) + \sum_{\kappa \succ \pi \cdot \rho} c_\kappa u(\kappa),$$

and again we say that $\pi \cdot \rho$ is the leading term of $u(\pi)r(\rho)$. Almost half of this paper is devoted to the proof of the following general form of relations among relations:

If leading terms of $u(\pi)r(\rho)$ and $u(\pi_1)r(\rho_1)$ are equal, then $u(\pi)r(\rho) - u(\pi_1)r(\rho_1)$ can be expressed as a linear combination of terms $u(\pi')r(\rho')$, $\pi' \cdot \rho' \succ \pi \cdot \rho$. Having this, by induction we see that the spanning set (9) of $M^1(\Lambda)$ can be reduced to a spanning set of vectors indexed by their leading terms. Since vectors $u(\pi \cdot \rho)v_\Lambda$ are elements of Poincaré-Birkhoff-Witt basis of $M(\Lambda)$, by using (10) we see that vectors $u(\pi)r(\rho)v_\Lambda$ of our reduced spanning set are linearly independent, and hence we have a basis of $M^1(k\Lambda_0)$ indexed in a desired way. Of course, the example $\Lambda = k\Lambda_0$ was taken only to avoid mentioning some technical difficulties in our brief discussion.

As it was pointed out, relations among relations are the central part of our proof of linear independence. This was done in a few steps: we start in Section 8 with generators of the kernel of $\tilde{\mathfrak{g}}$ -module map

$$(11) \quad U(\tilde{\mathfrak{g}}) \otimes_{U(\tilde{\mathfrak{g}}_{\geq 0})} \bar{R}\mathbf{1} \xrightarrow{\Psi_{N(k\Lambda_0)}} N^1(k\Lambda_0) \rightarrow 0,$$

next we formulate relations among relations in terms of colored partitions and leading terms (Lemma 9.2), and finally we prove the general case by induction and by solving some odd dozen of exceptional cases (Sections 9 and 10).

If we are to single out one major point from Sections 9 and 10, it should be Lemmas 9.4 and 10.7. They allow one to try to express $u(\pi)r(\rho) - u(\pi_1)r(\rho_1)$ as a linear combination of terms $u(\pi')r(\rho')$ first, and to worry about $\pi' \cdot \rho' \succ \pi \cdot \rho$ later.

Except for the infinite sums that pervade our theory, the work in Sections 9 and 10 appears to be of the same type as that required for producing a Groebner (or standard) basis of a suitable $\tilde{\mathfrak{g}}$ -module with respect to a total order on the monomials in a basis of $\tilde{\mathfrak{g}}$ (reducing S-pairs etc.). We have not attempted to investigate this connection in detail.

It is clear that combinatorics should enter sooner or later. Some of the combinatorial arguments here are carried through with a brute force, but we still hope that by using more of representation theory, say (11) for example, one can prove relations among relations in some simpler way, having to use combinatorial arguments at some later stage.

Of course, the combinatorial identity (4) is a consequence of (3), in full generality we discuss this in Section 11 by recalling Lepowsky-Wakimoto product formulas for certain specializations of the Weyl-Kac character formula.

It is a pleasure to the second author M.P. to thank the Department of Mathematics of University of Lund for their kind hospitality—a large portion of this work was done during his visit there.

1. FORMAL LAURENT SERIES AND RATIONAL FUNCTIONS

1.1. Power series expansions.

We shall throughout use the calculus of formal Laurent series as in [FLM, Chap. 2 and 8]. In particular $z_1, z_2, \dots$ denote “commuting formal variables”. We sometimes use the expansion convention that, for $n < 0$, $(z_0 + z_1 + \dots + z_k)^n$ denotes the power series expansion in nonnegative powers of $z_1, \dots, z_k$.

For any field F , denote by

$$(1.1.1) \quad F((z)) = \left\{ \sum_{n \in \mathbb{Z}} a_n z^n \mid a_n \in F, \exists N, a_n = 0 \text{ unless } n \geq N \right\}$$

the field of fractions of the formal power series algebra $F[[z]]$. Denote multiindices by

$$\alpha = (\alpha_1, \dots, \alpha_k), \quad |\alpha| = \alpha_1 + \dots + \alpha_k$$

and set

$$z^\alpha = z_1^{\alpha_1} \cdots z_k^{\alpha_k}.$$

Let $F\{z_1, \dots, z_k\}$ denote the vector space

$$F\{z_1, \dots, z_k\} = \left\{ \sum_{\alpha \in \mathbb{Z}^k} a_\alpha z^\alpha \mid a_\alpha \in F \right\}.$$

Let $\mathbb{F}$ be a field of characteristic 0. We have

$$\begin{aligned} \mathbb{F}((z_1)) \cdots ((z_k)) &= \left\{ \sum_{\alpha} a_\alpha z^\alpha \mid a_\alpha \in \mathbb{F}, \exists N_k, \dots, N_1(\alpha_2, \dots, \alpha_k) \text{ s.t. } a_\alpha = 0 \right. \\ &\quad \left. \text{unless } \alpha_k \geq N_k, \dots, \alpha_1 \geq N_1(\alpha_2, \dots, \alpha_k) \right\}, \end{aligned}$$

$$\begin{aligned} \mathbb{F}((z_k^{-1})) \cdots ((z_1^{-1})) &= \left\{ \sum_{\alpha} a_\alpha z^\alpha \mid a_\alpha \in \mathbb{F}, \exists N_1, \dots, N_k(\alpha_1, \dots, \alpha_{k-1}) \text{ s.t. } a_\alpha = 0 \right. \\ &\quad \left. \text{unless } \alpha_1 \leq N_1, \dots, \alpha_k \leq N_k(\alpha_1, \dots, \alpha_{k-1}) \right\}. \end{aligned}$$

There are canonical embeddings

$$\begin{aligned} \mathbb{F}[z_1, \dots, z_k] &= \mathbb{F}[z_1] \cdots [z_k] \rightarrow \mathbb{F}((z_1)) \cdots ((z_k)), \\ \mathbb{F}[z_1, \dots, z_k] &= \mathbb{F}[z_k] \cdots [z_1] \rightarrow \mathbb{F}((z_k^{-1})) \cdots ((z_1^{-1})). \end{aligned}$$

Since the target algebras are fields, the embeddings extend to embeddings

$$\begin{aligned} \iota : \mathbb{F}(z_1, \dots, z_k) &\rightarrow \mathbb{F}((z_1)) \cdots ((z_k)), \\ \iota' : \mathbb{F}(z_1, \dots, z_k) &\rightarrow \mathbb{F}((z_k^{-1})) \cdots ((z_1^{-1})). \end{aligned}$$

Lemma 1.1.1. *Let $(a_1, \dots, a_k) \in \mathbb{F}^k \setminus \{(0, \dots, 0)\}$. Then, in $\mathbb{F}\{z_1, \dots, z_k\}$,*

$$\iota\left(\sum_{i=1}^k a_i z_i\right)^{-1} = \iota'\left(\sum_{i=1}^k a_i z_i\right)^{-1}.$$

Proof. Let $a_i = 0$ for $i < m$, $a_m \neq 0$. Then

$$\begin{aligned} \iota'\left(\sum_{i=1}^k a_i z_i\right)^{-1} &= \iota'(a_m z_m (1 + \sum_{i=m+1}^k \frac{a_i z_i}{a_m z_m}))^{-1} \\ &= a_m^{-1} z_m^{-1} \sum_{n \geq 0} (-1)^n \left(\sum_{i=m+1}^k \frac{a_i z_i}{a_m z_m} \right)^n \in \mathbb{F}((z_1)) \cdots ((z_k)). \end{aligned}$$

Hence

$$\begin{aligned} \iota'\left(\sum_{i=1}^k a_i z_i\right)^{-1} &= \iota'\left(\sum_{i=1}^k a_i z_i\right)^{-1} \left(\sum_{i=1}^k a_i z_i \iota\left(\sum_{i=1}^k a_i z_i\right)^{-1}\right) \\ &= \left(\iota'\left(\sum_{i=1}^k a_i z_i\right)^{-1} \sum_{i=1}^k a_i z_i\right) \iota\left(\sum_{i=1}^k a_i z_i\right)^{-1} = \iota\left(\sum_{i=1}^k a_i z_i\right)^{-1} \end{aligned}$$

by the associative law in $\mathbb{F}((z_1)) \cdots ((z_k))$. $\square$

Set

$$(1.1.2) \quad S = \left\{ \sum_{i=1}^k a_i z_i \mid (a_1, \dots, a_k) \neq (0, \dots, 0) \right\},$$

$$(1.1.3) \quad \hat{\mathcal{R}}_k = \mathbb{F}[z_1, \dots, z_k]_{\langle S \rangle},$$

the localization.

Corollary 1.1.2. *The homomorphisms ι, ι' coincide on $\hat{\mathcal{R}}_k$.*

Remark. We have

$$\iota(1 - z_1)^{-1} = \sum_{n \geq 0} z_1^n \neq \iota'(1 - z_1)^{-1} = - \sum_{n \geq 0} z_1^{-n-1}$$

although $\iota(z_1 - z_2 z_3)^{-1} = \iota'(z_1 - z_2 z_3)^{-1}$. It seems difficult to determine for which $f \in \mathbb{F}(z_1, \dots, z_k)$, $\iota(f) = \iota'(f)$.

Define

$$(1.1.4) \quad \iota_{1, \dots, k} : \hat{\mathcal{R}}_k \rightarrow \mathbb{F}((z_1)) \cdots ((z_k))$$

to be $\iota_{1, \dots, k} = \iota = \iota'$. Also define the partial ι -maps

$$(1.1.5) \quad \iota_{\dots \ell+1, \dots, k} : \hat{\mathcal{R}}_k \rightarrow \hat{\mathcal{R}}_\ell((z_{\ell+1})) \cdots ((z_k));$$

the restriction of

$$\mathbb{F}(z_1, \dots, z_k) \rightarrow \mathbb{F}(z_1, \dots, z_\ell)((z_{\ell+1})) \cdots ((z_k)).$$

By considering the image of $(\sum_{i=1}^k a_i z_i)^{-1}$ one verifies that the image of $\hat{\mathcal{R}}_k$ actually lands in

$$\hat{\mathcal{R}}_\ell((z_{\ell+1})) \cdots ((z_k)).$$

For $m, n \in \mathbb{N}$, $m \leq n$, denote by $\hat{\mathcal{R}}_{[m, n]}$ the algebra analogous to $\hat{\mathcal{R}}_k$, based on the variables z_i , $m \leq i \leq n$. Define also

$$(1.1.6) \quad \iota_{1, \dots, \ell \dots} : \hat{\mathcal{R}}_k \rightarrow \hat{\mathcal{R}}_{[\ell+1, k]}((z_\ell^{-1})) \cdots ((z_1^{-1})),$$

the restriction of

$$\mathbb{F}(z_1, \dots, z_k) \rightarrow \mathbb{F}(z_{\ell+1}, \dots, z_k)((z_\ell^{-1})) \cdots ((z_1^{-1}));$$

here it is easier to see that the image is in $\hat{\mathcal{R}}_{[\ell+1, k]} \cdots$.

Proposition 1.1.3. *We have*

$$\iota_{1,\dots,k} = \iota_{1,\dots,\ell} \circ \iota_{\dots\ell+1,\dots,k} = \iota_{\ell+1,\dots,k} \circ \iota_{1,\dots,\ell\dots} : \hat{\mathcal{R}}_k \rightarrow \mathbb{F}((z_1)) \cdots ((z_k)).$$

Proof. The composition $\iota_{1,\dots,\ell} \circ \iota_{\dots\ell+1,\dots,k}$ is the restriction to $\hat{\mathcal{R}}_k$ of the canonical map

$$\iota : \mathbb{F}(z_1, \dots, z_k) \rightarrow \mathbb{F}((z_1)) \cdots ((z_k))$$

i.e. $\iota_{1,\dots,k}$. Also $\iota_{\ell+1,\dots,k} = \iota = \iota'_{[\ell+1,k]}$ on $\hat{\mathcal{R}}_{[\ell+1,k]}$ so that the composition $\iota_{\ell+1,\dots,k} \circ \iota_{1,\dots,\ell\dots}$ is the restriction to $\hat{\mathcal{R}}_k$ of the canonical map

$$\iota' : \mathbb{F}(z_1, \dots, z_k) \rightarrow \mathbb{F}((z_k^{-1})) \cdots ((z_1^{-1})).$$

By the corollary $\iota = \iota'$ on $\hat{\mathcal{R}}_k$. $\square$

Remark. It follows from the formula $\iota_{1,\dots,k} = \iota_{2,\dots,k} \circ \iota_{1,\dots}$ that the present definition of $\iota_{1,\dots,k}$ coincides with the one in [FLM, (8.10.37–42)].

Let $\sigma \in S_k = \text{Sym}\{1, \dots, k\}$. Denote also by σ the automorphism

$$\begin{aligned} \sigma : \mathbb{F}[z_1, \dots, z_k] &\rightarrow \mathbb{F}[z_1, \dots, z_k] \\ z_m &\mapsto z_{\sigma m}. \end{aligned}$$

Then $\sigma(S) = S$, hence σ extends to an automorphism

$$\sigma : \hat{\mathcal{R}}_k \rightarrow \hat{\mathcal{R}}_k.$$

Let σ also denote the vector space isomorphism

$$\begin{aligned} \sigma : \mathbb{F}\{z_1, \dots, z_k\} &\rightarrow \mathbb{F}\{z_1, \dots, z_k\} \\ \sum_{\alpha} a_{\alpha} z^{\alpha} &\mapsto \sum_{\alpha} a_{\alpha} z_{\sigma 1}^{\alpha_1} \cdots z_{\sigma k}^{\alpha_k} = \sum_{\alpha} a_{\alpha} z^{\alpha \circ \sigma^{-1}}. \end{aligned}$$

Then σ has the homomorphism property; if

$$x, y \in \mathbb{F}\{z_1, \dots, z_k\}$$

and xy is defined, then $\sigma(x)\sigma(y)$ is also defined and

$$\sigma(xy) = \sigma(x)\sigma(y).$$

Define also

$$(1.1.7) \quad \iota_{\sigma 1, \dots, \sigma k} = \sigma \circ \iota_{1, \dots, k} \circ \sigma^{-1},$$

$$(1.1.8) \quad \iota_{\dots\sigma(\ell+1), \dots, \sigma(k)} = \sigma \circ \iota_{\dots\ell+1, \dots, k} \circ \sigma^{-1},$$

$$(1.1.9) \quad \iota_{\sigma(1), \dots, \sigma(\ell)\dots} = \sigma \circ \iota_{1, \dots, \ell\dots} \circ \sigma^{-1}.$$

Then we have more generally

$$(1.1.10) \quad \begin{aligned} \iota_{\sigma 1, \dots, \sigma k} &= \iota_{\sigma 1, \dots, \sigma \ell} \circ \iota_{\dots\sigma(\ell+1), \dots, \sigma(k)} = \iota_{\sigma(\ell+1), \dots, \sigma(k)} \circ \iota_{\sigma(1), \dots, \sigma(\ell)\dots} \\ &: \hat{\mathcal{R}}_k \rightarrow \mathbb{F}((z_{\sigma 1})) \cdots ((z_{\sigma k})). \end{aligned}$$

1.2. Cauchy's theorem.

Set

$$(1.2.1) \quad \delta(z) = \sum_{n \in \mathbb{Z}} z^n.$$

For $k \in \mathbb{N}$ set

$$(1.2.2) \quad \mathcal{R}_k = \mathbb{F}[z_i, z_i^{-1}, (z_m - z_n)^{-1}; 1 \leq i \leq k, 1 \leq m < n \leq k],$$

$$(1.2.3) \quad \mathcal{L}_k = \mathbb{F}[z_i, z_i^{-1}; 1 \leq i \leq k].$$

The following Proposition was first formulated by H. Garland, cf. [FLM].

Proposition 1.2.1. *We have*

$$\delta(z)p(z) = \delta(z)p(1),$$

$$\delta(az_1/z_2)q(z_1, z_2) = \delta(az_1/z_2)q(z_1, az_1)$$

for all $p \in \mathbb{F}[z, z^{-1}]$, $a \in \mathbb{F}$, and $q \in \mathcal{L}_2$.

Proposition 1.2.2. *Let $g(z_0, z_1, z_2) \in \mathcal{L}_3$. Then*

$$\text{Res}_{z_0} z_0^{-1} \delta\left(\frac{z_1 - z_2}{z_0}\right) g(z_0, z_1, z_2) = \iota_{12} g(z_1 - z_2, z_1, z_2).$$

Proposition 1.2.3. *Let $f(z_1, z_2) \in \mathcal{R}_2$. Then*

$$\text{Res}_{z_1, z_2} \iota_{12} f - \text{Res}_{z_1, z_2} \iota_{21} f = \text{Res}_{z_0, z_2} \iota_{20} f(z_2 + z_0, z_2).$$

Proof. By partial fraction decomposition one may assume that

$$f \in \{z_1^m z_2^n, z_2^m (z_1 - z_2)^n \mid m, n \in \mathbb{Z}\}.$$

If $m + n \neq -2$ we have $0 - 0 = 0$. If $f = \frac{\partial}{\partial z_1} g$ then all terms vanish as well. Let $f = z_1^{-1} z_2^{-1}$. Then

$$\text{Res}_{z_1, z_2} \iota_{12} f - \text{Res}_{z_1, z_2} \iota_{21} f = 0,$$

$$\text{Res}_{z_0, z_2} \iota_{20} f(z_2 + z_0, z_2) = \text{Res}_{z_0, z_2} \iota_{20} (z_2 + z_0)^{-1} z_2^{-1} = 0.$$

Let $f = z_2^{-1} (z_1 - z_2)^{-1}$. Then

$$\text{Res}_{z_1, z_2} \iota_{12} f = \text{Res}_{z_1, z_2} z_2^{-1} \sum_{i \geq 0} \binom{-1}{i} z_1^{-1-i} (-z_2)^i = 1,$$

$$\text{Res}_{z_1, z_2} \iota_{21} f = \text{Res}_{z_1, z_2} z_2^{-1} \sum_{i \geq 0} \binom{-1}{i} (-z_2)^{-1-i} z_1^i = 0,$$

$$\text{Res}_{z_0, z_2} \iota_{20} f(z_2 + z_0, z_2) = \text{Res}_{z_0, z_2} z_2^{-1} z_0^{-1} = 1. \quad \square$$

Remark. Proposition 1.2.3 follows alternatively from Cauchy's theorem: Take real numbers ϵ, r, ρ, R , such that $0 < r < \rho < R, 0 < \epsilon < \min\{R - \rho, \rho - r\}$. Then (if f has coefficients in $\mathbb{C}$)

$$\begin{aligned}
& \text{Res}_{z_1, z_2} \iota_{12} f - \text{Res}_{z_1, z_2} \iota_{21} f \\
&= \frac{1}{(2\pi i)^2} \int_{C_R} \left(\int_{C_\rho} f(z_1, z_2) dz_2 \right) dz_1 - \frac{1}{(2\pi i)^2} \int_{C_\rho} \left(\int_{C_r} f(z_1, z_2) dz_1 \right) dz_2 \\
(1.2.4) \quad &= \frac{1}{(2\pi i)^2} \int_{C_\rho} \left(\int_{C_\epsilon(z_2)} f(z_1, z_2) dz_1 \right) dz_2 \\
&= \frac{1}{(2\pi i)^2} \int_{C_\rho} \left(\int_{C_\epsilon} f(z_2 + z_0, z_2) dz_0 \right) dz_2 \\
&= \text{Res}_{z_0, z_2} \iota_{20} f(z_2 + z_0, z_2).
\end{aligned}$$

An equivalent formulation of this result is:

Proposition 1.2.4. *Let $f(z_1, z_2) \in \mathcal{R}_2$. Then*

$$\text{Res}_{z_1, z_2} \iota_{12} f = \text{Res}_{z_1, z_2} (\iota_{21} - \iota_{12}) f(z_1, z_1 + z_2).$$

Lemma 1.2.5. *For all $f(z_1, z_2) \in \mathcal{R}_2$,*

$$T\iota_{34} f(z_4, z_3) = \text{Res}_{z_4} \iota_{34} z_2^{-1} \delta\left(\frac{z_3 - z_4}{-z_2}\right) f(z_4, z_3) = (\iota_{23} - \iota_{32}) f(z_2 + z_3, z_3).$$

Proof. It is enough to check the coefficients of $z_2^{-m-1} z_3^{-n-1}$ on both sides. Let $g(z_2, z_3) = z_2^m z_3^n \in \mathcal{L}_2$. Then

$$\begin{aligned}
& \text{Res}_{z_2, z_3} \text{Res}_{z_4} \iota_{34} z_2^{-1} \delta\left(\frac{z_3 - z_4}{-z_2}\right) f(z_4, z_3) g(z_2, z_3) \\
&= \text{Res}_{z_3, z_4} \iota_{34} f(z_4, z_3) g(-z_3 + z_4, z_3) \\
&= \text{Res}_{z_3, z_4} (\iota_{43} - \iota_{34}) f(z_3 + z_4, z_3) g(z_4, z_3) \\
&= \text{Res}_{z_2, z_3} (\iota_{23} - \iota_{32}) f(z_2 + z_3, z_3) g(z_2, z_3). \quad \square
\end{aligned}$$

2. GENERATING FIELDS

Definition 2.1. [B], [FLM], [FHL] A *vertex operator algebra* $(V, Y, \mathbf{1}, \omega)$ consists of a vector space V over $\mathbb{F}$, distinguished vectors $\mathbf{1}, \omega \in V$, and a linear map

$$\begin{aligned}
Y : V &\rightarrow (\text{End}_{\mathbb{F}} V)[[z, z^{-1}]], \\
v &\mapsto Y(v, z) = \sum_{n \in \mathbb{Z}} v_n z^{-n-1}
\end{aligned}$$

satisfying the following axioms:

$$\begin{aligned}
(2.1) \quad & \text{For all } u, v \in V \text{ there is an } N = N(u, v) \in \mathbb{Z} \text{ such that} \\
& u_n v = 0 \text{ for all } n > N,
\end{aligned}$$

and

$$(2.2) \quad v_{-1}\mathbf{1} = v, \quad v_n\mathbf{1} = 0$$

for $v \in V, n \geq 0$. Define L_n by

$$(2.3) \quad Y(\omega, z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2}$$

(i.e. $L_n = \omega_{n+1}$). Then

$$(2.4) \quad [L_{-1}, Y(v, z)] = \frac{d}{dz} Y(v, z) = Y(L_{-1}v, z)$$

for all $v \in V$, and

$$(2.5) \quad [L_m, L_n] = (m - n)L_{m+n} + \frac{m^3 - m}{12} \delta_{m+n,0} (\text{rank } V) 1_V$$

for some scalar $\text{rank } V \in \mathbb{F}$. Set, for $n \in \mathbb{Z}$,

$$(2.6) \quad V_n = \{v \in V \mid L_0 v = nv\}.$$

Then

$$(2.7) \quad V = \coprod_{n \in \mathbb{Z}} V_n,$$

$$(2.8) \quad \dim_{\mathbb{F}} V_n < \infty \text{ for all } n,$$

$$(2.9) \quad V_n = 0 \text{ for } n \text{ sufficiently small.}$$

Finally,

$$(2.10) \quad \begin{aligned} & z_0^{-1} \delta\left(\frac{z_1 - z_2}{z_0}\right) Y(u, z_1) Y(v, z_2) - z_0^{-1} \delta\left(\frac{z_2 - z_1}{-z_0}\right) Y(v, z_2) Y(u, z_1) \\ &= z_2^{-1} \delta\left(\frac{z_1 - z_0}{z_2}\right) Y(Y(u, z_0)v, z_2) \end{aligned}$$

for all $u, v \in V$ (the Jacobi identity).

We say that $Y(v, z)$ is a *vertex operator* or a *field* associated with a vector v . Later on we shall consider a formal Laurent series $Y(v, z)$ as a generating function for the coefficients $v_n, n \in \mathbb{Z}$.

In later sections we shall use the following consequences of the axioms [B], [FLM], [FHL]: For all $u, v \in V, n \in \mathbb{Z}$,

$$(2.11) \quad \begin{aligned} Y(u_n v, z) &= \text{Res}_{z_0} z_0^n (Y(u, z_0 + z) Y(v, z) \\ &\quad - Y(v, z) \text{Res}_{z_1} z_0^{-1} \delta\left(\frac{z - z_1}{-z_0}\right) Y(u, z_1)), \end{aligned}$$

with the notation

$$(2.12) \quad Y(v, z)^- = \sum_{n < 0} v_n z^{-n-1}, \quad Y(v, z)^+ = \sum_{n \geq 0} v_n z^{-n-1},$$

we have

$$(2.13) \quad \begin{aligned} Y(u_n v, z) &= \left(\frac{(\partial/\partial z)^{-n-1}}{(-n-1)!} Y(u, z)^- \right) Y(v, z) \\ &\quad + Y(v, z) \left(\frac{(\partial/\partial z)^{-n-1}}{(-n-1)!} Y(u, z)^+ \right) \end{aligned}$$

for $n \leq -1$, in particular

$$(2.14) \quad Y(u_{-1} v, z) = Y(u, z)^- Y(v, z) + Y(v, z) Y(u, z)^+,$$

furthermore

$$(2.15) \quad [Y(u, z_1), Y(v, z_2)] = \sum_{i=0}^{\infty} \frac{(-1)^i}{i!} \left(\frac{\partial}{\partial z_1} \right)^i z_2^{-1} \delta(z_1/z_2) Y(u_i v, z_2),$$

and

$$(2.16) \quad Y(u, z)v = e^{zL_{-1}} Y(v, -z)u.$$

The goal of this section is to derive Theorem 2.6 below giving a kind of “generators and relations” construction of vertex operator algebras.

Let V be a vector space over $\mathbb{F}$, and let $L_{-1}, L_0 \in \text{End}_{\mathbb{F}} V$, $\mathbf{1} \in V$. Set, for $n \in \mathbb{Z}$,

$$V_n = \{v \in V \mid L_0 v = nv\}.$$

Assume that

$$(2.17) \quad V = \coprod_{n \in \mathbb{Z}} V_n,$$

$$(2.18) \quad \mathbf{1} \in V_0, \quad \text{i.e. } L_0 \mathbf{1} = 0,$$

$$(2.19) \quad [L_0, L_{-1}] = L_{-1}, \quad \text{i.e. } L_{-1} V_n \subset V_{n+1}.$$

Definition 2.2. A generating function

$$a(z) = \sum_{n \in \mathbb{Z}} a_n z^{-n-1}$$

of operators $a_n \in \text{End}_{\mathbb{F}} V$ is said to be *admissible of conformal weight h* if

$$(2.20) \quad \text{for all } v \in V \text{ there is } N(a, v) \in \mathbb{Z} \text{ s.t. } a_n v = 0 \text{ if } n > N(a, v),$$

$$(2.21) \quad [L_{-1}, a(z)] = \frac{d}{dz} a(z),$$

$$(2.22) \quad [L_0, a(z)] = z \frac{d}{dz} a(z) + ha(z),$$

$$(2.23) \quad a(z) \mathbf{1} \in V[[z]], \quad \text{i.e. } a_n \mathbf{1} = 0 \text{ if } n \geq 0.$$

A generating function is *admissible* if it is a (finite) linear combination of admissible generating functions of operators with (various) conformal weights.

It follows from (2.18) and (2.22) that

$$(2.24) \quad a_{-1} \mathbf{1} \in V_h, \quad \text{i.e. } L_0 a_{-1} \mathbf{1} = h a_{-1} \mathbf{1}.$$

Proposition 2.3. *Let $a(z), b(z)$ be admissible generating functions. Define $d_n(z)$ for $n \in \mathbb{Z}$ by*

$$(2.25) \quad \sum_{n \in \mathbb{Z}} d_n(z_3) z_2^{-n-1} = a(z_2 + z_3) b(z_3) - b(z_3) T a(z_4),$$

where T denotes the operator

$$(2.26) \quad T = \text{Res}_{z_4} z_2^{-1} \delta\left(\frac{z_3 - z_4}{-z_2}\right)(\cdot)$$

(cf. [FLM, (8.8.31)]). Then $d_n(z)$ is admissible for all $n \in \mathbb{Z}$.

Proof. (2.20): Let $v \in V$. Then

$$(2.27) \quad \begin{aligned} (d_n)_k v &= \text{coeff}_{z_2^{-n-1} z_3^{-k-1}} \{a(z_2 + z_3) b(z_3) - b(z_3) [a(z_2 + z_3) - a(z_3 + z_2)]\} v \\ &= \sum_{r \leq n} \binom{-r-1}{n-r} a_r b_{k+n-r} v - \sum_{0 \leq r \leq n} \binom{-r-1}{n-r} b_{k+n-r} a_r v \\ &\quad + \sum_{r \geq 0} \binom{-r-1}{-n-1} b_{k+n-r} a_r v. \end{aligned}$$

There are $N_1, N_2 \in \mathbb{Z}$ such that $N_1 \geq 0$, $a_m v = 0$ if $m \geq N_1$, $b_m v = 0$ and $b_m a_r v = 0$ if $m \geq N_2$ and $0 \leq r \leq N_1 - 1$. If $N = \max\{-n, 0\} + N_1 + N_2$ we then have $(d_n)_k v = 0$ if $k \geq N$.

(2.21): It follows from

$$T a(z_4) = \sum_{j \in \mathbb{Z}} a_j \{(z_2 + z_3)^{-j-1} - (z_3 + z_2)^{-j-1}\}$$

that

$$[L_{-1}, T a(z_4)] = \frac{\partial}{\partial z_3} T a(z_4).$$

Hence

$$\begin{aligned} [L_{-1}, \sum_{n \in \mathbb{Z}} d_n(z_3) z_2^{-n-1}] &= [L_{-1}, a(z_2 + z_3)] b(z_3) + a(z_2 + z_3) [L_{-1}, b(z_3)] \\ &\quad - [L_{-1}, b(z_3)] T a(z_4) - b(z_3) [L_{-1}, T a(z_4)] \\ &= \frac{\partial}{\partial z_3} (\sum_{n \in \mathbb{Z}} d_n(z_3) z_2^{-n-1}). \end{aligned}$$

(2.22): We may assume that a and b have conformal weights h_a, h_b . Then

$$\begin{aligned} [L_0, \sum_{n \in \mathbb{Z}} d_n(z_3) z_2^{-n-1}] &= [L_0, a(z_2 + z_3)] b(z_3) + a(z_2 + z_3) [L_0, b(z_3)] \\ &\quad - [L_0, b(z_3)] T a(z_4) - b(z_3) [L_0, T a(z_4)] \\ &= (z_2 \frac{\partial}{\partial z_2} + z_3 \frac{\partial}{\partial z_3} + h_a + h_b) (\sum_{n \in \mathbb{Z}} d_n(z_3) z_2^{-n-1}) \\ &= \sum_{n \in \mathbb{Z}} \{(z_3 \frac{\partial}{\partial z_3} + h_a + h_b - n - 1) d_n(z_3)\} z_2^{-n-1} \end{aligned}$$

so that d_n has conformal weight $h_a + h_b - n - 1$.

(2.23): This follows from (2.27) and the fact that $a_k \mathbf{1} = 0 = b_k \mathbf{1}$ if $k \geq 0$. $\square$

Set

$$V' = \coprod_{n \in \mathbb{Z}} V_n^*,$$

where $V_n^* = \text{Hom}_{\mathbb{F}}(V_n, \mathbb{F})$ and denote the pairing between V' and V by

$$\langle \cdot, \cdot \rangle : V' \times V \rightarrow \mathbb{F}.$$

Definition 2.4. Let $a(z), b(z)$ be admissible generating functions. We say that $a(z), b(z)$ are mutually *local* or *commutative* if there exists $N \in \mathbb{N}$ such that for all $w \in V, w' \in V'$,

$$(z_1 - z_2)^N \langle w', a(z_1)b(z_2)w \rangle = (z_1 - z_2)^N \langle w', b(z_2)a(z_1)w \rangle$$

i.e. $(z_1 - z_2)^N [a(z_1), b(z_2)] = 0$.

Equivalently, $a(z)$ and $b(z)$ are mutually local iff there is $f(z_1, z_2) \in \mathcal{R}_2$ such that

$$\begin{aligned} \langle w', a(z_1)b(z_2)w \rangle &= \iota_{12}f, \\ \langle w', b(z_2)a(z_1)w \rangle &= \iota_{21}f, \end{aligned}$$

with ι as in Section 1.1, and there is an upper bound that is independent of w and w' for the order of the pole $(z_1 - z_2)^{-1}$ in such matrix coefficients.

Proposition 2.5. Let $a(z), b(z), c(z)$ be pairwise local admissible generating functions. Let $d_n(z)$ for $n \in \mathbb{Z}$ be defined as in (2.25), (2.26). Then $d_n(z)$ and $c(z)$ are mutually local for each $n \in \mathbb{Z}$.

Proof. Set

$$d(z_2, z_3) = a(z_2 + z_3)b(z_3) - b(z_3)Ta(z_4) \in \text{End}_{\mathbb{F}} V\{z_2, z_3\}.$$

Let $w \in V, w' \in V'$. We shall calculate the matrix coefficients $\langle w', c(z_1)d(z_2, z_3)w \rangle$ and $\langle w', d(z_2, z_3)c(z_1)w \rangle$. It follows from (2.17), (2.20), (2.22) that there is

$$f(z_1, z_2, z_3) \in \mathcal{R}_3$$

such that

$$\begin{aligned} \langle w', c(z_1)a(z_2)b(z_3)w \rangle &= \iota_{123}f, \\ \langle w', c(z_1)b(z_3)a(z_2)w \rangle &= \iota_{132}f. \end{aligned}$$

Then

$$\langle w', c(z_1)a(z_2 + z_3)b(z_3)w \rangle = \iota_{123}f(z_1, z_2 + z_3, z_3).$$

Let $\iota_{1\dots}f = \sum_{n \in \mathbb{Z}} g_n(z_2, z_3)z_1^n$. Using Lemma 1.2.5 and Proposition 1.1.3 we obtain

$$\begin{aligned} \langle w', c(z_1)b(z_3)Ta(z_4)w \rangle &= T\iota_{134}f(z_1, z_4, z_3) \\ &= \sum_{n \in \mathbb{Z}} T\iota_{34}g_n(z_4, z_3)z_1^n \\ &= \sum_{n \in \mathbb{Z}} (\iota_{23} - \iota_{32})g_n(z_2 + z_3, z_3)z_1^n \\ &= (\iota_{123} - \iota_{132})f(z_1, z_2 + z_3, z_3). \end{aligned}$$

Hence

$$(2.28) \quad \langle w', c(z_1)d(z_2, z_3)w \rangle = \iota_{132}f(z_1, z_2 + z_3, z_3).$$

Similarly,

$$\langle w', a(z_2 + z_3)b(z_3)c(z_1)w \rangle = \iota_{231}f(z_1, z_2 + z_3, z_3),$$

and

$$\begin{aligned} \langle w', b(z_3)Ta(z_4)c(z_1)w \rangle &= T\iota_{341}f(z_1, z_4, z_3) \\ &= \sum_{n \in \mathbb{Z}} T\iota_{34}h_n(z_4, z_3)z_1^n \\ &= \sum_{n \in \mathbb{Z}} (\iota_{23} - \iota_{32})h_n(z_2 + z_3, z_3)z_1^n \\ &= (\iota_{231} - \iota_{321})f(z_1, z_2 + z_3, z_3) \end{aligned}$$

if $\iota_{...1}f = \sum_{n \in \mathbb{Z}} h_n(z_2, z_3)z_1^n$. This gives

$$\langle w', d(z_2, z_3)c(z_1)w \rangle = \iota_{321}f(z_1, z_2 + z_3, z_3).$$

Since

$$f(z_1, z_2 + z_3, z_3) \in \mathbb{F}[z_1, z_2, z_3, z_1^{-1}, z_2^{-1}, z_3^{-1}, (z_1 - z_2 - z_3)^{-1}, (z_1 - z_3)^{-1}, (z_2 + z_3)^{-1}],$$

we have $\iota_{321}f(z_1, z_2 + z_3, z_3) = \iota_{312}f(z_1, z_2 + z_3, z_3)$ so that

$$(2.29) \quad \langle w', d(z_2, z_3)c(z_1)w \rangle = \iota_{312}f(z_1, z_2 + z_3, z_3).$$

Define $f_n(z_1, z_3) \in \mathcal{R}_{\{1,3\}}$ by

$$\iota_{...2}f(z_1, z_2 + z_3, z_3) = \sum_{n \in \mathbb{Z}} f_n(z_1, z_3)z_2^{-n-1}.$$

Then (2.28) and (2.29) imply

$$\begin{aligned} \langle w', c(z_1)d_n(z_3)w \rangle &= \iota_{13}f_n, \\ \langle w', d_n(z_3)c(z_1)w \rangle &= \iota_{31}f_n. \end{aligned}$$

Let

$$f(z_1, z_2, z_3) = \frac{p(z_1, z_2, z_3)}{(z_1 - z_2)^K (z_1 - z_3)^L (z_2 - z_3)^M}$$

with $p \in \mathcal{L}_3$. Then

$$\begin{aligned} \iota_{...2}f(z_1, z_2 + z_3, z_3) &= \iota_{...2} \frac{p(z_1, z_2 + z_3, z_3)}{(z_1 - z_2 - z_3)^K (z_1 - z_3)^L z_2^M} \\ &= \left(\sum_{j \geq 0} \binom{-K}{j} (z_1 - z_3)^{-K-j} (-z_2)^j \right) (z_1 - z_3)^{-L} \iota_{...2} \frac{p(z_1, z_2 + z_3, z_3)}{z_2^M}. \end{aligned}$$

Since $\iota_{...2}p(z_1, z_2 + z_3, z_3)$ only involves nonnegative powers of z_2 , the j -th term contributes to f_n only if $j - M \leq -n - 1$. Hence the order of the pole $(z_1 - z_3)^{-1}$ in f_n is at most $K + L + M - n - 1$, and since K, L, M can be taken to be independent of w, w' , we are done. $\square$

We shall now assume that

$$(2.30) \quad L_{-1}\mathbf{1} = 0.$$

The proof of the following theorem uses the ideas of P. Goddard in [G]. A similar result appears in [Xu, Theorem 2.4].

Theorem 2.6. *Let U be a subspace of V . Assume that there is given a linear map*

$$Y : U \rightarrow (\text{End}_{\mathbb{F}} V)[[z, z^{-1}]]$$

$$u \mapsto Y(u, z) = \sum_{n \in \mathbb{Z}} u_n z^{-n-1}$$

such that

$$(2.31) \quad Y(u, z) \text{ is admissible for all } u \in U,$$

$$(2.32) \quad u_{-1} \mathbf{1} = u \text{ for all } u \in U,$$

$$(2.33) \quad Y(u, z) \text{ and } Y(v, z) \text{ are local for each pair } u, v \in U,$$

$$(2.34) \quad V = \mathbb{F}\text{-span}\{u_{n_1}^{(1)} \cdots u_{n_k}^{(k)} \mathbf{1} \mid k \in \mathbb{N}, u^{(i)} \in U, n_i \in \mathbb{Z}\}.$$

Then Y extends uniquely to a linear map

$$Y : V \rightarrow (\text{End}_{\mathbb{F}} V)[[z, z^{-1}]]$$

that makes V into a vertex operator algebra (except that we have only L_0, L_{-1} out of the Virasoro algebra).

Proof. Let W be the space of all admissible generating functions

$$a(z) \in (\text{End}_{\mathbb{F}} V)[[z, z^{-1}]]$$

such that

$$(2.35) \quad a(z) \text{ and } Y(u, z) \text{ are local for each } u \in U.$$

Define

$$\phi : W \rightarrow V$$

$$a(z) \mapsto a_{-1} \mathbf{1}.$$

Step 1. (uniqueness) ϕ is injective.

Proof. Assume that $a(z) \in W$, $a_{-1} \mathbf{1} = 0$. It follows from (2.21) and (2.30) that

$$(2.36) \quad a(z) \mathbf{1} = e^{zL_{-1}} a_{-1} \mathbf{1} = 0.$$

Let $X = \{v \in V \mid a(z)v = 0\}$ and let $v \in X$, $u \in U$. Then there exist $N \in \mathbb{N}$ such that

$$(z_1 - z_2)^N \langle w', a(z_1)Y(u, z_2)v \rangle = (z_1 - z_2)^N \langle w', Y(u, z_2)a(z_1)v \rangle = 0$$

for all $w' \in V'$, hence $a(z_1)u_nv = 0$ and $u_n X \subset X$. Since by (2.36) $\mathbf{1} \in X$ it follows from (2.34) that $X = V$ so that $a(z) = 0$ and ϕ is injective.

By (2.31), (2.33), $Y(u, z) \in W$ for all $u \in U$, hence $\phi(W) \supset U$ by (2.32). Set

$$X = \phi(W), \quad U \subset X \subset V.$$

Since ϕ is injective we can define

$$Y : X \rightarrow (\text{End}_{\mathbb{F}} V)[[z, z^{-1}]]$$

by

$$Y(v, z) = \phi^{-1}(v)$$

for $v \in X$. By (2.32) the two meanings of $Y(u, z)$ for $u \in U$ denote the same series.

Step 2. $X = \phi(W) = V$.

Proof. For $u \in U$, $v \in X$, $n \in \mathbb{Z}$ define $Y(u_n v, z)$ by

$$\sum_{n \in \mathbb{Z}} Y(u_n v, z_3) z_2^{-n-1} = Y(u, z_2 + z_3) Y(v, z_3) - Y(v, z_3) T Y(u, z_4).$$

Then $Y(u_n v, z) \in W$ by Propositions 2.3 and 2.5, and we have

$$\phi Y(u_n v, z) = (u_n v)_{-1} \mathbf{1} = u_n v_{-1} \mathbf{1} = u_n v$$

by (2.27). Hence $u \in U$, $v \in X$, $n \in \mathbb{Z}$ implies

$$(2.37) \quad u_n v \in X.$$

Since $\text{id}_V z^0 \in W$ and $\text{coeff}_{z^0}(\text{id}_V z^0) \mathbf{1} = \mathbf{1}$ we have

$$(2.38) \quad \mathbf{1} \in X,$$

and

$$Y(\mathbf{1}, z) = \text{id}_V z^0.$$

Now (2.34), (2.37), (2.38) imply $X = V$, i.e. we have well defined vertex operators $Y(v, z)$ for all $v \in V$.

Step 3. $Y(u, z)$ and $Y(v, z)$ are mutually local for all pairs $u, v \in V$.

Proof. Fix $v^{(0)} \in V$ and set

$$X = \{v \in V \mid Y(v, z) \text{ and } Y(v^{(0)}, z) \text{ are mutually local}\}.$$

Clearly $\{\mathbf{1}\} \cup U \subset X$. If $u \in U$, $v \in X$, $n \in \mathbb{Z}$ then

$$Y(u_n v, z_3) = \text{coeff}_{z_2^{-n-1}}(Y(u, z_2 + z_3) Y(v, z_3) - Y(v, z_3) T Y(u, z_4))$$

together with Proposition 2.5 show that $u_n v \in X$. Hence (2.34) again implies $X = V$ as required.

We have showed that V satisfies the conditions of the axiomatic characterization of vertex operator algebras [FLM, A.2.5, A.2.8, A.3.1; FHL] (except that we only have L_0, L_{-1} out of the Virasoro algebra). $\square$

3. THE VERTEX OPERATOR ALGEBRA $N(k\Lambda_0)$

Let $\mathfrak{g}$ be a finite dimensional split simple Lie algebra over $\mathbb{F}$, and let $(\cdot, \cdot) : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{F}$ be the nondegenerate invariant symmetric bilinear form normalized so that $(\alpha, \alpha) = 2$ for each long root α . We then have the associated affine Kac-Moody algebra

$$(3.1) \quad \begin{aligned} \hat{\mathfrak{g}} &= \mathfrak{g} \otimes \mathbb{F}[t, t^{-1}] \oplus \mathbb{F}c, & \tilde{\mathfrak{g}} &= \hat{\mathfrak{g}} \oplus \mathbb{F}L_0, \\ c &\text{ central, } & c &\neq 0, & [L_0, x \otimes t^m] &= -m x \otimes t^m, \\ [x \otimes t^m, y \otimes t^n] &= [x, y] \otimes t^{m+n} + m\delta_{m+n,0}(x, y)c. \end{aligned}$$

Note that $d = -L_0$ defines the usual homogeneous grading on $\hat{\mathfrak{g}}$. We shall frequently denote $x \otimes t^m$ by $x(m)$ and $\mathfrak{g} \otimes t^m$ by $\mathfrak{g}(m)$. We often identify $\mathfrak{g}(0)$ and $\mathfrak{g}$. Set

$$(3.2) \quad \mathfrak{p} = \coprod_{n \geq 0} \mathfrak{g} \otimes t^n \oplus \mathbb{F}c, \quad \tilde{\mathfrak{g}}_{\geq 0} = \mathfrak{p} \oplus \mathbb{F}L_0.$$

For $k \in \mathbb{F}$ denote by $\mathbb{F}v_{k\Lambda_0}$ the 1-dimensional $\tilde{\mathfrak{g}}_{\geq 0}$ -module such that

$$(3.3) \quad c \cdot v_{k\Lambda_0} = kv_{k\Lambda_0}, \quad L_0 \cdot v_{k\Lambda_0} = 0, \quad (\mathfrak{g} \otimes t^n) \cdot v_{k\Lambda_0} = 0$$

for $n \geq 0$. Form the induced $\tilde{\mathfrak{g}}$ -module (a generalized Verma module)

$$(3.4) \quad N(k\Lambda_0) = U(\hat{\mathfrak{g}}) \otimes_{U(\mathfrak{p})} \mathbb{F}v_{k\Lambda_0} = U(\tilde{\mathfrak{g}}) \otimes_{U(\tilde{\mathfrak{g}}_{\geq 0})} \mathbb{F}v_{k\Lambda_0}.$$

Then $N(k\Lambda_0) \cong U(\tilde{\mathfrak{g}}_{<0})$ as vector spaces, where

$$\tilde{\mathfrak{g}}_{<0} = \coprod_{n < 0} \mathfrak{g} \otimes t^n.$$

We denote by $x(n)$ the operator on $N(k\Lambda_0)$ corresponding to $x \otimes t^n$. Set

$$(3.5) \quad \mathbf{1} = 1 \otimes v_{k\Lambda_0} \in N(k\Lambda_0).$$

Then L_0 satisfies

$$(3.6) \quad L_0 \cdot x_1(n_1) \cdots x_r(n_r)v_{k\Lambda_0} = \sum_j (-n_j)x_1(n_1) \cdots x_r(n_r)v_{k\Lambda_0}.$$

Let L_{-1} be the operator on $N(k\Lambda_0)$ determined by

$$(3.7) \quad L_{-1} \cdot x_1(n_1) \cdots x_r(n_r)v_{k\Lambda_0} = \sum_j (-n_j)x_1(n_1) \cdots x_j(n_j - 1) \cdots x_r(n_r)v_{k\Lambda_0}.$$

Then $N(k\Lambda_0)$, L_{-1} , L_0 , $\mathbf{1}$ satisfy (2.17–2.19, 2.30). For $x \in \mathfrak{g}$ consider the generating function of operators on $N(k\Lambda_0)$

$$(3.8) \quad x(z) = \sum_{n \in \mathbb{Z}} x(n)z^{-n-1}.$$

When we think of $x(n)$ as a coefficient of $x(z)$, we sometimes write x_n instead. It is easy to show that each $x(z)$ is admissible of conformal weight 1 in the sense of Definition 2.2. Set

$$(3.9) \quad U = \mathfrak{g}(-1) \otimes v_{k\Lambda_0} = N(k\Lambda_0)_1 \subset N(k\Lambda_0).$$

We shall often identify U with $\mathfrak{g}$. Define

$$(3.10) \quad Y : U \rightarrow (\text{End}_{\mathbb{F}} N(k\Lambda_0))[[z, z^{-1}]]$$

by

$$(3.11) \quad Y(x(-1) \otimes v_{k\Lambda_0}, z) = x(z),$$

for $x \in \mathfrak{g}$. Since for $x, y \in \mathfrak{g}$,

$$(3.12) \quad [x(z_1), y(z_2)] = z_2^{-1} \delta(z_1/z_2) [x, y](z_2) - k(x, y) z_2^{-1} \frac{\partial}{\partial z_1} \delta(z_1/z_2),$$

Y satisfies the hypotheses of Theorem 2.6. Hence Y extends to make $N(k\Lambda_0)$ a vertex operator algebra except that (so far) only L_{-1}, L_0 out of the Virasoro algebra act on $N(k\Lambda_0)$.

To complete the construction of a vertex operator algebra we need a conformal vector ω giving rise to the Virasoro algebra. For this we use the following so called Sugawara construction. Let $\{x^i\}_{i \in I}, \{y^i\}_{i \in I}$ be dual bases in $\mathfrak{g}$. Set

$$(3.13) \quad \phi = \sum_i (x^i)_{-1} (y^i)_{-1} \mathbf{1} = \sum_i (x^i)_{-1} y^i \in N(k\Lambda_0)_2.$$

There is a nondegenerate bilinear form on $\mathfrak{g} \otimes \mathfrak{g}$ determined by

$$(u \otimes v, x \otimes y) = (u, x)(v, y).$$

For $x \in \mathfrak{g}, m, n \in I$,

$$\begin{aligned} & \left(\sum_i ([x, x^i] \otimes y^i + x^i \otimes [x, y^i]), y^m \otimes x^n \right) \\ &= ([x, x^n], y^m) + ([x, y^m], x^n) = 0. \end{aligned}$$

Hence

$$\sum_i ([x, x^i] \otimes y^i + x^i \otimes [x, y^i]) = 0 \in \mathfrak{g} \otimes \mathfrak{g}.$$

It follows that

$$(3.14) \quad x_0 \phi = \sum_i \{ [x, x^i]_{-1} y^i + (x^i)_{-1} [x, y^i] \} = 0$$

for all $x \in \mathfrak{g}$. We have

$$\begin{aligned} (3.15) \quad x_1 \phi &= \sum_i \{ [[x, x^i], y^i] + k(x, x^i) y^i + k(x, y^i) x^i \} \\ &= 2kx + \sum_i [[x, x^i], y^i] \\ &= [2k + (\theta, \theta + 2\rho)]x \end{aligned}$$

if θ is the highest root and ρ half the sum of the positive roots. Now

$$(3.16) \quad (\rho, \theta) = \rho(\nu^{-1}(\theta)) = \rho(h_\theta) = \rho\left(\sum a_j^\vee \alpha_j^\vee\right) = \sum a_j^\vee = g - 1$$

(where g is the dual Coxeter number, and ν is the isomorphism between the Cartan subalgebra of $\mathfrak{g}$ and its dual defined via the bilinear form $(\cdot, \cdot)$, cf. [K]) so that

$$x_1\phi = 2(k + g)x.$$

Since there is no $\mathfrak{g}$ -invariant map $\mathfrak{g} \rightarrow \mathbb{F}$ we must have

$$x_2\phi = 0$$

for all $x \in \mathfrak{g}$, and by degree consideration $x_n\phi = 0$ for $n \geq 3$. The commutator formula [FLM, (8.6.3), (A.3.11)] gives

$$\begin{aligned} [x(z_1), Y(\phi, z_2)] &= -2(k + g)z_2^{-1} \frac{\partial}{\partial z_1} \delta(z_1/z_2)x(z_2), \\ [\phi_{m+1}, x_n] &= -2(k + g)nx_{m+n}. \end{aligned}$$

If $k \neq -g$ we define

$$(3.17) \quad \omega = \frac{1}{2(k + g)}\phi,$$

$$(3.18) \quad L_m = \omega_{m+1}$$

and have

$$(3.19) \quad [L_m, x_n] = -nx_{m+n}$$

so that L_0, L_{-1} coincide with the operators above. Also,

$$\begin{aligned} L_0\omega &= 2\omega, \\ L_1\omega &= \frac{1}{2(k + g)} \sum_i [x^i, y^i] = 0, \\ L_2\omega &= \frac{1}{2(k + g)} \sum_i k(x^i, y^i)\mathbf{1} = \frac{k \dim \mathfrak{g}}{2(k + g)}\mathbf{1} \end{aligned}$$

and $L_n\omega = 0$ for $n \geq 3$. It now follows from the commutator formula that

$$(3.20) \quad [L_m, L_n] = (m - n)L_{m+n} + \frac{m^3 - m}{12} \delta_{m+n,0} \frac{k \dim \mathfrak{g}}{k + g}$$

so that the Virasoro algebra acts on $N(k\Lambda_0)$. We can summarize the results of this section as:

Theorem 3.1. *For every $k \neq -g$, $(N(k\Lambda_0), Y, \mathbf{1}, \omega)$ is a vertex operator algebra with*

$$(3.21) \quad \text{rank } N(k\Lambda_0) = \frac{k \dim \mathfrak{g}}{k + g}.$$

4. MODULES OVER $N(k\Lambda_0)$

In this section we show that when $k \neq -g$ it is possible to extend the operators from $\tilde{\mathfrak{g}}$ on any highest weight module of level k so that the $\tilde{\mathfrak{g}}$ -module becomes a module over the vertex operator algebra $N(k\Lambda_0)$ in the sense of [FLM, 8.10].

Remark. If $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{F})$ and $M(\Lambda)$ is any Verma module over $\tilde{\mathfrak{g}}$ of level $-g = -2$ then there is no operator L_{-1} on $M(\Lambda)$ such that $[L_{-1}, u(n)] = -nu(n-1)$ for $u \in \mathfrak{g}$ and $[L_0, L_{-1}] = L_{-1}$. It is not even possible to define $L_{-1}v_\Lambda$ so as to satisfy

$$x(0)L_{-1}v_\Lambda = 0, \quad y(1)L_{-1}v_\Lambda = y(0)v_\Lambda.$$

To show this, note that since the weight of $L_{-1}v_\Lambda$ is determined, $L_{-1}v_\Lambda$ would have to be a linear combination of $h(-1)v_\Lambda$ and $x(-1)y(0)v_\Lambda$. It is easy to check that the above system of equations has no solution.

Fix a splitting Cartan subalgebra $\mathfrak{h}$ in $\mathfrak{g}$, let $\Phi \subset \mathfrak{h}^*$ be the associated root system and choose a set of simple roots $\{\alpha_1, \dots, \alpha_\ell\} \subset \Phi$. Let W be a level k module over $\hat{\mathfrak{g}} \rtimes (\mathbb{F}L_0 \oplus \mathbb{F}L_{-1}) = \tilde{\mathfrak{g}} \rtimes \mathbb{F}L_{-1}$, a Lie algebra with the action of L_0, L_{-1} on $\hat{\mathfrak{g}}$ determined by (3.19). Consider $\mathfrak{h}^e = \mathfrak{h}(0) \oplus \mathbb{F}c \oplus \mathbb{F}L_0$ as Cartan subalgebra in $\hat{\mathfrak{g}} \rtimes (\mathbb{F}L_0 \oplus \mathbb{F}L_{-1})$. Assume that

$$(4.1) \quad W = \coprod_{\mu \in \mathfrak{h}^{e*}} W_\mu,$$

$$(4.2) \quad \dim W_\mu < \infty$$

and that there is $\tau \in \mathbb{F}$ such that for every weight μ of W

$$(4.3) \quad \mu(L_0) \in \tau + \mathbb{N}.$$

Remark. An easy calculation (cf. [H]) shows that the relation

$$[L_m, x_n] = -nx_{m+n},$$

from the Sugawara construction in Section 3, takes place in the completed enveloping algebra $\overline{U(\tilde{\mathfrak{g}})}$ [MP] when $k \neq -g$. Hence the Sugawara operator L_{-1} can be used to show that when $k \neq -g$ any highest weight module of level k satisfies the above conditions.

Set

$$W' = \coprod_{\mu \in \mathfrak{h}^{e*}} W_\mu^*$$

and denote the pairing between W' and W also by

$$\langle \cdot, \cdot \rangle : W' \times W \rightarrow \mathbb{F}.$$

For simplicity, set $V = N(k\Lambda_0)$. Our main goal is to construct for each $w \in W$ a so called intertwining operator

$$Y(w, z) \in (\text{Hom}_{\mathbb{F}}(V, W))[[z, z^{-1}]].$$

For

$$u = x(-1) \otimes v_{k\Lambda_0} \in U,$$

we change the meaning of $Y(u, z)$ so as to denote the generating function

$$Y(u, z) = x(z) = \sum_{n \in \mathbb{Z}} x(n) z^{-n-1}$$

of operators on $V \oplus W$ leaving invariant V and W . The intertwining operator should be local with respect to this extension i.e.

$$(z_1 - z_2)^N [Y(u, z_1), Y(w, z_2)] = 0$$

for some $N \in \mathbb{N}$, and all $u \in U$.

With $T(\hat{\mathfrak{g}})$ the tensor algebra, consider the associative algebra

$$T_k = T(\hat{\mathfrak{g}})/(c - k).$$

Then T_k has a basis consisting of monomials

$$x_1(m_1) \otimes \cdots \otimes x_r(m_r)$$

with the x_j chosen from a basis of $\mathfrak{g}$. Let $u_2, \dots, u_n \in U$, $w \in W$, $w' \in W'$. Set

$$P = \prod_{2 \leq \ell < m \leq n} (z_\ell - z_m)^2.$$

Then

$$\begin{aligned} & P \langle w', Y(u_2, z_2) \cdots Y(u_n, z_n) e^{z_1 L_{-1}} w \rangle \\ &= P \langle w', e^{z_1 L_{-1}} Y(u_2, z_2 - z_1) \cdots Y(u_n, z_n - z_1) w \rangle \\ &= P \langle w', e^{z_1 L_{-1}} Y(u_{\pi 2}, z_{\pi 2} - z_1) \cdots Y(u_{\pi n}, z_{\pi n} - z_1) w \rangle \end{aligned}$$

for all $\pi \in \text{Sym}\{2, \dots, n\}$. It follows that there is $N \in \mathbb{N}$ such that if

$$Q = \prod_{\ell=2}^n (z_\ell - z_1)^N,$$

then

$$(4.4) \quad PQ \langle w', Y(u_2, z_2) \cdots Y(u_n, z_n) e^{z_1 L_{-1}} w \rangle \in \mathbb{F}[z_1, \dots, z_n].$$

We have also that

$$\langle w', Y(u_2, z_2) \cdots Y(u_n, z_n) e^{z_1 L_{-1}} w \rangle \in \mathbb{F}((z_2)) \cdots ((z_n))((z_1)),$$

hence there is $f \in \mathcal{R}_n$ such that

$$(4.5) \quad \langle w', Y(u_2, z_2) \cdots Y(u_n, z_n) e^{z_1 L_{-1}} w \rangle = \iota_{2 \dots n 1} f.$$

Clearly the map

$$\begin{aligned} \mathfrak{g} \times \cdots \times \mathfrak{g} &\rightarrow \mathcal{R}_n \\ (u_2, \dots, u_n) &\mapsto f \end{aligned}$$

is multilinear. Consideration of weights with respect to $\mathfrak{h}^e$ now show that we can define

$$Y(w, z) \in (\text{Hom}_{\mathbb{F}}(T_k, W))[[z, z^{-1}]]$$

by the condition that

$$(4.6) \quad \langle w', Y(w, z_1)Y(u_2, z_2) \cdots Y(u_n, z_n) \rangle = \iota_{12 \dots n} f$$

for all $w' \in W'$ with f as in (4.5). Here we use $\dim W_\mu < \infty$ in order that $W_\mu^{**} = W_\mu$.
Set

$$U_k(\hat{\mathfrak{g}}) = U(\hat{\mathfrak{g}})/(c - k), \quad U_k(\tilde{\mathfrak{g}}) = U(\tilde{\mathfrak{g}})/(c - k).$$

By the construction of the enveloping algebra, $U_k(\hat{\mathfrak{g}})$ is then the quotient of T_k by the two-sided ideal generated by

$$x(m)y(n) - y(n)x(m) - [x, y](m + n) - m\delta_{m+n,0}(x, y)k$$

for all $x, y \in \mathfrak{g}$, $m, n \in \mathbb{Z}$. We want to show that the components of $Y(w, z)$ induce operators $U_k(\hat{\mathfrak{g}}) \rightarrow W$. We thus consider the effect of interchanging the operators $Y(u_j, z_j)$, and $Y(u_{j+1}, z_{j+1})$ in (4.5). For clarity we raise the indices on ι so that $\iota_{a,b,\dots,m} = \iota(a, b, \dots, m)$. Multiplication by $(z_j - z_{j+1})^2$ show that

$$\begin{aligned} \langle w', Y(u_2, z_2) \cdots Y(u_{j+1}, z_{j+1})Y(u_j, z_j) \cdots Y(u_n, z_n) e^{z_1 L - 1} w \rangle = \\ = \iota(2, \dots, j+1, j, \dots, n, 1) f \end{aligned}$$

with f as in (4.5). By (4.4) the order of the pole $(z_j - z_{j+1})^{-1}$ in f is at most 2. It is easy to see that there is a unique representation

$$(4.7) \quad f = \frac{f_1}{z_j - z_{j+1}} + \frac{f_2}{(z_j - z_{j+1})^2} + p$$

with $f_1, f_2 \in \mathcal{R}_{\{1, \dots, \hat{j}, \dots, n\}}$ (i.e. independent of z_j) and with p having the poles of functions in $\mathcal{R}_n$ except that p does not have the pole $(z_j - z_{j+1})^{-1}$. It follows that

$$\begin{aligned} (4.8) \quad \{ \iota(2, \dots, n, 1) - \iota(2, \dots, j+1, j, \dots, n, 1) \} f = \\ = z_{j+1}^{-1} \delta(z_j/z_{j+1}) \iota(2, \dots, \hat{j}, \dots, n, 1) f_1 \\ - z_{j+1}^{-1} \frac{\partial}{\partial z_j} \delta(z_j/z_{j+1}) \iota(2, \dots, \hat{j}, \dots, n, 1) f_2. \end{aligned}$$

Let $x, y \in \mathfrak{g}$ so that

$$u_j = x(-1) \otimes v_{k\Lambda_0}, \quad u_{j+1} = y(-1) \otimes v_{k\Lambda_0}$$

and set

$$\bar{u} = [x, y](-1) \otimes v_{k\Lambda_0} \in U.$$

There are $g \in \mathcal{R}_{\{1, \dots, \hat{j}, \dots, n\}}$, $h \in \mathcal{R}_{\{1, \dots, \hat{j}, \widehat{j+1}, \dots, n\}}$ such that

$$\begin{aligned} \langle w', Y(u_2, z_2) \cdots Y(u_{j-1}, z_{j-1}) Y(\bar{u}, z_{j+1}) \cdots Y(u_n, z_n) e^{z_1 L-1} w \rangle = \\ = \iota(2, \dots, \hat{j}, \dots, n, 1)g, \end{aligned}$$

and

$$\begin{aligned} \langle w', Y(u_2, z_2) \cdots Y(u_{j-1}, z_{j-1}) Y(u_{j+2}, z_{j+2}) \cdots Y(u_n, z_n) e^{z_1 L-1} w \rangle \\ = \iota(2, \dots, \hat{j}, \widehat{j+1}, \dots, n, 1)h. \end{aligned}$$

Since W is a $\hat{\mathfrak{g}}$ -module we have

$$\begin{aligned} \{\iota(2, \dots, n, 1) - \iota(2, \dots, j+1, j, \dots, n, 1)\}f = \\ = z_{j+1}^{-1} \delta(z_j/z_{j+1}) \iota(2, \dots, \hat{j}, \dots, n, 1)g \\ - k(x, y) z_{j+1}^{-1} \frac{\partial}{\partial z_j} \delta(z_j/z_{j+1}) \iota(2, \dots, \hat{j}, \widehat{j+1}, \dots, n, 1)h. \end{aligned}$$

Comparison with (4.8) yields

$$f_1 = g, \quad f_2 = k(x, y)h$$

(after multiplication by $(z_j - z_{j+1})$). (Thus f_2 does not depend on z_{j+1} .) Hence

$$\begin{aligned} \{\iota(1, 2, \dots, n) - \iota(1, 2, \dots, j+1, j, \dots, n)\}f = \\ = z_{j+1}^{-1} \delta(z_j/z_{j+1}) \iota(1, 2, \dots, \hat{j}, \dots, n)g \\ - k(x, y) z_{j+1}^{-1} \frac{\partial}{\partial z_j} \delta(z_j/z_{j+1}) \iota(1, 2, \dots, \hat{j}, \widehat{j+1}, \dots, n)h \end{aligned}$$

so that

$$\begin{aligned} \langle w', Y(w, z_1) Y(u_2, z_2) \cdots Y(u_n, z_n) \rangle - \langle w', Y(w, z_1) \cdots Y(u_{j+1}, z_{j+1}) Y(u_j, z_j) \cdots \rangle \\ = z_{j+1}^{-1} \delta(z_j/z_{j+1}) \langle w', Y(w, z_1) Y(u_2, z_2) \cdots Y(\bar{u}, z_{j+1}) \cdots Y(u_n, z_n) \rangle \\ - k(x, y) z_{j+1}^{-1} \frac{\partial}{\partial z_j} \delta(z_j/z_{j+1}) \langle w', Y(w, z_1) \cdots Y(u_{j-1}, z_{j-1}) Y(u_{j+2}, z_{j+2}) \cdots \rangle. \end{aligned}$$

This means that each component of $Y(w, z)$ annihilates the two-sided ideal which is the kernel of $T_k \rightarrow U_k(\hat{\mathfrak{g}})$ and hence we have well defined operators

$$Y(w, z) \in (\text{Hom}_{\mathbb{F}}(U_k(\hat{\mathfrak{g}}), W))[[z, z^{-1}]].$$

Furthermore, from (4.4) there are no poles z_n^{-1} in f satisfying (4.5), hence

$$\iota_{12 \dots n} f = \sum_{d \leq -1} f_d z_n^{-d-1}$$

with $f_d \in \mathbb{F}((z_1)) \cdots ((z_{n-1}))$ so that

$$\langle w', Y(w, z_1)Y(u_2, z_2) \cdots Y(u_{n-1}, z_{n-1})(u_n)_d \rangle = 0$$

if $d \geq 0$. Hence all components of $Y(w, z)$ annihilate all vectors in $U_k(\hat{\mathfrak{g}})$ of the form

$$x_2(m_2) \cdots x_{n-1}(m_{n-1})x_n(d)$$

with $d \geq 0$. Since these span the left ideal in $U_k(\hat{\mathfrak{g}})$ which is the kernel of

$$U_k(\hat{\mathfrak{g}}) \rightarrow N(k\Lambda_0),$$

$Y(w, z)$ induces a well defined operator

$$Y(w, z) \in (\text{Hom}_{\mathbb{F}}(N(k\Lambda_0), W))[[z, z^{-1}]].$$

This is the intertwining operator that we want. We have proved:

Proposition 4.1. *Let $u_2, \dots, u_n \in U$, $w \in W$, $w' \in W'$.*

(i) *There is $f \in \mathcal{R}_n$ such that*

$$\langle w', Y(u_2, z_2) \cdots Y(u_n, z_n)e^{z_1 L^{-1}}w \rangle = \iota_{2 \dots n 1} f.$$

(ii) *Using these matrix coefficients f , there is a well defined operator*

$$Y(w, z) \in (\text{Hom}_{\mathbb{F}}(N(k\Lambda_0), W))[[z, z^{-1}]]$$

determined by

$$(4.9) \quad \langle w', Y(w, z_1)Y(u_2, z_2) \cdots Y(u_n, z_n)\mathbf{1} \rangle = \iota_{12 \dots n} f.$$

Now it is easy to show that $Y(w, z)$ is local with respect to the operators $Y(u, z)$, $u \in U$:

Proposition 4.2. *Let $u_0, u_2, \dots, u_n \in U$, $w \in W$, $w' \in W'$. Then*

$$(4.10) \quad \langle w', Y(u_0, z_0)Y(w, z_1)Y(u_2, z_2) \cdots Y(u_n, z_n)\mathbf{1} \rangle = \iota_{012 \dots n} f$$

if $f \in \mathcal{R}_{n+1}$ is such that

$$\langle w', Y(u_0, z_0)Y(u_2, z_2) \cdots Y(u_n, z_n)e^{z_1 L^{-1}}w \rangle = \iota_{02 \dots n 1} f.$$

If $N \in \mathbb{N}$ is such that $(u_0)_d w = 0$ when $d \geq N$, then the order of the pole $(z_0 - z_1)^{-1}$ in f is at most N . Hence $Y(w, z)$ is local with respect to $Y(u_0, z)$ for all $u_0 \in U$.

Proof. Let

$$\iota_{0 \dots n} f = \sum_{d \in \mathbb{Z}} f_d z_0^{-d-1}.$$

Then

$$\langle w', (u_0)_d Y(u_2, z_2) \cdots Y(u_n, z_n)e^{z_1 L^{-1}}w \rangle = \iota_{2 \dots n 1} f_d$$

by Proposition 1.1.3. The definition of $Y(w, z_1)$ gives

$$\langle w', (u_0)_d Y(w, z_1) Y(u_2, z_2) \cdots Y(u_n, z_n) \mathbf{1} \rangle = \iota_{12 \cdots n} f_d,$$

and (4.10) now follows from Proposition 1.1.3. From

$$\begin{aligned} & \prod_{\ell=2}^n (z_0 - z_\ell)^2 \iota_{02 \cdots n} f = \\ &= \prod_{\ell=2}^n (z_0 - z_\ell)^2 \langle w', Y(u_0, z_0) Y(u_2, z_2) \cdots Y(u_n, z_n) e^{z_1 L_{-1}} w \rangle \\ &= \prod_{\ell=2}^n (z_0 - z_\ell)^2 \langle w', e^{z_1 L_{-1}} Y(u_2, z_2 - z_1) \cdots Y(u_n, z_n - z_1) Y(u_0, z_0 - z_1) w \rangle \end{aligned}$$

we can determine the order of the pole $(z_0 - z_1)^{-1}$ in f . $\square$

Theorem 4.3. *Let $k \neq -g$, and let W be a $\hat{\mathfrak{g}} \rtimes (\mathbb{F}L_0 \oplus \mathbb{F}L_{-1})$ -module of level k satisfying (4.1–3). (In particular W could be any highest weight module or a module from the category $\mathcal{O}$.) Then there is a unique extension of the operators $Y(u, z)$, $u \in U$ on W that make W into a module over the vertex operator algebra $N(k\Lambda_0)$.*

Proof. Let $\tilde{V}$ be the space of all admissible generating functions

$$a(z) = \sum_{n \in \mathbb{Z}} a_n z^{-n-1}$$

of operators a_n on $V \oplus W$ stabilizing both V and W such that

$$a(z) \text{ is local w.r.t. } \{Y(u, z) \mid u \in U\} \cup \{Y(w, z) \mid w \in W\}.$$

Then the proof of Theorem 2.6 applies with only minor modifications to show that

1. the map $\tilde{V} \rightarrow V$, $a(z) \mapsto a_{-1} \mathbf{1}$ is injective,
2. the image of this map is all of V ,
3. any two $a(z), b(z) \in \tilde{V}$ are mutually local.

Given this, the argument of [FLM, A.2.5] can be employed, using the intertwining operator $Y(w, z)$ in the rôle of one vertex operator, to show that W satisfies the “associativity condition” [FLM, (8.10)] and hence the Jacobi identity. $\square$

We have seen that for level k highest weight modules, or modules from the category $\mathcal{O}$, the action of $\tilde{\mathfrak{g}}$ determined by $x(z) = \sum x_n z^{-n-1}$ extends, by using normal order products (2.14), to the action of $N(k\Lambda_0)$ defined by $Y(v, z) = \sum v_n z^{-n-1}$, $v \in N(k\Lambda_0)$. While x_n are elements in $U_k(\tilde{\mathfrak{g}})$, in general the coefficients v_n are infinite sums of elements in $U_k(\tilde{\mathfrak{g}})$ and should be seen in some kind of completion $\overline{U_k(\tilde{\mathfrak{g}})}$. We prefer to think in terms of completion defined in [MP]: a net $(y_i)_{i \in I}$ converges to y if for each $\tilde{\mathfrak{g}}$ -module V in the category $\mathcal{O}$ and each vector v in V there is an index $i_0 \in I$ such that $i \geq i_0$ implies $y_i \cdot v = y_{i_0} \cdot v = y \cdot v$.

Remark. Constructions and results analogous to Theorem 3.1 and Theorem 4.3 have appeared in [FF] and in relation to the Wess-Novikov-Zumino-Witten model in the physics literature. Theorems 3.1 and 4.3 contain the precise results we require in the following sections.

It is easy to see that normal order product (2.14) coincides with the normal order product defined in [H] if one identifies $\mathfrak{g} \cong \mathfrak{g}(-1)\mathbf{1}$ and $S(\tilde{\mathfrak{g}}_{<0}) \cong U(\tilde{\mathfrak{g}}_{<0}) \cong N(k\Lambda_0)$ via the symmetrization map. Hence we work with the same set of fields, but the “index set” $N(k\Lambda_0)$ brings into the game the action of $\tilde{\mathfrak{g}}$.

Before our work was completed, other proofs of Theorems 3.1 and 4.3 appeared: in [FZ] by using quite different ideas, and recently in [Li] by using the locality in a similar way, but from a different point of view, and proving in an elegant way stronger and more general results.

5. RELATIONS ON STANDARD MODULES

We identify $\tilde{\mathfrak{g}}$ with a Kac-Moody algebra in the usual way (cf. [K]): first choose root vectors $x_\alpha \in \mathfrak{g}_\alpha$, such that $h_\alpha = [x_\alpha, x_{-\alpha}]$ satisfy $\alpha(h_\alpha) = 2$ for $\alpha \in \Phi$, and then set $e_i = x_{\alpha_i}(0)$, $f_i = x_{-\alpha_i}(0)$, for $i = 1, \dots, \ell$, $e_0 = x_{-\theta}(1)$, $f_0 = x_\theta(-1)$ (θ the highest root as in Section 3) and $h_i = [e_i, f_i]$ for $i = 0, 1, \dots, \ell$. Then we have the usual triangular decompositions $\mathfrak{g} = \mathfrak{n}_- + \mathfrak{h} + \mathfrak{n}_+$ and $\tilde{\mathfrak{g}} = \tilde{\mathfrak{n}}_- + \mathfrak{h}^e + \tilde{\mathfrak{n}}_+$, where $\mathfrak{h}^e = \mathfrak{h} + \mathbb{F}c + \mathbb{F}L_0$. Denote by

$$(5.1) \quad P_+ = \{\Lambda \in \mathfrak{h}^{e*} \mid \Lambda(h_i) \in \mathbb{N} \text{ for } i = 0, 1, \dots, \ell\},$$

the set of dominant integral weights, and let for $i = 0, 1, \dots, \ell$, $\Lambda_i \in P_+$ be the fundamental weight determined by $\Lambda_i(h_j) = \delta_{ij}$, $\Lambda_i(L_0) = 0$.

For $\Lambda \in \mathfrak{h}^{e*}$ we shall use the notation $M(\Lambda)$ for the Verma module with highest weight Λ and $L(\Lambda)$ for the irreducible quotient of $M(\Lambda)$. In this section we show that the standard $\tilde{\mathfrak{g}}$ -modules $L(\Lambda)$ of level k , i.e. $\Lambda \in P_+$, $\Lambda(c) = k$, satisfy the relation $x_\theta(z)^{k+1} = 0$, and that in some sense this relation determines the structure of $L(\Lambda)$, see Theorem 5.14. The relation $x_\theta(z)^{k+1} = 0$ and its consequences will be studied further in the following sections.

For a given nontrivial finite dimensional irreducible $\mathfrak{g}$ -module W we have the associated loop module $\bar{W} = W \otimes \mathbb{C}[t, t^{-1}]$ with the action of $\tilde{\mathfrak{g}}$ defined by (cf. [CP]):

$$x \otimes t^n \cdot w \otimes t^m = x \cdot w \otimes t^{n+m}, \quad c \cdot w \otimes t^m = 0, \quad L(0) \cdot w \otimes t^m = -m w \otimes t^m.$$

The results in this section are mostly about the loop module $\bar{R}$ constructed from the coefficients of the annihilating field $x_\theta(z)^{k+1}$.

Theorem 5.1. *Let $k \in \mathbb{F}$ and let R be a subspace of $N(k\Lambda_0)$ invariant under $\mathfrak{g} = \mathfrak{g}(0)$ and L_0 . Set*

$$(5.2) \quad \bar{R} = \mathbb{F}\text{-span}\{r_n \mid r \in R, n \in \mathbb{Z}\} \subset \text{End}_{\mathbb{F}} N(k\Lambda_0).$$

(i) *The following are equivalent:*

(a) *For all $x \in \mathfrak{g}$, $r \in R$, $m, n \in \mathbb{Z}$,*

$$(5.3) \quad [x(m), r_n] = (x(0)r)_{m+n}$$

so that $\bar{R}$ is a loop module under the “adjoint” action of $\tilde{\mathfrak{g}}$.

(b) *We have $\mathfrak{g}(n)R = 0$ for all positive integers n .*

(ii) *If (a) and (b) are satisfied then $\bar{R}$ is irreducible iff R is a nontrivial irreducible $\mathfrak{g}$ -module.*

Proposition 5.2. *There is a bijection between*

- (a) *the irreducible loop modules $\bar{R}$ associated to subspaces R of $N(k\Lambda_0)$ invariant under $\mathfrak{g}$ and L_0 as in Theorem 5.1,*
- (b) *the singular vectors (up to nonzero scalar multiples) in $N(k\Lambda_0)$ that are not invariant under $\mathfrak{g}$.*

Proof. If $\bar{R}$ is an irreducible loop module then R is a nontrivial irreducible $\mathfrak{g}$ -module by Theorem 5.1(ii) and is hence by the theorem of the highest weight determined by a highest weight vector with respect to $(\mathfrak{h}, \Phi^+)$. This vector is by condition (b) in Theorem 5.1 also a singular vector. We must also show that R is determined by $\bar{R}$. Since R is irreducible under $\mathfrak{g}$, R must be homogeneous. It then follows from $Y(r, z)\mathbf{1} = e^{zL_{-1}}r$ that

$$\bar{R}\mathbf{1} = \coprod_{n \geq 0} L_{-1}^n R$$

and hence R is determined as the homogeneous component of highest degree that is nonzero in $\bar{R}\mathbf{1}$. $\square$

Proof of Theorem 5.1. (i) For $x \in \mathfrak{g}$, $r \in R$,

$$[x(z_1), Y(r, z_2)] = \sum_{i \geq 0} \frac{(-1)^i}{i!} \left(\frac{\partial}{\partial z_1} \right)^i z_2^{-1} \delta(z_1/z_2) Y(x(i)r, z_2).$$

Hence (b) implies (a). Furthermore, if (a) holds and $N \in \mathbb{N}$ such that $x(i)r = 0$ for $i > N$, then

$$\sum_{i=1}^N \frac{(-1)^i}{i!} \left(\frac{\partial}{\partial z_1} \right)^i \delta(z_1/z_2) Y(x(i)r, z_2) = 0,$$

and

$$\begin{aligned} \text{coeff}_{z_1^0} \sum_{i=1}^N \frac{(-1)^i}{i!} \left(\frac{\partial}{\partial z_1} \right)^i \delta(z_1/z_2) Y(x(i)r, z_2) \mathbf{1} &= \\ &= \sum_{i=1}^N (-1)^i z_2^{-i} (x(i)r + z_2(\cdots)) = 0, \end{aligned}$$

so that $x(N)r = 0$. Repeating this argument shows successively that $x(N-1)r = 0$, $\dots$, $x(1)r = 0$, and hence (b).

(ii) If S is a nonzero proper submodule of R then clearly $\bar{S}$ is a nonzero proper submodule of $\bar{R}$. If R is a 1-dimensional trivial $\mathfrak{g}$ -module, then $\mathbb{F}r_n$ is a $\tilde{\mathfrak{g}}$ -submodule of $\bar{R}$ for each $r \in R, n \in \mathbb{Z}$. Conversely, assume that R is nontrivial and irreducible and let Z be a nonzero $\tilde{\mathfrak{g}}$ -submodule of $\bar{R}$. From $[L_0, Z] \subset Z$ it follows that $Z = \oplus_{n \in \mathbb{Z}} Z(n)$ with $Z(n) = \{p \in Z \mid [L_0, p] = np\}$. For some n , $Z(n) \neq 0$ and then $[\mathfrak{g}, Z(n)] \subset Z(n)$ implies that

$$Z(n) = \{r_m \mid r \in R\}$$

for an appropriate m . It now follows from (5.3) that $Z = \bar{R}$. $\square$

Proposition 5.3. [K, Corollary 10.4] *Let $\Lambda \in P_+$. Then the annihilator in $U(\tilde{\mathfrak{g}})$ of a highest weight vector of $L(\Lambda)$ is the left ideal generated by e_i , $f_i^{\Lambda(h_i)+1}$, and $h - \Lambda(h)$ where $i = 0, 1, \dots, \ell$ and $h \in \mathfrak{h}^e$.*

For $k \in \mathbb{F}$ let $N^1(k\Lambda_0)$ denote the maximal $\tilde{\mathfrak{g}}$ -submodule of $N(k\Lambda_0)$.

In the remainder of this section we shall assume that k is a positive integer.

Corollary 5.4. *The maximal submodule $N^1(k\Lambda_0)$ of $N(k\Lambda_0)$ is generated by the vector $x_\theta(-1)^{k+1}\mathbf{1}$.*

Proof. This follows immediately from Proposition 5.3 since for $i = 1, \dots, \ell$,

$$f_i^{k\Lambda_0(h_i)+1}\mathbf{1} = f_i\mathbf{1} = x_{-\alpha_i}(0)\mathbf{1} = 0,$$

and we have $f_0^{k\Lambda_0(h_0)+1}\mathbf{1} = f_0^{k+1}\mathbf{1} = x_\theta(-1)^{k+1}\mathbf{1}$. $\square$

Set

$$(5.4) \quad R = U(\mathfrak{g})x_\theta(-1)^{k+1}\mathbf{1} \subset N(k\Lambda_0).$$

Since $\mathfrak{n}_+(0)$ annihilates $x_\theta(-1)^{k+1}\mathbf{1}$, and $N(k\Lambda_0)$ is a sum of finite dimensional $\mathfrak{g}(0)$ -modules, R is an irreducible $\mathfrak{g}$ -module.

Lemma 5.5. *The vector $x_\theta(-1)^{k+1}\mathbf{1} \in N(k\Lambda_0)$ is a singular vector. Hence $U(\tilde{\mathfrak{g}})x_\theta(-1)^{k+1}\mathbf{1} = U(\tilde{\mathfrak{n}}_-)x_\theta(-1)^{k+1}\mathbf{1}$.*

Proof. It remains to show that $x_{-\theta}(1)x_\theta(-1)^{k+1}\mathbf{1} = 0$. From

$$[x_{-\theta}(1), x_\theta(-1)] = -h_\theta + c,$$

and $(-h_\theta + c)\mathbf{1} = k\mathbf{1}$ it follows by a standard calculation in the representation theory of $\mathfrak{sl}(2, \mathbb{F})$ that $x_{-\theta}(1)x_\theta(-1)^{k+1}\mathbf{1} = 0$. $\square$

For $r \in R$, define $r(n)$ by

$$(5.5) \quad Y(r, z) = \sum_{n \in \mathbb{Z}} r(n)z^{-n-k-1}.$$

Then $r(n)$ has degree n and $r(n) = r_{n+k}$. By Theorem 5.1

$$(5.6) \quad [x(m), r(n)] = (x(0)r)(m+n)$$

for $x \in \mathfrak{g}$, $r \in R$, $m, n \in \mathbb{Z}$, so that

$$(5.7) \quad \bar{R} = \mathbb{F}\text{-span}\{r(n) \mid r \in R, n \in \mathbb{Z}\}$$

is a loop module. Let us record this as:

Corollary 5.6. *The coefficients of $Y(ux_\theta(-1)^{k+1}\mathbf{1}, z)$ with u ranging through $U(\mathfrak{g})$ span a loop module.*

Remark. The above statement can also be verified by direct calculation of commutators.

By the result of V. Chari and A. Pressley [CP, Theorem 4.5] we have that $\bar{R} \otimes L(\Lambda)$ is irreducible for each standard module $L(\Lambda)$ of level k , and hence the map

$$\begin{aligned} \bar{R} \otimes L(\Lambda) &\rightarrow L(\Lambda) \\ u \otimes v &\mapsto uv \end{aligned}$$

must be zero, i.e. $\bar{R}$ annihilates $L(\Lambda)$. We shall give another proof of this fact:

Lemma 5.7. *Let*

$$(5.8) \quad x = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad y = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

be the usual basis of $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{F})$, and set $\mathfrak{n}_+ = \mathbb{F}x$. Then

$$x^{k+1}y^{k+1} \in ch(h-1) \cdots (h-k) + U(\mathfrak{g})\mathfrak{n}_+,$$

for some nonzero integer c .

Proof. Clearly $u = x^{k+1}y^{k+1} \in p(h) + U(\mathfrak{g})\mathfrak{n}_+$ for some polynomial p of degree $k+1$. By applying u to a highest weight vector of a $(j+1)$ -dimensional irreducible $\mathfrak{g}$ -module, $j \in \{0, \dots, k, k+1\}$ we get $p(j) = 0$ for $j = 0, \dots, k$, $p(k+1) \neq 0$, and hence the lemma. $\square$

Lemma 5.8. *Let $\bar{W} = \coprod_{n \in \mathbb{Z}} W(n)$ be a loop module for $\tilde{\mathfrak{g}}$ which acts on a Verma module $M(\Lambda)$. Then*

$$\bar{W}M(\Lambda) = M(\Lambda) \quad \text{iff} \quad W(0)_0 v_\Lambda \neq 0,$$

where $W(0)_0$ denotes the zero-weight subspace of $\bar{W}$.

Proof. Note first that $\bar{W}M(\Lambda) = M(\Lambda)$ if and only if $v_\Lambda \in \bar{W}M(\Lambda)$. Since

$$\bar{W}M(\Lambda) = \bar{W}U(\tilde{\mathfrak{n}}_-)v_\Lambda = U(\tilde{\mathfrak{n}}_-)\bar{W}v_\Lambda,$$

the Λ weight subspace of $\bar{W}M(\Lambda)$ equals $W(0)_0 v_\Lambda$, and hence the lemma holds. $\square$

Theorem 5.9. *Let $L(\Lambda)$ be a standard $\tilde{\mathfrak{g}}$ -module of level k . Then*

$$\bar{R}L(\Lambda) = 0.$$

Proof. Consider first the case $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{F})$. Then

$$R(0) = U(\mathfrak{g}) \cdot \left(\sum_{j_1 + \dots + j_{k+1} = 0} y(j_1) \cdots y(j_{k+1}) \right)$$

(with $\cdot$ the “adjoint” action as above and y as in (5.8)) and for $V = M(\Lambda)$,

$$(5.9) \quad \begin{aligned} R(0)_0 v_\Lambda &= \mathbb{F} \left(x^{k+1} \cdot \left(\sum_{j_1 + \dots + j_{k+1} = 0} y(j_1) \cdots y(j_{k+1}) \right) \right) v_\Lambda \\ &= \mathbb{F} x^{k+1} y(0)^{k+1} v_\Lambda \\ &= \mathbb{F} p(\Lambda(h)) v_\Lambda, \end{aligned}$$

where by Lemma 5.7 $p(h) = h(h-1) \cdots (h-k)$. If $\Lambda(h) \in \{0, \dots, k\}$, then $R(0)_0 v_\Lambda = 0$ and by Lemma 5.8 $\bar{R}M(\Lambda)$ is a proper submodule, in particular $\bar{R}M(\Lambda) \subset M^1(\Lambda)$. Since we have an exact sequence

$$0 \rightarrow M^1(\Lambda) \rightarrow M(\Lambda) \rightarrow L(\Lambda) \rightarrow 0,$$

$\bar{R}$ acts trivially on $L(\Lambda)$.

In the general case we let $\mathfrak{a}$ be the subalgebra isomorphic to $\mathfrak{sl}(2, \mathbb{F})$ spanned by $x = x_\theta$, $y = x_{-\theta}$, $h = h_\theta$, and consider $L(\Lambda)$ as an $\mathfrak{a}$ -module by restriction. By [K, Theorem 10.7] $L(\Lambda)$ is a direct sum of standard level k $\mathfrak{a}$ -modules. By the $\mathfrak{sl}(2, \mathbb{F})$ -case above, $x_\theta(z)^{k+1}$ annihilates $L(\Lambda)$ and bracketing with $\tilde{\mathfrak{g}}$ now shows that $\bar{R}$ annihilates $L(\Lambda)$. $\square$

Definition 5.10. A subspace I of a vertex operator algebra V is said to be an *ideal* in V if and only if

$$Y(v, z)I \subset I[[z, z^{-1}]]$$

for all $v \in V$, i.e. I is invariant under the action of all components of all vertex operators.

Equivalently, I is an ideal if and only if I is a submodule of V considered as a module over itself.

Proposition 5.11. *Let I be an ideal of a vertex operator algebra V . Then the vertex operator algebra structure on V induces a structure of vertex operator algebra on V/I .*

Proof. Since there is a conformal vector $\omega \in V$ such that

$$Y(\omega, z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2},$$

I is invariant under the Virasoro algebra. From the relation

$$(5.10) \quad Y(u, z)v = e^{zL_{-1}}Y(v, -z)u$$

valid in any vertex operator algebra [FLM, (8.8.7), A.3.2; FHL] it now follows that $Y(u, z)V \subset I[[z, z^{-1}]]$ if $u \in I$, so that the induced operator $Y(v, z)$ on V/I only depends on $v \bmod I$. $\square$

Proposition 5.12. *Any $\tilde{\mathfrak{g}}$ -submodule of $N(k\Lambda_0)$ is also a vertex operator algebra ideal of $N(k\Lambda_0)$. There is in particular an induced vertex operator algebra structure on $L(k\Lambda_0)$.*

Proof. Let I be a $\tilde{\mathfrak{g}}$ -submodule of $N(k\Lambda_0)$. It then follows from

$$(5.11) \quad \begin{aligned} Y(u_n v, z) &= \\ &= \text{Res}_{z_0} z_0^n \{Y(u, z_0 + z)Y(v, z) - \text{Res}_{z_1} z_0^{-1} \delta\left(\frac{z - z_1}{-z_0}\right)Y(v, z)Y(u, z_1)\} \end{aligned}$$

for $u, v \in N(k\Lambda_0)$ (cf. (2.25), [FLM, (8.8.31)]) and induction that

$$Y(v, z)I \subset I[[z, z^{-1}]]$$

for all $v \in N(k\Lambda_0)$. $\square$

Proposition 5.13. *Let $\mathfrak{g}$ be a finite dimensional split simple Lie algebra over $\mathbb{F}$ and let $L(\Lambda)$ be a standard $\tilde{\mathfrak{g}}$ -module of level k .*

- (i) *We have $x_\theta(z)^{k+1} = 0$ on $L(\Lambda)$.*
- (ii) *The $\tilde{\mathfrak{g}}$ -module structure on $L(\Lambda)$ extends to make $L(\Lambda)$ a module over the vertex operator algebra $L(k\Lambda_0)$.*
- (iii) *If $\beta \in \Phi$ is a short root such that $(\beta, \beta) = \frac{1}{2}(\theta, \theta)$ then*

$$(5.12) \quad x_\beta(z)^{2k+1} = 0 \text{ on } L(\Lambda).$$

- (iv) *If $\gamma \in \Phi$ is a short root such that $(\gamma, \gamma) = \frac{1}{3}(\theta, \theta)$ then*

$$(5.13) \quad x_\gamma(z)^{3k+1} = 0 \text{ on } L(\Lambda).$$

Proof. Part (i) is contained in Theorem 5.9. Part (ii) follows from Theorem 4.3, Corollary 5.4, (5.11) and part (i).

(iii) By (ii)

$$Y(u, z) = 0 \text{ on } L(\Lambda)$$

for each $u \in N^1(k\Lambda_0)$, so it is enough to show that $x_\beta(-1)^{2k+1}\mathbf{1} \in N^1(k\Lambda_0)$. Since $x_\beta(-1)$ is a root vector of a real root we have $x_\beta(-1)^n v_{k\Lambda_0} = 0$ in $L(k\Lambda_0)$ for n sufficiently large [K] (here $v_{k\Lambda_0}$ denotes a highest weight vector in $L(k\Lambda_0)$). From $x_{-\beta}(1)v_{k\Lambda_0} = 0$, $[x_{-\beta}(1), x_\beta(-1)] = h_\beta + 2c$, $(h_\beta + 2c)v_{k\Lambda_0} = 2kv_{k\Lambda_0}$ it follows from the representation theory of $\mathfrak{sl}(2, \mathbb{F})$ that we must have $x_\beta(-1)^{2k+1}v_{k\Lambda_0} = 0$ in $L(k\Lambda_0)$ so that (in $N(k\Lambda_0)$)

$$x_\beta(-1)^{2k+1}\mathbf{1} \in N^1(k\Lambda_0).$$

Part (iv) is proved similarly. $\square$

Theorem 5.14. *Let $\mathfrak{g}$ be a finite dimensional split simple Lie algebra over $\mathbb{F}$ and let Λ be dominant integral with $\Lambda(c) = k$. Then*

$$(5.14) \quad \bar{R}M(\Lambda) = M^1(\Lambda)$$

where $M^1(\Lambda)$ denotes the maximal submodule of the Verma module $M(\Lambda)$.

Proof. Set $V = \bar{R}M(\Lambda)$. By Theorem 5.9 $V \subset M^1(\Lambda)$. By Proposition 5.3 it is enough to show that

$$\{f_i^{\Lambda(h_i)+1}v_\Lambda \mid i = 0, \dots, \ell\} \subset V,$$

with v_Λ a highest weight vector in $M(\Lambda)$. Let α_i be a long simple root. Then $x_{-\alpha_i}(-1)^{k+1}\mathbf{1} \in R$ (e.g. by using the action of the Weyl group of $\mathfrak{g}$) and it follows that

$$x_{-\alpha_i}(z)^{k+1}v_\Lambda \in V[[z, z^{-1}]].$$

In particular

$$f_i^{k+1}v_\Lambda = x_{-\alpha_i}(0)^{k+1}v_\Lambda = \text{coeff}_{z^{-k-1}} x_{-\alpha_i}(z)^{k+1}v_\Lambda \in V,$$

and now the representation theory of $\mathfrak{sl}(2, \mathbb{F})$ implies that $f_i^{\Lambda(h_i)+1}v_\Lambda \in V$. If α_j is a short simple root

$$x_{-\alpha_j}(-1)^{nk+1}\mathbf{1} \in N^1(k\Lambda_0) = U(\hat{\mathfrak{g}})x_\theta(-1)^{k+1}\mathbf{1},$$

for $n = 2$ or 3 by the argument of the proof of Proposition 5.13(iii). It follows from (5.11) and induction that

$$x_{-\alpha_j}(z)^{nk+1}v_\Lambda \in V[[z, z^{-1}]]$$

so that (coefficient of z^{-nk-1}) $f_j^{nk+1}v_\Lambda \in V$ and hence $f_j^{\Lambda(h_j)+1}v_\Lambda \in V$. Finally,

$$f_0^{k+1}v_\Lambda = x_\theta(-1)^{k+1}v_\Lambda = \text{coeff}_{z^0} x_\theta(z)^{k+1}v_\Lambda \in V$$

and thus $f_0^{\Lambda(h_0)+1}v_\Lambda \in V$. $\square$

Theorem 5.15. *Let V be a highest weight module of level k . The following are equivalent:*

- (a) V is a standard module,
- (b) $\bar{R}$ annihilates V .

Proof. By Theorem 5.9 (a) implies (b). Conversely, assume that $\bar{R}V = 0$, let Λ be the highest weight of V and v_Λ a highest weight vector. Then the same reasoning as in the proof of Theorem 5.14 shows that for each $i = 0, 1, \dots, \ell$, $f_i^{n_k+1}v_\Lambda \in \bar{R}V = 0$ for $n = 1, 2$ or 3 . Thus V is a standard module. $\square$

Remark. As was mentioned in the introduction, Theorem 5.9 may be viewed as a consequence of the Frenkel-Kac vertex operator formula (cf. [FK]). We omitted such a proof in order to avoid additional notation that would not be needed later. Besides that, the ideas used in the above proof (Lemma 5.8 in particular) work for some nonintegrable representations of affine Lie algebras, where there is no (known) vertex operator formula analogous to the integrable case. Similar ideas have been used in some other situations, see for example [FNO].

Theorem 5.14 was inspired by [MP, Theorem 9.28] in the principal picture, except that here it is a consequence of the Weyl-Kac character formula. Moreover, the Kac-Wakimoto character formula can be used in a similar way for some modular representations (cf. [Ad]).

Theorems 5.9 and 5.14 put together give Theorem 5.15, which in terms of the vertex operator algebra $L(k\Lambda_0)$ states that the level k standard $\mathfrak{g}$ -modules are its only irreducible highest weight modules. Results of this type have been proved by different methods in [FZ], [DL] and [Li].

Theorem 5.1 and Proposition 5.2 are similar to some results in [H] at the critical level.

6. COLORED PARTITIONS, LEADING TERMS AND THE MAIN RESULTS

In Sections 6–10 we specialize to the case $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{F})$ and level $k \in \mathbb{N}$.

6.1. Colored partitions.

Let A be a nonempty set and denote by $\mathcal{P}(A)$ the set of all maps $\pi : A \rightarrow \mathbb{N}$, where $\pi(a)$ equals zero for all but finitely many $a \in A$. Clearly π is determined by its values $(\pi(a) \mid a \in A)$ and we shall also write π as a monomial

$$(6.1.1) \quad \pi = \prod_{a \in A} a^{\pi(a)}.$$

We shall say that π is a *partition* and for $\pi(a) > 0$ we shall say that a is a *part* of π . We define the *length* $\ell(\pi)$ of π by

$$\ell(\pi) = \sum_{a \in A} \pi(a).$$

For $\rho, \pi \in \mathcal{P}(A)$ we write $\rho \subset \pi$ (or $\pi \supset \rho$) if $\rho(a) \leq \pi(a)$ for all $a \in A$ and we say that ρ is *contained* in π . For $\rho, \pi \in \mathcal{P}(A)$ we define $\pi\rho$ in $\mathcal{P}(A)$ by $(\pi\rho)(a) = \pi(a) + \rho(a)$, $a \in A$. If $\rho \subset \pi$, then we define π/ρ in $\mathcal{P}(A)$ by $(\pi/\rho)(a) = \pi(a) - \rho(a)$, $a \in A$. Clearly $\rho(\pi/\rho) = \pi$.

We also define $\pi \cup \nu$ and $\pi \cap \nu$ by

$$\begin{aligned}(\pi \cup \nu)(a) &= \max\{\pi(a), \nu(a)\}, \\(\pi \cap \nu)(a) &= \min\{\pi(a), \nu(a)\}.\end{aligned}$$

We shall say that $1 = \prod_{a \in A} a^0$ is the partition with no parts and length 0 (and we shall sometimes denote it as $\emptyset$). Clearly $\mathcal{P}(A)$ is a monoid, graded by length. We shall usually use the multiplicative notation (6.1.1). We consider elements of A as partitions of length 1, i.e. $A \subset \mathcal{P}(A)$.

For lack of a better terminology, we shall say that $\mathcal{I} \subset \mathcal{P}(A)$ is an *ideal* in the monoid $\mathcal{P}(A)$ if $\rho \in \mathcal{I}$ and $\pi \in \mathcal{P}(A)$ implies $\rho\pi \in \mathcal{I}$. This should not be confused with a notion of partition ideal which we shall use in Section 11.

In the case when A is $\mathbb{Z}, \mathbb{Z}_{<0}$ or $\mathbb{Z}_{\leq 0}$ we shall call elements of $\mathcal{P}(A)$ *plain partitions*. For a plain partition $\pi \in \mathcal{P}(\mathbb{Z})$ we define the *degree* $|\pi|$ of π by

$$|\pi| = \sum_{n \in \mathbb{Z}} n \pi(n),$$

and we say that a part n of π has *degree* n .

Set $x = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, $h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and $y = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, so that $B = \{x, h, y\}$ is the usual basis of $\mathfrak{g}$. Set

$$\begin{aligned}\bar{B} &= \{b(n) \mid b \in B, n \in \mathbb{Z}\}, \\ \bar{B}_- &= \bar{B}_{<0} \cup \{y(0)\}, \\ \bar{B}_{<0} &= \{b(n) \mid b \in B, n \in \mathbb{Z}_{<0}\},\end{aligned}$$

so that $\bar{B}, \bar{B}_-, \bar{B}_{<0}$ parameterize bases of Lie algebras $\hat{\mathfrak{g}}/\mathbb{F}c$, $\tilde{\mathfrak{n}}_-$ and $\tilde{\mathfrak{g}}_{<0}$ (respectively). We shall say that elements of $\mathcal{P}(\bar{B})$, $\mathcal{P}(\bar{B}_-)$ or $\mathcal{P}(\bar{B}_{<0})$ are *colored partitions*. For an element $a = b(n)$ in $\bar{B}$ we say that it is of *degree* $|a| = n$, *color* b and *h-weight* $\text{wt}(a) = \text{wt}(b)$, where $\text{wt}(x) = \alpha$, $\text{wt}(h) = 0$ and $\text{wt}(y) = -\alpha$. For $\pi \in \mathcal{P}(\bar{B})$ we define the *degree* $|\pi|$ and *weight* $\text{wt}(\pi)$ by

$$|\pi| = \sum_{a \in \bar{B}} \pi(a) |a|, \quad \text{wt}(\pi) = \sum_{a \in \bar{B}} \pi(a) \text{wt}(a).$$

We also define the *shape* $\text{sh}(\pi)$ of π as the plain partition

$$\text{sh}(\pi) = \prod_{a \in \bar{B}} |a|^{\pi(a)}.$$

6.2. An order on colored partitions.

We choose the order $\prec$ on B defined as $y \prec h \prec x$, and the order $\prec$ on $\bar{B}$ defined by

$$(6.2.1) \quad b_1(j_1) \prec b_2(j_2) \quad \text{iff} \quad j_1 < j_2 \quad \text{or} \quad j_1 = j_2, \quad b_1 \prec b_2.$$

Then we may think of the colored partition $\pi \in \mathcal{P}(\bar{B})$ of the form $\pi = \prod b_i(j_i)$ as

$$(6.2.2) \quad \pi = (b_1(j_1), \dots, b_s(j_s)), \quad b_1(j_1) \prec \dots \prec b_s(j_s),$$

where $b_i(j_i) \in \bar{B}$ are parts of π and $s \geq 0$ is the length of π .

We may visualize a colored partition $\pi \in \mathcal{P}(\bar{B}_{<0})$ by its Young diagram consisting of s rows of boxes: in the first row $-j_1$ boxes of color b_1 , in the second row $-j_2$ boxes of color b_2 , etc. Similarly for the shape of π — just without colors:

$$\begin{array}{llll} \text{color} & b_1 & \square\square\cdots\square\cdots\square\square & -j_1 \text{ boxes} \\ & b_2 & \square\square\cdots\square\cdots\square & -j_2 \\ & & \cdots & \\ & b_s & \square\square\cdots\square & -j_s . \end{array}$$

Let $\pi = (a_1, \dots, a_s)$, $\pi' = (a'_1, \dots, a'_{s'})$, $\pi, \pi' \in \mathcal{P}(\bar{B})$ (i.e. π and π' are written in the form (6.2.2)). We extend the order $\preccurlyeq$ on $\bar{B} \subset \mathcal{P}(\bar{B})$ defined by (6.2.1) to the order on $\mathcal{P}(\bar{B})$ defined by

$$\pi \prec \pi'$$

if $\pi \neq \pi'$ and one of the following statements hold:

- (i) $\ell(\pi) > \ell(\pi')$,
- (ii) $\ell(\pi) = \ell(\pi')$, $|\pi| < |\pi'|$,
- (iii) $\ell(\pi) = \ell(\pi')$, $|\pi| = |\pi'|$ and there is i , $\ell(\pi) \geq i \geq 1$, such that $|a_j| = |a'_j|$ for $\ell(\pi) \geq j > i$ and $|a_i| < |a'_i|$,
- (iv) $\text{sh } \pi = \text{sh } \pi'$ and there is i , $\ell(\pi) \geq i \geq 1$, such that $a_j = a'_j$ for $\ell(\pi) \geq j > i$ and $a_i \prec a'_i$.

We also order plain partitions $\mathcal{P}(\mathbb{Z})$ by requirements (i)–(iii). So if (i), (ii) or (iii) holds, we have that $\text{sh } \pi \prec \text{sh } \pi'$ and (hence) $\pi \prec \pi'$.

As an example we may take

$$y(-2)y(-2)x(-2) \prec y(-3)x(-2)y(-1) \prec y(-3)y(-2)x(-1) \prec x(-4)y(-1)y(-1),$$

all partitions being of degree -6 and weight $-\alpha$.

Remark. In what follows all notions will depend on the order chosen. The same results may hold for a different choice of order as well. For example, if we define $\pi < \pi'$ by $\pi \neq \pi'$ and

- (i) $\ell(\pi) > \ell(\pi')$,
- (ii) $\ell(\pi) = \ell(\pi')$, $|\pi| < |\pi'|$,
- (iii) $\ell(\pi) = \ell(\pi')$, $|\pi| = |\pi'|$, and there is i , $\ell(\pi) \geq i \geq 1$, such that $a_j = a'_j$ for $\ell(\pi) \geq j > i$ and $a_i \prec a'_i$,

then all the statements in this section hold. However, some proofs in later sections cannot be applied as they stand. Note that with this order we have

$$y(-2)y(-2)x(-2) < y(-3)x(-2)y(-1) < x(-4)y(-1)y(-1) < y(-3)y(-2)x(-1).$$

Lemma 6.2.1. (i) The relation $\preccurlyeq$ is a linear order on $\mathcal{P}(\bar{B})$. The element 1 is the largest element of $\mathcal{P}(\bar{B})$.

(ii) Let $\ell \geq 0$, $n \in \mathbb{Z}$ and let $S \subset \mathcal{P}(\bar{B})$ be a nonempty subset such that all π in S have length $\ell(\pi) \leq \ell$ and degree $|\pi| \geq n$. Then S has a minimal element.

(iii) The relation $\preccurlyeq$ is a (reverse) well order on $\mathcal{P}(\bar{B}_-)$.

Lemma 6.2.2. *Let $\mu, \nu, \pi \in \mathcal{P}(\bar{B})$. Let $\mu \preccurlyeq \nu$. Then $\pi\mu \preccurlyeq \pi\nu$.*

6.3. Filtrations on highest weight modules and leading terms.

For $\pi \in \mathcal{P}(\bar{B})$ of the form (6.2.2) set

$$u(\pi) = b_1(j_1) \dots b_s(j_s) \in U_k(\tilde{\mathfrak{g}}).$$

This defines a map $u : \mathcal{P}(\bar{B}) \rightarrow U_k(\tilde{\mathfrak{g}})$. For a highest weight $\tilde{\mathfrak{g}}$ -module V with a highest weight vector v_0 and for $\pi \in \mathcal{P}(\bar{B}_-)$ define

$$\begin{aligned} V_{[\pi]} &= \mathbb{F}\text{-span}\{u(\pi')v_0 \mid \pi' \in \mathcal{P}(\bar{B}_-), \pi' \succcurlyeq \pi\}, \\ V_{(\pi)} &= \mathbb{F}\text{-span}\{u(\pi')v_0 \mid \pi' \in \mathcal{P}(\bar{B}_-), \pi' \succ \pi\} \\ &= \bigcup_{\pi' \succ \pi} V_{[\pi']}. \end{aligned}$$

Clearly we have

$$(6.3.1) \quad V_{[\pi]} \supset V_{(\pi)} \supset V_{[\pi']} \supset V_{(\pi')} \quad \text{for } \pi \prec \pi',$$

$$(6.3.2) \quad \bigcup_{\pi \in \mathcal{P}(\bar{B}_-)} V_{[\pi]} = V, \quad \bigcap_{\pi \in \mathcal{P}(\bar{B}_-)} V_{(\pi)} = 0, \quad \dim V_{[\pi]}/V_{(\pi)} \leq 1,$$

$$u(\pi')V_{[\pi]} \subset V_{[\pi'\pi]}, \quad u(\pi')V_{(\pi)} \subset V_{(\pi'\pi)}.$$

Moreover, if $\varphi : V \rightarrow V'$ is a surjective homomorphism of highest weight $\tilde{\mathfrak{g}}$ -modules, then

$$\varphi(V_{[\pi]}) = V'_{[\pi]} \quad , \quad \varphi(V_{(\pi)}) = V'_{(\pi)}.$$

For $v \in V \setminus \{0\}$ define the *leading term* $\ell t(v) = \ell t_V(v) \in \mathcal{P}(\bar{B}_-)$ by

$$\ell t(v) = \pi \quad \text{iff} \quad v \in V_{[\pi]} \setminus V_{(\pi)}.$$

Because of (6.3.1), (6.3.2) and Lemma 6.2.1(iii) the leading term $\ell t(v)$ is well defined for every $v \neq 0$. For a subset $S \subset V$ set

$$\ell t(S) = \{\ell t(v) \mid v \in S \setminus \{0\}\}.$$

Note that $\ell t(M(\Lambda)) = \mathcal{P}(\bar{B}_-)$, $\ell t(N(k\Lambda_0)) = \mathcal{P}(\bar{B}_{<0})$. Clearly we have

Proposition 6.3.1. (i) *For $\pi \in \mathcal{P}(\bar{B}_-)$ and $v \in M(\Lambda)$ we have $\ell t(u(\pi)v) = \pi \ell t(v)$. In particular, for a $\tilde{\mathfrak{g}}$ -submodule $V' \subset M(\Lambda)$ we have that $\ell t_{M(\Lambda)}(V')$ is an ideal in the monoid $\mathcal{P}(\bar{B}_-)$.*

(ii) *For $\pi \in \mathcal{P}(\bar{B}_{<0})$ and $v \in N(k\Lambda_0)$ we have $\ell t(u(\pi)v) = \pi \ell t(v)$. In particular, for a $\tilde{\mathfrak{g}}$ -submodule $V' \subset N(k\Lambda_0)$ we have that $\ell t_{N(k\Lambda_0)}(V')$ is an ideal in the monoid $\mathcal{P}(\bar{B}_{<0})$.*

Proposition 6.3.2. *If V' is a $\tilde{\mathfrak{g}}$ -submodule of a highest weight $\tilde{\mathfrak{g}}$ -module V , then*

$$\ell t(V/V') = \ell t(V) \setminus \ell t_V(V').$$

Proposition 6.3.3. *For an $\mathfrak{h}^e$ -invariant subspace $V' \subset V$ and $\mu \in \mathfrak{h}^{e*}$ we have*

$$\dim V'_\mu = \#\{\pi \in \ell_V(V') \mid \text{wt}(\pi) + |\pi|\delta = \mu\}.$$

As usual (cf. [K]), here δ denotes the imaginary root related to the homogeneous grading of $\tilde{\mathfrak{g}}$.

Remark. For a highest weight module V with highest weight Λ we have $V = M(\Lambda)/M$ for some submodule M of the Verma module $M(\Lambda)$. By Propositions 6.3.1–3 we can write a character formula

$$(6.3.3) \quad e^{-\Lambda} \text{ch } V = \sum_{\pi \in \mathcal{P}(\bar{B}_-) \setminus \ell_{M(\Lambda)}(M)} e^{\text{wt}(\pi) + |\pi|\delta}.$$

The idea to parameterize bases of standard modules by classes of partitions originated in the work [LW] on a Lie theoretic interpretation of the Rogers-Ramanujan identities.

Proposition 6.3.4. *Let V be a highest weight $\tilde{\mathfrak{g}}$ -module. Then*

- (i) $\{u(\pi)v_0 \mid \pi \in \ell(V)\}$ is a basis of V .
- (ii) For a linear subspace $V' \subset V$ and a map $u': \ell_V(V') \rightarrow V'$ such that $\ell_V(u'(\pi)) = \pi$ the set

$$\{u'(\pi) \mid \pi \in \ell_V(V')\}$$

is a basis of V' .

6.4. A filtration on a completion of the enveloping algebra and leading terms.

Recall that the category $\mathcal{O}$ consists of $\tilde{\mathfrak{g}}$ -modules V such that V is a direct sum of finite dimensional weight spaces and such that the set of weights of V is contained in a finite union of sets of the form $\lambda - \sum \mathbb{N}\alpha_i$ (cf. [K]).

Recall that we denote by $\overline{U}_k(\tilde{\mathfrak{g}})$ the completed enveloping algebra such that $x_i \rightarrow x$ if for each $\tilde{\mathfrak{g}}$ -module V in the category $\mathcal{O}$ and each vector v in V there is i_0 such that $i \geq i_0$ implies $x_i v = x_{i_0} v = xv$.

Lemma 6.4.1. *Let $\pi \in \mathcal{P}(\bar{B})$. Then there exist a $\tilde{\mathfrak{g}}$ -module M of level k in the category $\mathcal{O}$, a vector $v_\pi \in M$ and a functional $v_\pi^* \in M^*$ such that*

$$\begin{aligned} \langle v_\pi^*, u(\pi)v_\pi \rangle &\neq 0, \\ \langle v_\pi^*, u(\pi')v_\pi \rangle &= 0 \quad \text{if } \pi' \in \mathcal{P}(\bar{B}), \pi' \neq \pi, \ell(\pi') \leq \ell(\pi). \end{aligned}$$

Proof. Let $\pi = \pi_- \pi_0 \pi_+$, where $\pi_- \in \mathcal{P}(\bar{B}_-)$, $\pi_0 = h^q$ for some $q \geq 0$ and the parts of π_+ are in $\tilde{\mathfrak{n}}_+$. For an element $b \in \bar{B}_-$ denote by $\bar{b}$ the element in $\bar{B} \cap \tilde{\mathfrak{n}}_+$ such that

$$[\bar{b}, b] \in (\mathbb{F}h + \mathbb{F}c) \setminus \{0\}.$$

Let $p = \ell(\pi_-)$, $r = \ell(\pi_+)$, $\ell = \ell(\pi) = p + q + r$ and let M be a level k module of the form

$$M = (\otimes_{i=1}^p M(0)) \otimes (\oplus_{i=1}^{q+1} M(\lambda_i)) \otimes (\otimes_{i=1}^r M(\mu_i)),$$

where $\lambda_i, \mu_i \in \mathfrak{h}^{e*}$ are chosen so that $\lambda_i(h) \neq \lambda_j(h)$ for $i \neq j$, $\mu_i([\bar{b}_i, b_i]) > 0$ (i.e. positive rational numbers) for all $i = 1, \dots, r$, and where $b_i \in \bar{B}_-$ are such that

$$(6.4.1) \quad \pi_+ = (\bar{b}_1, \dots, \bar{b}_r), \quad \bar{b}_1 \preccurlyeq \dots \preccurlyeq \bar{b}_r.$$

Let as usual $v_\Lambda \in M(\Lambda)$ denote a highest weight vector in a Verma module $M(\Lambda)$. Then the vectors

$$w_i = \lambda_1(h)^i v_{\lambda_1} + \dots + \lambda_{q+1}(h)^i v_{\lambda_{q+1}}, \quad i = 0, 1, \dots, q$$

form a basis of $\mathbb{F}\text{-span}\{v_{\lambda_1}, \dots, v_{\lambda_{q+1}}\}$ and

$$\{u(\rho)w_i \mid \rho \in \mathcal{P}(\bar{B}_-), i = 0, \dots, q\}$$

is a basis of $\oplus_{i=1}^{q+1} M(\lambda_i)$. For other Verma modules $M(\Lambda)$ choose the basis $u(\rho)v_\Lambda$ $\rho \in \mathcal{P}(\bar{B}_-)$, and for M the corresponding tensor products. Note that

$$h^i w_0 = w_i \quad \text{for } i = 0, 1, \dots, q.$$

Let v_π be a basis vector

$$v_\pi = (\otimes_{i=1}^p v_0) \otimes w_0 \otimes (\otimes_{i=1}^r b_i v_{\mu_i})$$

where $b_i \in \bar{B}_-$ are defined by (6.4.1).

Let v_π^* be the element of dual basis corresponding to the vector

$$v = (\otimes_{i=1}^p d_i v_0) \otimes h^q w_0 \otimes (\otimes_{i=1}^r v_{\mu_i}),$$

where $d_i \in \bar{B}_-$ are parts of $\pi_- = (d_1, \dots, d_p)$, $d_1 \preccurlyeq \dots \preccurlyeq d_p$. It is clear from our choice that

$$v_\pi^*(u(\pi)v_\pi) > 0.$$

Now let $\pi' \in \mathcal{P}(\bar{B})$, $\ell(\pi') \leq \ell$, and assume that $u(\pi')v_\pi$ has a nontrivial component along the basis vector v . Write $\pi' = \pi'_- \pi'_0 \pi'_+$, notation being as before. In order to have a component $cv = c(\otimes_{i=1}^p d_i v_0) \otimes h^q w_0 \otimes (\otimes_{i=1}^r v_{\mu_i})$, $c \neq 0$, π'_- must contain all parts d_i of π and $\ell(\pi'_0)$ must be at least q . Hence $\ell(\pi'_+) \leq r$. But to have $u(\pi'_+)(\otimes_{i=1}^r b_i v_{\mu_i}) = c_1(\otimes_{i=1}^r v_{\mu_i}) + \dots$, $c_1 \neq 0$, we need $\ell(\pi'_+) \geq r$. Hence $\ell(\pi'_-) = r$, $\pi'_0 = \pi_0$, $\ell(\pi'_+) = r$. But then $\pi'_- = \pi_-$ and $\pi'_+ = \pi_+$, i.e. $\pi' = \pi$. $\square$

Remark. The above proof is a slight refinement of the proof of Lemma 4.1 in [MP].

For $\pi \in \mathcal{P}(\bar{B})$ set

$$U_{[\pi]} = \overline{\mathbb{F}\text{-span}\{u(\pi') \mid \pi' \succcurlyeq \pi\}},$$

$$U_{(\pi)} = \overline{\mathbb{F}\text{-span}\{u(\pi') \mid \pi' \succ \pi\}},$$

the closure taken in $\overline{U_k(\tilde{\mathfrak{g}})}$.

Lemma 6.4.2. For $\pi \in \mathcal{P}(\bar{B})$ we have $U_{[\pi]} = \mathbb{F}u(\pi) + U_{(\pi)}$. Moreover,

$$\dim U_{[\pi]}/U_{(\pi)} = 1.$$

Proof. Clearly $\mathbb{F}u(\pi) + U_{(\pi)} \subset U_{[\pi]}$. We shall prove the converse inclusion as well: Let $u = \lim_{i \in I} u_i \in U_{[\pi]}$. Choose $v_\pi \in M$ and $v_\pi^* \in M^*$ as in Lemma 6.4.1. Then for some $i_0 \in I$ we have

$$\begin{aligned} u_i v_\pi &= u_{i_0} v_\pi \quad \text{for } i \geq i_0, \\ v_\pi^*(u_i v_\pi) &= v_\pi^*(u_{i_0} v_\pi) = c \quad \text{for } i \geq i_0. \end{aligned}$$

Since

$$u_i = c_i u(\pi) + \sum_{\pi' \succ \pi} c_{\pi'}^i u(\pi'),$$

by Lemma 6.4.1 we have $c_i = c$ for $i \geq i_0$. Hence

$$u'_i = u_i - cu(\pi) \in U_{(\pi)} \quad \text{for } i \geq i_0,$$

the limit $\lim_{i \in I} u'_i$ exists and $u - cu(\pi) \in U_{(\pi)}$. Now $U_{[\pi]} = \mathbb{F}u(\pi) + U_{(\pi)}$ implies $\dim U_{[\pi]}/U_{(\pi)} \leq 1$. By Lemma 6.4.1 $\langle v_\pi^*, U_{(\pi)} v_\pi \rangle = 0$, $\langle v_\pi^*, U_{[\pi]} v_\pi \rangle \neq 0$, and the lemma follows. $\square$

Set

$$U_k(\tilde{\mathfrak{g}})_{\text{loc}} = \mathbb{F}\text{-span}\{v_n \mid v \in N(k\Lambda_0), n \in \mathbb{Z}\},$$

where v_n denotes a coefficient in $Y(v, z)$. From the commutator formula (2.15) we see that $U_k(\tilde{\mathfrak{g}})_{\text{loc}}$ is a Lie algebra. From the normal order product formula (2.14) we see that $U_k(\tilde{\mathfrak{g}})_{\text{loc}} \subset \bigcup_{\pi \in \mathcal{P}(\bar{B})} U_{[\pi]}$. Let us denote by $U'(U_k(\tilde{\mathfrak{g}})_{\text{loc}})$ the associative subalgebra of $\overline{U_k(\tilde{\mathfrak{g}})}$ generated by $U_k(\tilde{\mathfrak{g}})_{\text{loc}}$. Then we have:

$$U'(U_k(\tilde{\mathfrak{g}})_{\text{loc}}) \subset \bigcup_{\pi \in \mathcal{P}(\bar{B})} U_{[\pi]}.$$

For $u \in U_{[\pi]}$, $u \notin U_{(\pi)}$ we define the *leading term*

$$\ell(u) = \pi.$$

Remark. From Lemma 6.4.1 we see that if the leading term $\ell(u)$ of u in $U'(U_k(\tilde{\mathfrak{g}})_{\text{loc}})$ exists, it is unique.

Proposition 6.4.3. Every element $u \in U'(U_k(\tilde{\mathfrak{g}})_{\text{loc}})$, $u \neq 0$, has a unique leading term $\ell(u)$. Moreover, for any highest weight module V we have that

$$uV_{[\pi]} \subset V_{[\pi \ell(u)]}, \quad uV_{(\pi)} \subset V_{(\pi \ell(u))}.$$

Proof. We should prove that each element in $A = U'(U_k(\tilde{\mathfrak{g}})_{\text{loc}})$ has a leading term.

First note that A is graded by degree, say $A = \coprod_{n \in \mathbb{Z}} A_n$.

Recall that by the definition of convergence in $\overline{U_k(\tilde{\mathfrak{g}})}$ we have $x_i \rightarrow x$ if for each module V in the category $\mathcal{O}$ and each vector v in V there is i_0 such that

$i \geq i_0$ implies $x_i v = x_{i_0} v = x v$. In this proof we shall use the notion of *uniform convergence*: we say that $x_i \rightarrow x$ uniformly if for each $N \in \mathbb{N}$ there is i_0 such that for each module V in the category $\mathcal{O}$ and each v in V , $|\deg v| < N$, $i \geq i_0$ implies $x_i v = x_{i_0} v = x v$. Here by $|\deg v| < N$ we mean that for a weight vector $v \in V_\mu$ the degree of any maximal weight of V differs from the degree of μ by at most N .

From the normal order product formula (2.13) we see by induction that each $x \in A_n$ is a limit of a sequence $(x_i)_{i \in \mathbb{N}}$ such that $x_i \in U_k(\tilde{\mathfrak{g}})_{\ell, n}$ (where ℓ denotes a filtration and n a degree) and that $x_i \rightarrow x$ uniformly.

Set $\mathcal{P}_{\ell, n} = \{\pi \mid \ell(\pi) = \ell, |\pi| = n\}$. Then $\mathcal{P}_{\ell_1, n} \prec \mathcal{P}_{\ell_2, n}$ for $\ell_1 > \ell_2$, and each $\mathcal{P}_{\ell, n} \cong \mathbb{N}$ as an ordered set.

Now we show in three steps that $x \in A_n \cap U_{[\pi]}$ is the uniform limit of a sequence (x_i) , $x_i \in U_k(\tilde{\mathfrak{g}})_{[\pi]} = \mathbb{F}\text{-span}\{u(\pi') \mid \pi' \succ \pi\}$, $\deg x_i = n$:

(1) Let π_0 be the smallest element in $\mathcal{P}_{\ell, n}$, $\ell = \ell(\pi)$, $n = |\pi|$. We (may) assume that $x \in U_{[\pi]} \cap A_n$ is a uniform limit of a sequence (x_i) , $x_i \in U_k(\tilde{\mathfrak{g}})_{\ell, n}$. Note that the segment $[\pi_0, \pi]$ is finite, and write

$$x_i = \sum_{\pi_0 \prec \kappa \prec \pi} C_{\kappa, i} u(\kappa) + \sum_{\nu \succ \pi} C_{\nu, i} u(\nu).$$

For $\kappa \prec \pi$ take v_κ, v_κ^* as in Lemma 6.4.1. Then $x \in U_{[\pi]}$ implies $v_\kappa^*(x v_\kappa) = 0$. Now $x_i \rightarrow x$ implies that there is i_0 such that $x_i v_\kappa = x_{i_0} v_\kappa = x v_\kappa$ for $i \geq i_0$, and hence $C_{\kappa, i} = 0$ for $i \geq i_0$. So in a finite number of steps we see that for some i_0 the sequence $(x_i)_{i \geq i_0}$, $x_i \in U_k(\tilde{\mathfrak{g}})_{[\pi]}$, converges uniformly to x .

(2) Let $\ell = \ell(\pi) = \ell(\pi')$, $n = |\pi| = |\pi'|$, $\pi \prec \pi'$ and $x \in U_{[\pi]}$, $x \notin U_{(\pi')}$. Since the interval $[\pi, \pi']$ is finite, there is $\pi \prec \tau \prec \pi'$ such that $x \in U_{[\tau]} \setminus U_{(\tau)}$, i.e. u has a leading term.

(3) Let $x \in A_n$, $x \in U_{[\pi]}$ for all π such that $\ell(\pi) = \ell$. For each π we may choose a sequence (x_i^π) in $U_k(\tilde{\mathfrak{g}})_{[\pi]}$ such that $x_i^\pi \rightarrow x$ uniformly. Let

$$\begin{aligned} x_i^\pi &= \sum_{\ell(\kappa) = \ell} C_{\kappa, i}^\pi u(\kappa) + x_{i, \ell-1}^\pi \\ &= x_{i, \ell}^\pi + x_{i, \ell-1}^\pi, \end{aligned}$$

where $x_{i, \ell-1}^\pi \in U_k(\tilde{\mathfrak{g}})_{\ell-1, n}$. For a sequence $\pi^{(1)} \prec \pi^{(2)} \prec \dots$ such that

$$U_{[\pi^{(1)}]} \supset U_{[\pi^{(2)}]} \supset \dots \supset \bigcap_{\ell(\pi) = \ell} U_{[\pi]} \cap A_n = \bigcup_{\ell(\pi) < \ell} U_{[\pi]} \cap A_n$$

we can construct a diagonal sequence

$$x_{i_k}^{\pi^{(k)}} \quad , \quad k = 1, 2, \dots$$

such that (a) i_k is such that for each module V and each vector v in V , $|\deg v| < k$, the inequality $i \geq i_k$ implies $x_i^{\pi^{(k)}} v = x_{i_k}^{\pi^{(k)}} v = x v$, and (b) $k < k'$ implies $i_k < i_{k'}$. Then $x_{i_k}^{\pi^{(k)}} \rightarrow x$ uniformly. Since $x_{i_k}^{\pi^{(k)}} \in U_{[\pi^{(k)}]}$, we have $x_{i_k, \ell}^{\pi^{(k)}} \rightarrow 0$ and we may take a subsequence $x_{i_{k_m}, \ell}^{\pi^{(k_m)}}$ which converges to 0 uniformly. Hence we have constructed a sequence

$$x_m = x_{i_{k_m}, \ell-1}^{\pi^{(k_m)}} \in \bigcup_{\ell(\kappa) < \ell} U_{[\kappa]} \cap A_n$$

such that $x_m \rightarrow x$ uniformly.

Now in a finite number of steps we see that either $x = 0$ or x has a leading term. $\square$

Remark. The above proof shows the following: Let $U_k(\tilde{\mathfrak{g}}) \subset A \subset \overline{U_k(\tilde{\mathfrak{g}})}$ be a graded subspace. Then every x in A has a leading term if every x in A is a uniform limit of a sequence in $U_k(\tilde{\mathfrak{g}})_\ell$ for some fixed ℓ (depending on x).

6.5. The main results.

In the remainder of this section we shall assume that Λ is dominant integral and that $k = \Lambda(c)$.

First let us recall the construction of the loop module $\bar{R}$: Since we assume that $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{F})$, we have $\theta = \alpha$ if we write

$$x = x_\alpha = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad y = x_{-\alpha} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad h = \alpha^\vee = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Then R defined by (5.4), i.e.

$$R = U(\mathfrak{g})x_\alpha(-1)^{k+1}\mathbf{1} \subset N(k\Lambda_0),$$

is a $(2k+3)$ -dimensional $\mathfrak{sl}(2, \mathbb{F})$ -module and $\bar{R}$ is defined as a linear span of the coefficients of the fields

$$Y(r, z) = \sum_{n \in \mathbb{Z}} r(n)z^{-n-k-1}$$

for $r \in R$. As it has been remarked, these coefficients $r(n)$ are infinite sums of elements in $U_k(\tilde{\mathfrak{g}})$, but are well defined operators on every highest weight $\tilde{\mathfrak{g}}$ -module V of level k .

Note that every vector $v \neq 0$ in a Verma $\tilde{\mathfrak{g}}$ -module $M(\Lambda)$ can be written as a linear combination of terms $u(\pi)v_\Lambda$, where

$$u(\pi) = b_1(j_1) \dots b_s(j_s) \in U_k(\tilde{\mathfrak{g}}) \quad \text{for} \quad \pi = (b_1(j_1) \preccurlyeq \dots \preccurlyeq b_s(j_s)) \in \mathcal{P}(\bar{B}_-)$$

and v_Λ is a highest weight vector, and then $\ell(v)$ is just the smallest π which appears nontrivially in this linear combination. The existence of the leading term $\rho = \ell(r(n))$ is a bit more subtle: it is guaranteed by Proposition 6.4.3, but as we shall see later on, $r(n) \in \bar{R}$ can be written in the form

$$r(n) = c_\rho u(\rho) + \sum_{\pi \succ \rho} c_\pi u(\pi),$$

where $c_\rho, c_\pi \in \mathbb{C}$, $c_\rho \neq 0$, and again the leading term is the smallest colored partition which appears nontrivially in this (infinite) linear combination.

For a subset $S \subset \mathcal{P}(A)$ denote by $(S) \subset \mathcal{P}(A)$ the ideal in the monoid $\mathcal{P}(A)$ generated by S . To avoid ambiguities because of notation, let us agree with a convention that

$$\begin{aligned} \ell(\bar{R}) &\subset \ell(U'(U_k(\tilde{\mathfrak{g}})_{\text{loc}})) = \mathcal{P}(\bar{B}), \\ \ell_{M(\Lambda)}(\bar{R}v_\Lambda) &\subset \ell_{M(\Lambda)}(M(\Lambda)) = \mathcal{P}(\bar{B}_-), \\ \ell_{N(k\Lambda_0)}(\bar{R}\mathbf{1}) &\subset \ell_{N(k\Lambda_0)}(N(k\Lambda_0)) = \mathcal{P}(\bar{B}_{<0}), \end{aligned}$$

so that $(\ell(\bar{R})) \subset \mathcal{P}(\bar{B})$, $(\ell(\bar{R}v_\Lambda)) \subset \mathcal{P}(\bar{B}_-)$, $(\ell(\bar{R}\mathbf{1})) \subset \mathcal{P}(\bar{B}_{<0})$.

Let $U_0 \subset U_1 \subset U_2 \subset \dots$ be the filtration of $U_k(\tilde{\mathfrak{g}})$. Since by Theorem 5.14 $\bar{R}M(\Lambda)$ is the maximal submodule $M^1(\Lambda)$ of a Verma module $M(\Lambda)$, we have:

Lemma 6.5.1. *Let v_Λ be a highest weight vector of Verma module $M(\Lambda)$. Then*

- (i) $\ell(\bar{R}v_\Lambda) \subset \ell(U_1\bar{R}v_\Lambda) \subset \cdots \subset \ell(U_j\bar{R}v_\Lambda) \subset \cdots \subset \ell_{M(\Lambda)}(M^1(\Lambda)),$
- (ii) $(\ell(\bar{R}v_\Lambda)) \subset (\ell(U_1\bar{R}v_\Lambda)) \subset \cdots \subset (\ell(U_j\bar{R}v_\Lambda)) \subset \cdots \subset \ell_{M(\Lambda)}(M^1(\Lambda)),$
- (iii) $\bigcup_{j \geq 0} (\ell(U_j\bar{R}v_\Lambda)) = \ell_{M(\Lambda)}(M^1(\Lambda)).$

We think of elements of $U_j\bar{R}$ as relations that hold on $L(\Lambda) = M(\Lambda)/M^1(\Lambda)$. Note that (iii) shows that relations $U_j\bar{R}$, $j \geq 0$, determine the character of $L(\Lambda)$ written in the form (6.3.3), i.e

$$e^{-\Lambda} \text{ch } L(\Lambda) = \sum_{\pi \in \mathcal{P}(\bar{B}_-) \setminus \ell_{M(\Lambda)}(M^1(\Lambda))} e^{\text{wt}(\pi) + |\pi|\delta}.$$

In what follows we shall describe the ideal $\ell_{M(\Lambda)}(M^1(\Lambda))$ by explicitly constructing a basis of $M^1(\Lambda) = \bar{R}M(\Lambda)$ parameterized by colored partitions.

We shall write ℓ instead of $\ell_{M(\Lambda)}$ when no confusion can arise. Clearly there exists a map

$$\ell(\bar{R}v_\Lambda) \rightarrow \bar{R}, \quad \rho \mapsto r(\rho),$$

such that $\ell(r(\rho)v_\Lambda) = \rho$, i.e. for each colored partition $\rho \in \ell(\bar{R}v_\Lambda)$ we choose a relation $r(\rho)$ in $\bar{R}$ which will produce a vector $r(\rho)v_\Lambda$ in $M^1(\Lambda)$ with the leading term ρ . With given such a map we define

$$u(\rho \subset \pi) = u(\pi/\rho)r(\rho)$$

for $\rho \in \ell(\bar{R}v_\Lambda)$ and $\pi \in (\ell(\bar{R}v_\Lambda))$, $\rho \subset \pi$. Since our order $\preceq$ behaves well with respect to multiplication, we have $\ell(u(\rho \subset \pi)v_\Lambda) = \pi$. For each π in the ideal $(\ell(\bar{R}v_\Lambda))$ there is at least one $\rho \in \ell(\bar{R}v_\Lambda)$ such that $\rho \subset \pi$. Hence there is a map

$$(\ell(\bar{R}v_\Lambda)) \rightarrow \ell(\bar{R}v_\Lambda), \quad \pi \mapsto \rho(\pi),$$

such that $\rho(\pi) \subset \pi$.

In other words, for each colored partition π in the ideal $(\ell(\bar{R}v_\Lambda))$ we can choose $\rho = \rho(\pi)$ such that $\pi = (\pi/\rho) \cdot \rho$. Moreover, with this choice of ρ we can construct a vector $u(\rho \subset \pi)v_\Lambda = u(\pi/\rho)r(\rho)v_\Lambda$ in $M^1(\Lambda)$ with the leading term π . As we shall see later on, the main difficulty in our construction is that for a given π this can be done in several ways. For example, we may have $\pi = x(-3)x(-2)x(-1)$ and vectors $x(-3)r(x(-2)x(-1))v_\Lambda$ and $x(-1)r(x(-3)x(-2))v_\Lambda$ in $M^1(\Lambda)$ with the same leading term π . Our main result for $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{F})$ is that for each π we need only one such vector in order to obtain a basis of $M^1(\Lambda)$:

Theorem 6.5.2. *For any fixed map $(\ell(\bar{R}v_\Lambda)) \ni \pi \mapsto \rho(\pi) \in \ell(\bar{R}v_\Lambda)$ such that $\rho(\pi) \subset \pi$ the set of vectors*

$$(6.5.1) \quad u(\rho(\pi) \subset \pi)v_\Lambda, \quad \pi \in (\ell(\bar{R}v_\Lambda)),$$

is a basis of the maximal submodule $M^1(\Lambda)$ of the Verma module $M(\Lambda)$.

Remark. The above theorem is analogous to [MP, Theorem 9.26] in the principal picture. Although in [MP] a notion of the leading term was not introduced, the

general ideas are quite similar to the ones used here. At this point let us only mention that the order used in [MP] on the set of partitions $\mathcal{P}$ slightly differs from our present order on $\mathcal{P}(\mathbb{Z})$, but in both cases the minimal elements used in our arguments, in [MP] denoted by $(p;n)$, are the same. So in the present terminology Lemma 9.9 in [MP] defines the leading term $X(p;n)$ of a relation $R_p(n)$, and $X(\mu)$ in Theorem 9.26 in [MP] is analogous to $u(\rho(\pi) \subset \pi)$ for some specific choice of the map $\pi \mapsto \rho(\pi)$.

Theorem 6.5.2 will be proved in Section 10 as a consequence of Theorem 10.1, and almost all of the rest of this paper is devoted to the proof. At this point we state the following immediate consequences of Theorem 6.5.2 and Proposition 6.3.2:

Corollary 6.5.3. $\ell_{M(\Lambda)}(M^1(\Lambda)) = (\ell(\bar{R}v_\Lambda)).$

Corollary 6.5.4. $\ell(L(\Lambda)) = \mathcal{P}(\bar{B}_-) \setminus (\ell(\bar{R}v_\Lambda)).$

Theorem 6.5.5. *The set of vectors*

$$(6.5.2) \quad u(\pi)v_\Lambda, \quad \pi \in \mathcal{P}(\bar{B}_-) \setminus (\ell(\bar{R}v_\Lambda)),$$

is a basis of the standard $\mathfrak{g}$ -module $L(\Lambda)$.

Proof. Since on $L(\Lambda)$ relations $r(\rho)$, $\rho \in \ell(\bar{R})$, vanish, we see by induction that (6.5.2) is a spanning set of $L(\Lambda)$. But then by Corollary 6.5.4 and Proposition 6.3.3 it must be a basis. $\square$

Remark. Corollary 6.5.3 states that all inclusions in Lemma 6.5.1(ii) are equalities. In the case of $\mathfrak{g} = \mathfrak{sl}(3, \mathbb{F})$ and $\Lambda = \Lambda_0$ take B to be the ordered basis

$$[f_1, f_2] \prec f_1 \prec f_2 \prec h_2 \prec h_1 \prec e_2 \prec e_1 \prec [e_1, e_2],$$

where e_i, h_i, f_i denote the Chevalley generators of $\mathfrak{g}$, and let the order on $\mathcal{P}(\bar{B})$ be defined as above. Then one can see that $(\ell(\bar{R}v_{\Lambda_0})) \subsetneq (\ell(U_1 \bar{R}v_{\Lambda_0}))$, and some evidence suggest that one should expect equalities

$$(\ell(U_1 \bar{R}v_{\Lambda_0})) = \cdots = (\ell(U_j \bar{R}v_{\Lambda_0})) = \cdots = \ell_{M(\Lambda_0)}(M^1(\Lambda_0)).$$

6.6. Difference and initial conditions.

Now we will describe the sets $\ell(\bar{R})$, $\ell(\bar{R}v_\Lambda)$ and $\ell(\bar{R}\mathbf{1})$: In Section 5 we denoted by

$$R = U(\mathfrak{g})x(-1)^{k+1}\mathbf{1} \subset N(k\Lambda_0)$$

and by $\bar{R}$ the corresponding loop module. R is a $(2k+3)$ -dimensional $\mathfrak{g}$ -module with a basis $\{r_{i\alpha} \mid -k-1 \leq i \leq k+1\}$, where

$$(6.6.1) \quad r_{i\alpha} = \frac{1}{(k+1+i)!} x(0)^{k+1+i} \cdot y(-1)^{k+1}\mathbf{1}.$$

In particular $r_{-(k+1)\alpha} = y(-1)^{k+1}\mathbf{1}$, and

$$Y(r_{-(k+1)\alpha}, z) = \sum_{n \in \mathbb{Z}} r_{-(k+1)\alpha}(n) z^{-n-k-1},$$

$$r_{-(k+1)\alpha}(n) = \sum_{j_1 + \cdots + j_{k+1} = n} y(j_1) \cdots y(j_{k+1}).$$

Clearly $r_{i\alpha}(n)\mathbf{1} \neq 0$ for $n \leq -k-1$. Set $\pi_{i\alpha}(n) = \ell(r_{i\alpha}(n))$. It is also clear that for $n = (k+1)j - s$, $0 \leq s \leq k+1$,

$$(6.6.2) \quad \text{sh}(\pi_{-(k+1)\alpha}(n)) = (j-1)^s j^{k+1-s}.$$

By the adjoint action of $x(0)$ we obtain from $r_{-(k+1)\alpha}(n)$ all $r_{i\alpha}(n)$, and from (6.6.2) we see that

$$\text{sh}(\pi_{i\alpha}(n)) = (j-1)^s j^{k+1-s}.$$

It is easy to see the smallest possible colorings of the shape $(j-1)^s j^{k+1-s}$ which appears in $r_{i\alpha}(n)$: For example, for $k=4$ and the shape $(-2)^2(-1)^3$ the elements $\pi_{i\alpha}(-7)$, $i = -5, \dots, 5$, are

		$y y h h x x x x x x$
		$y h h x x x x x x x$
		$y y y y y y y h h h x$
		$y y y y y y h h h x x$
		$y y y y y h h h x x x$

In general we have:

Proposition 6.6.1. *The set $\ell(\bar{R})$ consists of elements $\pi_{m\alpha}(n) = \ell(r_{m\alpha}(n))$, $n \in \mathbb{Z}$, of the shape $(j-1)^a j^b$, $a(j-1) + bj = n$, $a+b = k+1$, $0 \leq a, b \leq k+1$,*

$$\pi_{m\alpha}(n) = \begin{cases} y(j-1)^r h(j-1)^{a-r} y(j)^b & a \geq r \geq 0, \quad m = -k-1 + a - r, \\ h(j-1)^r x(j-1)^{a-r} y(j)^b & a > r \geq 0, \quad m = -k-1 + 2a - r, \\ x(j-1)^a y(j)^r h(j)^{b-r} & b \geq r \geq 0, \quad m = k+1 - b - r, \\ x(j-1)^a h(j)^r x(j)^{b-r} & b > r \geq 0, \quad m = k+1 - r. \end{cases}$$

The set $\ell(\bar{R}\mathbf{1})$ consists of the elements $\pi_{m\alpha}(n)$, $n \leq -k-1$, $m = -k-1, \dots, k+1$.

Remark. Later on we shall use the map

$$\ell(\bar{R}) \rightarrow \bar{R}, \quad \rho \mapsto r(\rho),$$

defined by $r(\pi_{m\alpha}(n)) = r_{m\alpha}(n)$ for $n \in \mathbb{Z}$, $|m| \leq k+1$. Clearly $\ell(r(\rho)) = \rho$.

In the case of Verma module $M(\Lambda)$ the same argument as above applies for leading terms of $r_{i\alpha}(n)v_\Lambda$, $n \leq -k-1$. We may state this as the following:

Proposition 6.6.2. *For $n \leq -k-1$, $i = -k-1, \dots, k+1$, we have*

$$\ell_{M(\Lambda)}(r_{i\alpha}(n)v_\Lambda) = \ell_{N(k\Lambda_0)}(r_{i\alpha}(n)\mathbf{1}) = \pi_{i\alpha}(n).$$

Hence the ideals $(\ell(\bar{R}v_\Lambda))$ have in common (for different Λ 's) the generators $\pi_{i\alpha}(n)$, $n \leq -k-1$. These generators (colored partitions) have “very small” differences between the largest and the smallest part. Since a partition π from $\mathcal{P}(\bar{B}_-) \setminus (\ell(\bar{R}v_\Lambda))$ does not contain any of these generators, we may say that π satisfies difference conditions defined by $\ell(\bar{R}\mathbf{1})$. In this vein we may also say that the set $\ell(\bar{R}v_\Lambda) \setminus \ell(\bar{R}\mathbf{1})$ represents the initial conditions which π from $\mathcal{P}(\bar{B}_-) \setminus (\ell(\bar{R}v_\Lambda))$ satisfies (since these are the conditions on the “initial part” of colored partition).

Write $\Lambda = k_0\Lambda_0 + k_1\Lambda_1$, $k_0 + k_1 = k$, $k_0, k_1 \in \mathbb{N}$.

Proposition 6.6.3. *The set $\ell(\bar{R}v_\Lambda) \setminus \ell(\bar{R}\mathbf{1})$ consists of elements denoted by $\pi_{m\alpha}^\Lambda(n) = \ell_{M(\Lambda)}(r_{m\alpha}(n)v_\Lambda)$, where $-k-1 < n \leq 0$ and*

$$\pi_{m\alpha}^\Lambda(n) = \begin{cases} y(-1)^r h(-1)^{a-r} y^b & a+b = k+1, a \geq r \geq 0, \\ & m = -r-b, n = -a, \\ h(-1)^r x(-1)^{a-r} y^b & a+b = k+1, a > r \geq 0, \\ & m = -b+a-r, n = -a, \\ x(-1)^a y^r & a+r < k+1, a > k_0 \text{ or } r > k_1, \\ & m = a-r, n = -a. \end{cases}$$

Proposition 6.6.3 will be proved in Section 10, see Proposition 10.2.

Example. *For Λ of level 1 we have the difference conditions:*

$$(6.6.3) \quad \begin{array}{cc} \cdots \square & y \ y \ h \ h \ x \\ \cdots \square & y \ h \ h \ x \ x, \end{array} \quad \begin{array}{cc} \cdots \square \square & y \ h \ x \ x \ x \\ \cdots \square & y \ y \ y \ h \ x. \end{array}$$

For $\Lambda = \Lambda_0$ the initial conditions are $y^2, y, y(-1)y, h(-1)y, x(-1)y$.

For $\Lambda = \Lambda_1$ the initial conditions are $y^2, y(-1)y, h(-1)y, x(-1)y, x(-1)$.

In particular, Theorem 6.5.5 states that for the basic $\mathfrak{g}$ -module $L(\Lambda_0)$ we have a basis consisting of vectors of the form

$$u(\pi)v_{\Lambda_0},$$

where $\pi \in \mathcal{P}(\bar{B}_{<0})$ satisfies the difference condition defined by $\ell(\bar{R}\mathbf{1})$, i.e. π does not contain any of the partitions listed in (6.6.3).

Remark. As we have already mentioned, a notion of the leading term was not introduced in [MP], but still we can consider $X(p; n)$ to be the leading term of the relation $R_p(n)$ in the principal picture. In the case of level 3 modules the corresponding Young diagrams for $X(2; n)$ have only one “color” X and can be identified with

$$\begin{array}{cc} \cdots \square & \cdots \square \square \\ \cdots \square & \cdots \square \end{array} \quad \text{or} \quad \begin{array}{cc} \cdots \square \square & \\ \cdots \square & \end{array}.$$

Partitions which do not contain any of these leading terms are precisely the partitions satisfying Rogers-Ramanujan’s difference 2 conditions.

Note that in Theorem 6.5.5 as well as in [MP, Theorem 8.7] it is easy to show the spanning result for the set of vectors parameterized by partitions which satisfy difference and initial conditions, i.e. partitions which do not contain the leading terms of relations which vanish on a given standard module. It is the linear independence which is difficult to prove, and that involves, one way or the other, the combinatorial identities of Rogers-Ramanujan type.

At this point it may be interesting to note that the combinatorial identities for the $(1, 2)$ -specialization of characters of level one $\mathfrak{sl}(2, \mathbb{C})^\sim$ -modules given by Theorem 6.5.5 are identical with the combinatorial identities obtained by S. Capparelli for level 3 modules in the principal picture for the type $A_2^{(2)}$ (cf. Section 11.3).

7. COLORED PARTITIONS ALLOWING AT LEAST TWO EMBEDDINGS

In Subsection 6.6 we described the set $\ell(\bar{R})$, and, for example, for level 1 modules we have elements $\rho_1 = x(-3)x(-2)$ and $\rho_2 = x(-2)x(-1)$ in $\ell(\bar{R})$. For colored partition $\pi = x(-3)x(-2)x(-1)$ we can construct two vectors $u(\rho_1 \subset \pi)v_\Lambda = x(-1)r(x(-3)x(-2))v_\Lambda$ and $u(\rho_2 \subset \pi)v_\Lambda = x(-3)r(x(-2)x(-1))v_\Lambda$, and by Theorem 6.5.2 it is enough to take just one of them for building a basis of the maximal submodule $M^1(\Lambda)$ of the Verma module $M(\Lambda)$. To prove Theorem 6.5.2 we shall need relations which relate all possible choices $u(\rho_1 \subset \pi)v_\Lambda$ and $u(\rho_2 \subset \pi)v_\Lambda$. In this section we classify all possible choices for $\rho_1 \subset \pi$ and $\rho_2 \subset \pi$.

7.1. Embeddings.

Let $\pi \in \mathcal{P}(\bar{B})$, $\rho \in \ell(\bar{R})$ and let $\rho \subset \pi$. Then we shall say that ρ is *embedded* in π , or that $\rho \subset \pi$ is an *embedding*.

If $\rho \subset \pi$ is an embedding and $\kappa \in \mathcal{P}(\bar{B})$, then $\rho \subset \kappa\pi$ is an embedding as well. For $\rho, \rho' \in \ell(\bar{R})$ we have $\rho, \rho' \subset \rho \cup \rho'$. Since for $\rho \in \ell(\bar{R})$ we have $\ell(\rho) = k + 1$, it is clear from Proposition 6.6.1 that:

Lemma 7.1.1. *Let $\pi = \rho \cup \rho'$. Then either $\rho \cap \rho' = \emptyset$, $\ell(\pi) = 2k + 2$ and $\pi = \rho\rho'$ or $k + 2 \leq \ell(\pi) \leq 2k + 1$ and the shape $\text{sh}(\pi)$ has a form*

$$(j - 1)^a j^b (j + 1)^c$$

for some $j \in \mathbb{Z}$, $b > 0$, $a, c \geq 0$, $a + b + c = \ell(\pi)$.

For $b \in \bar{B}$ define $b^* \in \bar{B}$ by

$$x(j)^* = y(-j), \quad h(j)^* = h(-j), \quad y(j)^* = x(-j).$$

For $\pi = b_1 b_2 \dots b_s \in \mathcal{P}(\bar{B})$ we define a *dual* colored partition $\pi^* = b_s^* b_{s-1}^* \dots b_1^*$.

For $N \in \mathbb{Z}$ and a partition $\pi: \bar{B} \rightarrow \mathbb{Z}$ we may define a *translated* partition $\pi_N = \pi \circ \tau_N$, where

$$\tau_N(x(i)) = x(i - N), \quad \tau_N(h(i)) = h(i - N), \quad \tau_N(y(i)) = y(i - N).$$

It follows from Proposition 6.6.1 that

Proposition 7.1.2. *The set $\ell(\bar{R})$ is translation invariant and self-dual (i.e. for $\rho \in \ell(\bar{R})$ we have $\rho^* \in \ell(\bar{R})$).*

We shall sometimes also say that π and π' are dual to each other if π' is dual to some translation of π . In this way we shall also speak of dual embeddings.

7.2. Colored partitions of length $k + 2$ allowing at least two embeddings.

From Proposition 6.6.1 we get the following list of partitions of length $k + 2$ allowing two or more than two embeddings (here N denotes the number of embeddings):

$$(1) \quad \text{sh } \pi = j^{k+2},$$

$$h(j)^a x(j)^{k+2-a}, \quad 1 \leq a \leq k + 1, \quad N = 2,$$

$$y(j)h(j)^k x(j), \quad N = 2,$$

$$y(j)^{k+2-a} h(j)^a, \quad 1 \leq a \leq k + 1, \quad N = 2,$$

$$(2) \quad \text{sh } \pi = (j-1)^a j^b, \quad a, b > 0, \quad a + b = k + 2,$$

$$\begin{array}{ll}
x(j-1)^a x(j)^b, & N = 2, \\
x(j-1)^a h(j)^c x(j)^{b-c}, \quad 1 \leq c \leq b-1, & N = 3, \\
x(j-1)^a h(j)^b, & N = 2, \\
(*) \quad x(j-1)^a y(j) h(j)^{b-2} x(j), & N = 2, \\
x(j-1)^a y(j)^c h(j)^{b-c}, \quad 1 \leq c \leq b-1, & N = 3, \\
x(j-1)^a y(j)^b, & N = 2, \\
h(j-1) x(j-1)^{a-1} y(j)^{b-1} h(j), & N = 2, \\
h(j-1)^c x(j-1)^{a-c} y(j)^b, \quad 1 \leq c \leq a-1, & N = 3, \\
h(j-1)^a y(j)^b, & N = 2, \\
(*) \quad y(j-1) h(j-1)^{a-2} x(j-1) y(j)^b, & N = 2, \\
y(j-1)^c h(j-1)^{a-c} y(j)^b, \quad 1 \leq c \leq a-1, & N = 3, \\
y(j-1)^a y(j)^b, & N = 2.
\end{array}$$

Note that for $a \geq 2$ and $b \geq 2$ elements denoted by (*) are well defined. Also note that elements are listed so that their weights descend from $(k+2)\alpha$ to $-(k+2)\alpha$, where for each element denoted by (*) there is another element of the same weight. In the case $b = 1$ (and $a = k + 1$) the list should be modified by elements

$$\begin{cases} x(j-1)^a h(j), & N = 2, \\ h(j-1) x(j-1)^{a-1} x(j), & N = 2, \end{cases}$$

and in the case $a = 1$ (and $b = k + 1$) by

$$\begin{cases} h(j-1) y(j)^b, & N = 2, \\ y(j-1) y(j)^{b-1} h(j), & N = 2. \end{cases}$$

$$(3) \quad \text{sh } \pi = (j-1) j^k (j+1),$$

$$\begin{array}{ll}
x(j-1) x(j)^k x(j+1)^a h(j+1)^b, \quad a, b \geq 0, a + b = 1, & N = 2, \\
x(j-1) h(j)^a x(j)^{k-a} y(j+1), \quad 0 \leq a \leq k, & N = 2, \\
x(j-1) y(j)^a h(j)^{k-a} y(j+1), \quad 1 \leq a \leq k, & N = 2, \\
y(j-1)^a h(j-1)^b y(j)^k y(j+1), \quad a, b \geq 0, a + b = 1, & N = 2.
\end{array}$$

Note that the above list of embeddings is self dual.

7.3. Linked embeddings and exceptional cases.

Let us call two embeddings $\rho, \rho' \subset \pi$ *linked* if there is a sequence

$$\rho_1 = \rho \subset \pi, \rho_2 \subset \pi, \dots, \rho_{n-1} \subset \pi, \rho_n = \rho' \subset \pi$$

of embeddings such that $\ell(\rho_{i-1} \cup \rho_i) < \ell(\rho \cup \rho')$ for each $i = 2, \dots, n$.

The above definition is designed for inductive arguments. However, there are *exceptional cases* that we will have to consider separately:

Lemma 7.3.1. *Let π be a colored partition, $\rho_1, \rho_2 \subset \pi$ leading terms of relations such that $\pi = \rho_1 \cup \rho_2$, $\rho_1 \cap \rho_2 \neq \emptyset$, $\ell(\pi) \geq k + 3$, and the two embeddings $\rho_1 \subset \pi$, $\rho_2 \subset \pi$ are not linked. Then π is one of the following colored partitions:*

- (1) $y(j)^a h(j)^{k+1-a} x(j)^a$,
 $2 \leq a \leq k$,
- (2) $x(j-1)^{k+1-a-b} y(j)^a h(j)^b x(j)^a$,
 $a \geq 2, a+b \leq k$,
- (3) $h(j-1)^{k+1-a-b} x(j-1)^a y(j)^b x(j)^{k+1-a}$,
 $1 \leq a \leq k-1$,
- (4) $h(j-1)^{k+1-a-b} x(j-1)^a y(j)^b h(j)^{k+1-a-b}$,
 $1 \leq a+b \leq k-1$,
- (5) $y(j-1)^{k+1-a} x(j-1)^b y(j)^a h(j)^{k+1-a-b}$,
 $1 \leq a \leq k-1$,
- (6) $y(j-1)^a h(j-1)^b x(j-1)^a y(j)^{k+1-a-b}$,
 $a \geq 2, a+b \leq k$,
- (7) $y(j-1)^{k+1-a-c} h(j-1)^c x(j-1)^b y(j)^a h(j)^{k+1-a-b}$,
 $a, c \geq 1, a+b+c \leq k$,
- (8) $h(j-1)^{k+1-a-b} x(j-1)^a y(j)^b h(j)^c x(j)^{k+1-a-c}$,
 $a, c \geq 1, a+b+c \leq k$,
- (9) $x(j-2)^a y(j-1)^b h(j-1)^{k+1-a-b} y(j)^a$,
 $2 \leq a \leq k$,
- (10) $x(j-2)^a h(j-1)^{k+1-a-b} x(j-1)^b y(j)^a$,
 $2 \leq a \leq k$,
- (11) $x(j-2)^{k+1-a-b} h(j-1)^a x(j-1)^b y(j)^c h(j)^{k+1-b-c}$,
 $1 \leq b \leq k-1, a+b+c \leq k$,
- (12) $x(j-2)^{k+1-a-b} h(j-1)^a x(j-1)^b h(j)^c x(j)^{k+1-b-c}$,
 $1 \leq b \leq k-1, a+b \leq k$,
- (13) $h(j-2)^{k+1-b-c} x(j-2)^c y(j-1)^b h(j-1)^a y(j)^{k+1-a-b}$,
 $1 \leq b \leq k-1, a+b+c \leq k$,
- (14) $y(j-2)^{k+1-b-c} h(j-2)^c y(j-1)^b h(j-1)^a y(j)^{k+1-a-b}$,
 $1 \leq b \leq k-1, a+b \leq k$.

Here (and later on) it goes without saying that X^a , $X \in \bar{B}$, is defined only for integers $a \geq 0$ (in particular $a = 0$ is allowed unless explicitly ruled out).

Note that the following cases are dual to each other:

$$(1)^* = (1), (2)^* = (6), (3)^* = (5), (4)^* = (4),$$

$$(7)^* = (8), (9)^* = (10), (11)^* = (13), (12)^* = (14).$$

We could change case (7) to include $c=0$ covering (5), and we could change case (8) to include $c=0$ covering (3), but for later proofs it will be convenient to keep them separated. Note also that in each of the cases (1)–(14) there are only two embeddings $\rho_1 \subset \pi, \rho_2 \subset \pi$ of leading terms of relation, so that ρ_1, ρ_2 are determined up to permutation.

Proof. We shall often use the following “replacement of representative”:

$$(A) \left\{ \begin{array}{l} \text{Let } \pi = \pi' y(j)^d, \rho_i = \rho' y(j)^c \text{ (} i = 1 \text{ or } 2 \text{), where } \text{sh}(\rho_i) = (j-1)^{k+1-c} j^c \\ \text{and } y(j)^d, y(j)^c \text{ are the maximal powers of } y(j) \text{ contained in } \pi \text{ and } \rho_i \\ \text{(respectively). Then we may, by replacement of } \rho_i \text{ by } \rho_3 = \rho'' y(j)^d \text{ with} \\ \rho'' \subset \rho' \text{ if necessary, assume that } c = d, \text{ that is } \rho_i \supset y(j)^d. \\ \text{Dually, if } \pi \supset x(j-1)^d, \rho_i \supset x(j-1)^c, \text{sh}(\rho_i) = (j-1)^c j^{k+1-c}, \\ \text{then we may assume that } \rho_i \supset x(j-1)^d. \end{array} \right.$$

Assume first that $\text{sh}(\pi) = j^\ell$, $k+3 \leq \ell \leq 2k+1$. Then $\pi = y(j)^a h(j)^b x(j)^c$. Clearly ρ_1, ρ_2 are linked if a or $c = 0$, so $a, c \geq 1$. We may then assume $\rho_1 \subset y(j)^a h(j)^b$, $\rho_2 \subset h(j)^b x(j)^c$. If $a+b$ or $b+c > k+1$, ρ_1 and ρ_2 are linked. Thus π must be of type (1).

Consider now the case that $\text{sh}(\pi) = (j-1)^\ell j^m$ for some $\ell, m \geq 1$. Then

$$\pi = y(j-1)^a h(j-1)^b x(j-1)^c y(j)^d h(j)^e x(j)^f$$

for some $a, \dots, f$. It cannot be that $\text{sh}(\rho_1) = (j-1)^{k+1}$ and $\text{sh}(\rho_2) = j^{k+1}$ since $\rho_1 \cap \rho_2 \neq \emptyset$. We may thus assume that $\text{sh}(\rho_1) = (j-1)^p j^{k+1-p}$ for some $1 \leq p \leq k$. By duality and (A) we may assume that $\rho_1 \supset y(j)^d$. By (A) we may also assume that ρ_2 satisfies one of the following:

$$\begin{aligned} (\alpha) \quad & \rho_2 \supset y(j)^d, \quad \text{sh}(\rho_2) = (j-1)^{k+1-d} j^d, \\ (\beta) \quad & \rho_2 \supset x(j-1)^c, \quad c \geq 1, \quad \text{sh}(\rho_2) = (j-1)^c j^{k+1-c}, \\ (\gamma) \quad & \rho_2 \subset y(j)^d h(j)^e, \\ (\delta) \quad & \rho_2 \subset y(j-1)^a h(j-1)^b x(j-1)^c. \end{aligned}$$

Case (α): If $\rho_1 = y(j-1)^a h(j-1)^{b'} y(j)^d$, $\rho_2 = h(j-1)^{b''} x(j-1)^c y(j)^d$, then ρ_1 and ρ_2 are linked unless $b' = b = b''$ in which case π is of type (6). If $a = 0$ or $c = 0$, ρ_1 and ρ_2 are linked.

Case (β): If $\rho_1 \subset y(j-1)^a h(j-1)^b y(j)^d$ and $\rho_2 \subset x(j-1)^c h(j)^e x(j)^f$ then $\rho_1 \cap \rho_2 = \emptyset$. By duality we may thus assume that $\rho_1 = h(j-1)^b x(j-1)^{c'} y(j)^d$ and $a = 0$. Then ρ_1 and ρ_2 are linked if $c' < c$ so $c' = c$. If $\rho_2 = x(j-1)^c h(j)^e x(j)^f$ we have π of type (8) or (3). In case π of type (8) we must have $a+b+c \leq k$, otherwise ρ_1 and ρ_2 are linked via $\rho_3 = x(j-1)^a y(j)^b h(j)^{k+1-a-b}$. If $\rho_2 = x(j-1)^c y(j)^{d'} h(j)^e$ then ρ_1 and ρ_2 are linked if $d' < d$, so $d' = d$ and π is of type (4).

Case (γ): If $c \geq 1$, then ρ_1 and ρ_2 are linked via $\rho_3 = x(j-1) y(j)^d h(j)^{k-d}$, so $c = 0$. If $\rho_2 = y(j)^{d'} h(j)^e$ with $d' < d$, then ρ_1 and ρ_2 are linked via $\rho_3 = y(j)^d h(j)^{k+1-d}$. Thus $d' = d$ and π is of type (7).

Case (δ): If $\rho_2 = y(j-1)^{k+1-b'} h(j-1)^{b'}$ and $\rho_1 = h(j-1)^{b''} x(j-1)^c y(j)^d$ then it is easy to see that ρ_1 and ρ_2 are linked unless $b' = b = b''$. In that case ρ_1 and ρ_2

are linked via $y(j-1)^{k-b}h(j-1)^by(j)$. If $c = 0$ and $\rho_1 = y(j-1)^{a'}h(j-1)^{b'}y(j)^d$, then ρ_1 and ρ_2 are linked via either $\rho_3 = y(j-1)^{a'+1}h(j-1)^{b'}y(j)^{d-1}$ or $\rho_3 = y(j-1)^{a'}h(j-1)^{b'+1}y(j)^{d-1}$. Similarly ρ_1 and ρ_2 are linked in all cases where $\rho_2 \subset h(j-1)^bx(j-1)^c$.

There remains the case that $\text{sh}(\pi) = (j-2)^p(j-1)^qj^r$ for some $p, q, r \geq 1$. We may then assume $\text{sh}(\rho_1) = (j-2)^p(j-1)^{k+1-p}$, $\text{sh}(\rho_2) = (j-1)^{k+1-r}j^r$. Since $\rho_1 \cap \rho_2 \neq \emptyset$ it cannot be that $\rho_1 \supset y(j-1)^{k+1-p}$, $\rho_2 \supset x(j-1)^{k+1-r}$. By duality we may thus assume that $\rho_1 \supset x(j-2)^p$. Then either

$$\begin{aligned} (\varepsilon) \quad & \rho_2 \supset x(j-1)^{k+1-r}, \text{ or} \\ (\zeta) \quad & \rho_2 \supset y(j)^r. \end{aligned}$$

Case (ε) : By (A) $x(j-1)^{k+1-r}$ is the maximal power of $x(j-1)$ contained in π . We must have $\rho_1 = x(j-2)^ph(j-1)^{k+1-p-a}x(j-1)^a$ for some a . If $a < k+1-r$, then ρ_1 and ρ_2 are linked via $x(j-2)^{p-1}h(j-1)^{k+1-p-a}x(j-1)^{a+1}$. Thus $a = k+1-r$ and π is of type (10), (11) or (12). If π is of type (11) and $a \geq 1$, then $a+b+c \leq k$ since otherwise ρ_1 and ρ_2 are linked via $\rho_3 = h(j-1)^{a'}x(j-1)^by(j)^{k+1-a'-b}$ for suitable $a' \geq 1$.

Case (ζ) : If $\rho_1 = x(j-2)^py(j-1)^ah(j-1)^{k+1-a-p}$ and $\rho_2 = h(j-1)^{k+1-b-r}x(j-1)^by(j)^r$ and $1 \leq a \leq b$, then ρ_1 and ρ_2 are linked via $\rho_3 = x(j-2)^ph(j-1)^{k+1-a-p}x(j-1)^a$. If instead $1 \leq b \leq a$, then ρ_1 and ρ_2 are linked via $\rho_3 = y(j-2)^bh(j-1)^{k+1-b-r}y(j)^r$. Similarly ρ_1 and ρ_2 are linked if $\rho_1 = x(j-2)^ph(j-1)^{k+1-a-p}x(j-1)^a$, $\rho_2 = y(j-1)^bh(j-1)^{k+1-r-b}y(j)^r$. We must therefore have $\pi = x(j-2)^py(j-1)^ah(j-1)^by(j)^r$ or $\pi = x(j-2)^ph(j-1)^ax(j-1)^by(j)^r$, and now it is easy to see that ρ_1 and ρ_2 are linked except in case π is of type (9) or (10). $\square$

8. RELATIONS AMONG RELATIONS

By Theorem 5.14 we have $M^1(\Lambda) = U_k(\tilde{\mathfrak{g}})\bar{R}v_\Lambda$, so $M^1(\Lambda)$ is spanned by elements of the form $u(\rho \subset \pi)v_\Lambda = u(\pi/\rho)r(\rho)v_\Lambda$, where ρ 's are leading terms of "relations" $r(\rho)$ in $\bar{R}$ and π 's are in the ideal $(\ell(\bar{R}))$. We will prove that for two elements of the form $u(\rho \subset \pi)$ and $u(\rho' \subset \pi)$ there are relations that hold on every highest weight module of level k , and we shall refer to them as *relations among relations*.

Proposition 8.1. *We have*

$$(8.1) \quad \sum_{j \in \mathbb{Z}} ((k+2)j - n) x(j) r_{(k+1)\alpha}(n-j) = 0$$

in $\overline{U_k(\tilde{\mathfrak{g}})}$ for all $n \in \mathbb{Z}$.

Note that $x(j)$ and $r_{(k+1)\alpha}(n-j)$ commute so that the infinite sum is convergent.

Proof. It follows from $[L_{-1}, x(-1)] = x(-2)$ and $L_{-1}\mathbf{1} = 0$ that

$$(8.2) \quad L_{-1}x(-1)^{k+2}\mathbf{1} = (k+2)x(-2)x(-1)^{k+1}\mathbf{1}.$$

By (2.4), (2.13) and (2.14)

$$\begin{aligned} Y(L_{-1}x(-1)^{k+2}\mathbf{1}, z) &= \frac{d}{dz} \left(x(z)Y(r_{(k+1)\alpha}, z) \right), \\ Y(x(-2)x(-1)^{k+1}\mathbf{1}, z) &= \left(\frac{d}{dz} x(z) \right) Y(r_{(k+1)\alpha}, z). \end{aligned}$$

Thus (8.2) implies

$$(8.3) \quad \frac{d}{dz} \left(x(z)Y(r_{(k+1)\alpha}, z) \right) = (k+2) \left(\frac{d}{dz} x(z) \right) Y(r_{(k+1)\alpha}, z).$$

The coefficient of z^{-n-k-3} in (8.3) gives (8.1). $\square$

Proposition 8.2. *We have*

$$\begin{aligned} (8.4) \quad & \sum_{j<0} ((k+2)j-n) \{x(j)r_{(i-1)\alpha}(n-j) - h(j)r_{i\alpha}(n-j) - y(j)r_{(i+1)\alpha}(n-j)\} \\ & + \sum_{j\geq 0} ((k+2)j-n) \{r_{(i-1)\alpha}(n-j)x(j) - r_{i\alpha}(n-j)h(j) - r_{(i+1)\alpha}(n-j)y(j)\} = 0 \end{aligned}$$

in $\overline{U_k(\tilde{\mathfrak{g}})}$ for all $n \in \mathbb{Z}$, $i = -k-2, \dots, k+2$.

Proof. The case $i = k+2$ is (8.1). If we denote the left hand side in (8.4) by s_i then one finds

$$[y(0), s_i] = (k+3-i)s_{i-1}$$

and (8.4) follows by induction. $\square$

Proposition 8.3. *We have*

$$\begin{aligned} (8.5) \quad & (k+2-i) \left(\sum_{j<0} x(j)r_{(i-1)\alpha}(n-j) + \sum_{j\geq 0} r_{(i-1)\alpha}(n-j)x(j) \right) \\ & + i \left(\sum_{j<0} h(j)r_{i\alpha}(n-j) + \sum_{j\geq 0} r_{i\alpha}(n-j)h(j) \right) \\ & + (k+2+i) \left(\sum_{j<0} y(j)r_{(i+1)\alpha}(n-j) + \sum_{j\geq 0} r_{(i+1)\alpha}(n-j)y(j) \right) \\ & = (k+2)(-n-k-1)r_{i\alpha}(n) \end{aligned}$$

in $\overline{U_k(\tilde{\mathfrak{g}})}$ for all $n \in \mathbb{Z}$, $i = -k-1, \dots, k+1$.

Proof. By (3.17), (3.18) and (6.6.1)

$$\begin{aligned} L_{-1}r_{i\alpha} &= \frac{1}{2(k+2)} (2x(-1)y(0) + h(-1)h(0) + 2y(-1)x(0))r_{i\alpha} \\ &= \frac{1}{k+2} ((k+2-i)x(-1)r_{(i-1)\alpha} + ih(-1)r_{i\alpha} + (k+2+i)y(-1)r_{(i+1)\alpha}). \end{aligned}$$

From (2.4) and (2.14) it follows that

$$\begin{aligned}
& (k+2-i)(x(z)^-Y(r_{(i-1)\alpha}, z) + Y(r_{(i-1)\alpha}, z)x(z)^+) \\
& \quad + i(h(z)^-Y(r_{i\alpha}, z) + Y(r_{i\alpha}, z)h(z)^+) \\
& + (k+2+i)(y(z)^-Y(r_{(i+1)\alpha}, z) + Y(r_{(i+1)\alpha}, z)y(z)^+) \\
& = (k+2)\frac{d}{dz}Y(r_{i\alpha}, z).
\end{aligned}$$

The coefficient of z^{-n-k-2} gives (8.5). $\square$

Let $j \in \mathbb{Z}$, $j < 0$, $0 \leq p \leq k+2$ and consider the shape

$$\sigma = (j-1)^p j^{k+2-p}.$$

Set $n = |\sigma| = (k+2)j - p$. By taking linear combinations of (8.4) and (8.5) we can produce relations in $\overline{U_k(\tilde{\mathfrak{g}})}$ of the form

$$\begin{aligned}
(8.6) \quad & C^x x(j-1)r_{(i-1)\alpha}(n-j+1) + C^h h(j-1)r_{i\alpha}(n-j+1) + C^y y(j-1)r_{(i+1)\alpha}(n-j+1) \\
& + C_x x(j)r_{(i-1)\alpha}(n-j) + C_h h(j)r_{i\alpha}(n-j) + C_y y(j)r_{(i+1)\alpha}(n-j) \\
& + \text{terms of higher shape} = 0.
\end{aligned}$$

Calculations of some 2×2 determinants show the following:

Lemma 8.4. *There is a relation (8.6) with*

$$\begin{aligned}
\{C^x, C_x\} &= \{0, 1\} \quad \text{unless } i = k+2, \\
\{C^y, C_y\} &= \{0, 1\} \quad \text{unless } i = -k-2, \\
\{C^x, C^h\} &= \{0, 1\} \quad \text{unless } \sigma = (j-1)^{k+2}, \\
\{C^h, C^y\} &= \{0, 1\} \quad \text{unless } \sigma = (j-1)^{k+2}, \\
\{C_x, C_h\} &= \{0, 1\} \quad \text{unless } \sigma = j^{k+2}, \\
\{C_h, C_y\} &= \{0, 1\} \quad \text{unless } \sigma = j^{k+2}.
\end{aligned}$$

Remark. The identities (8.4) and (8.5) are analogous to [MP, Proposition 9.3] in the principal picture.

Remarks. Since we have exact sequences

$$0 \rightarrow M^1(\Lambda) \rightarrow M(\Lambda) \rightarrow L(\Lambda) \rightarrow 0,$$

$$0 \rightarrow N^1(k\Lambda_0) \rightarrow N(k\Lambda_0) \rightarrow L(k\Lambda_0) \rightarrow 0,$$

we may call elements of $M^1(\Lambda) = U(\tilde{\mathfrak{g}})\bar{R}v_\Lambda$ and/or of $N^1(k\Lambda_0) = U(\tilde{\mathfrak{g}})\bar{R}\mathbf{1}$ the relations (which define the standard module $L(\Lambda)$ and/or $L(k\Lambda_0)$).

Obviously $\bar{R}v_\Lambda \subset M(\Lambda)$ is a $(\mathfrak{h}^e + \tilde{\mathfrak{n}}_+)$ -module, and $\bar{R}\mathbf{1} \subset N(\Lambda)$ is a $\tilde{\mathfrak{g}}_{\geq 0}$ -module, so we have exact sequences

$$0 \rightarrow \ker \Psi_{M(\Lambda)} \rightarrow U(\tilde{\mathfrak{g}}) \otimes_{U(\mathfrak{h}^e + \tilde{\mathfrak{n}}_+)} \bar{R}v_\Lambda \xrightarrow{\Psi_{M(\Lambda)}} M^1(\Lambda) \rightarrow 0,$$

$$0 \rightarrow \ker \Psi_{N(k\Lambda_0)} \rightarrow U(\tilde{\mathfrak{g}}) \otimes_{U(\tilde{\mathfrak{g}}_{\geq 0})} \bar{R}\mathbf{1} \xrightarrow{\Psi_{N(k\Lambda_0)}} N^1(k\Lambda_0) \rightarrow 0,$$

where $\Psi_{M(\Lambda)}(u \otimes rv_\Lambda) = urv_\Lambda$, $\Psi_{N(k\Lambda_0)}(u \otimes r\mathbf{1}) = ur\mathbf{1}$. Since $\ker \Psi$ defines the quotient (“which are relations”), we may call the elements of $\ker \Psi$ relations among relations.

One can see that the $U_k(\tilde{\mathfrak{g}})$ -module $\ker \Psi_{N(k\Lambda_0)}$ is generated by the elements $q^1(n)$, $n \leq -k-3$, and $q^2(n)$, $n \leq -k-2$, where

$$\begin{aligned} q^1(n) &= \sum_{j \leq -1} ((k+2)j - n)x(j) \otimes r_{(k+1)\alpha}(n-j)\mathbf{1}, \\ q^2(n) &= - \sum_{i \leq -1} (k+1)h(i) \otimes r_{(k+1)\alpha}(n-i)\mathbf{1} \\ &\quad - \sum_{i \leq -1} x(i) \otimes r_{k\alpha}(n-i)\mathbf{1} \\ &\quad - (k+2)(n+k+1)\mathbf{1} \otimes r_{(k+1)\alpha}(n)\mathbf{1}. \end{aligned}$$

So in some sense $q^1(n)$ and $q^2(n)$ (or the corresponding (8.4) and (8.5)) form a “complete generating set” of relations among relations.

For an element of the form $u \otimes r\mathbf{1}$ in $U(\tilde{\mathfrak{g}}_{<0}) \otimes \bar{R}\mathbf{1}$ define a “leading term” $\ell t(u \otimes r\mathbf{1}) = (\ell t(u), \ell t(r\mathbf{1}))$. We may identify a “leading term” (κ, ρ) with the embedding $\rho \subset \kappa\rho$. So each nonzero element q in $U(\tilde{\mathfrak{g}}) \otimes_{U(\tilde{\mathfrak{g}}_{\geq 0})} \bar{R}\mathbf{1} \cong U(\tilde{\mathfrak{g}}_{<0}) \otimes \bar{R}\mathbf{1}$ can be written in the form

$$q = \sum_{\rho' \subset \pi} C_{\rho', \pi} u(\pi/\rho') \otimes r(\rho')\mathbf{1} + \sum_{\substack{\rho' \subset \pi' \\ \pi' \succ \pi}} C_{\rho', \pi'} u(\pi'/\rho') \otimes r(\rho')\mathbf{1}$$

for some $\pi \in \mathcal{P}(\bar{B}_{<0})$ and $C_{\rho', \pi}, C_{\rho', \pi'} \in \mathbb{F}$, where at least one coefficient $C_{\rho', \pi}$ is nonzero. Then $q \in \ker \Psi_{N(k\Lambda_0)}$, i.e.

$$\sum_{\rho' \subset \pi} C_{\rho', \pi} u(\rho' \subset \pi)\mathbf{1} + \sum_{\substack{\rho' \subset \pi' \\ \pi' \succ \pi}} C_{\rho', \pi'} u(\rho' \subset \pi')\mathbf{1} = 0,$$

implies $\sum_{\rho' \subset \pi} C_{\rho', \pi} = 0$. Hence for such q there must be at least two embeddings $\rho' \subset \pi$, $\rho'' \subset \pi$ with the corresponding coefficients being nonzero.

9. RELATIONS AMONG RELATIONS FOR TWO EMBEDDINGS

We fix a map (analogous to the one in Section 6.5)

$$\ell t(\bar{R}) \rightarrow \bar{R} \quad , \quad \rho \mapsto r(\rho),$$

such that $\ell t(r(\rho)) = \rho$. For $\rho \in \ell t(\bar{R})$ and $\pi \in (\ell t(\bar{R}))$, $\rho \subset \pi$, we define

$$u(\rho \subset \pi) = u(\pi/\rho)r(\rho).$$

For $\pi \in \mathcal{P}(\bar{B})$ set

$$\mathcal{R}_{(\pi)} = \overline{\mathbb{F}\text{-span}\{u(\rho \subset \pi') \mid \rho \in \ell t(\bar{R}), \pi' \in (\ell t(\bar{R})), \rho \subset \pi', \pi \prec \pi'\}},$$

the closure taken in $\overline{U_k(\tilde{\mathfrak{g}})}$.

The main result in this section is the following consequence of relations among relations constructed in Section 8:

Theorem 9.1. *Let $\pi \in (\ell(\bar{R}))$, $\rho_1, \rho_2 \in \ell(\bar{R})$, $\rho_1 \subset \pi$, $\rho_2 \subset \pi$. Then*

$$(9.1) \quad u(\rho_1 \subset \pi) \in \mathbb{F}^\times u(\rho_2 \subset \pi) + \mathcal{R}_{(\pi)}.$$

Remark. Since $N^1(k\Lambda_0) = U(\tilde{\mathfrak{g}}_{<0})\bar{R}\mathbf{1}$, the set of vectors $u(\rho \subset \pi)\mathbf{1}$, where $\rho \in \ell(\bar{R}\mathbf{1})$, $\pi \in (\ell(\bar{R}\mathbf{1}))$, $\rho \subset \pi$, is a spanning set of $N^1(k\Lambda_0)$. Let $(\ell(\bar{R}\mathbf{1})) \ni \pi \mapsto \rho(\pi) \in \ell(\bar{R}\mathbf{1})$ be such that $\rho(\pi) \subset \pi$. Using Theorem 9.1 and induction we see that

$$u(\rho(\pi) \subset \pi)\mathbf{1}, \quad \pi \in (\ell(\bar{R}\mathbf{1})),$$

is a spanning set as well. Clearly this set is linearly independent, i.e. it is a basis of $N^1(k\Lambda_0)$. As a consequence we have

$$\ell_{N(k\Lambda_0)}(N^1(k\Lambda_0)) = (\ell(\bar{R}\mathbf{1})),$$

which in turn implies (see the proof of Theorem 6.5.5) that the set of vectors

$$u(\pi)v_{k\Lambda_0}, \quad \pi \in \mathcal{P}(\bar{B}_{<0}) \setminus (\ell(\bar{R}\mathbf{1})),$$

is a basis of the standard $\tilde{\mathfrak{g}}$ -module $L(k\Lambda_0) = N(k\Lambda_0)/N^1(k\Lambda_0)$.

The rest of this section is devoted to the proof of Theorem 9.1.

We may and we will assume that the map $\rho \mapsto r(\rho)$ is such that $r(\rho)$ is homogeneous of weight $\text{wt}(\rho)$ and degree $|\rho|$. For example, we may take the map $\pi_{m\alpha}(n) \mapsto r_{m\alpha}(n)$, $n \in \mathbb{Z}$, $|m| \leq k+1$ (cf. Section 6.6).

Lemma 9.2. *Let $\rho_1, \rho_2 \in \ell(\bar{R})$, $\pi = \rho_1 \cup \rho_2$, $\ell(\pi) = k+2$. Then*

$$u(\rho_1 \subset \pi) \in \mathbb{F}^\times u(\rho_2 \subset \pi) + \mathcal{R}_{(\pi)}.$$

Proof. In Section 7.2 we have listed all π of length $k+2$ which allow more than one embedding. For a fixed degree and weight only one of three distinct cases may occur: among all colored partitions of length $k+2$ of fixed degree and weight a) there is one π with two embeddings, b) there are two π_1 and π_2 with two embeddings each, and c) there is one π with three embeddings. Note that we have a) for $\text{wt}(\pi) = \pm(k+2)\alpha$, and b) and c) for $\text{wt}(\pi) = j\alpha$, $-k-1 \leq j \leq k+1$. Also note that cases (1) and (3) in Section 7.2 go together for $|\pi| \equiv 0 \pmod{k+2}$.

In Section 8 we have constructed relations among relations (which hold on every highest weight module of level k) of the form

$$(9.2) \quad \sum c_i u(\rho_i \subset \pi_i) = 0,$$

where (for the sake of notation we assume that) $c_i \neq 0$ and $\pi_i = \pi_j$ implies $\rho_i \neq \rho_j$. Note that relation (8.4) holds for weights $j\alpha$, $-k-2 \leq j \leq k+2$, and that (8.5) holds for weights $j\alpha$, $-k-1 \leq j \leq k+1$. So in the case a) we have one linearly independent relation, and in the case b) or c) there are two linearly independent relations.

Fix a degree and weight and write a relation in the form (9.2). Let π be the smallest among π_i 's which appear in the sum, say $\pi = \pi_1$. Then

$$c_1 u(\rho_1 \subset \pi) + \sum_{\substack{i \neq 1 \\ \pi \preceq \pi_i}} c_i u(\rho_i \subset \pi_i) = 0.$$

In the case that $\pi \prec \pi_i$ for all other π_i we would have that the expression on the left has a leading term π , and then the expression on the left would be different from 0. Hence there are $\rho_2 \subset \pi_2, \dots, \rho_s \subset \pi_s$ such that $\pi_2 = \dots = \pi_s = \pi$ and we have

$$(9.3) \quad c_1 u(\rho_1 \subset \pi) + \dots + c_s u(\rho_s \subset \pi) + \sum_{\pi \prec \pi_i} c_i u(\rho_i \subset \pi_i) = 0.$$

So in particular there exists π with at least two embeddings $\rho_1 \subset \pi, \rho_2 \subset \pi, \dots$

Assume that we have the case a). Then $\rho_1 \subset \pi$ and $\rho_2 \subset \pi$ are uniquely determined and (9.3) is the statement of our lemma.

Assume that we have the case b). Then we have two linearly independent relations: R of the form (9.2) and R' of the form

$$(9.4) \quad \sum c'_i u(\rho'_i \subset \pi'_i) = 0,$$

where $c'_i \neq 0$, and $\pi'_i = \pi'_j$ implies $\rho'_i \neq \rho'_j$. We also have two colored partitions, say π_1 and π_2 with two embeddings each.

Each relation (9.2) and (9.4) can be written in the form (9.3) (with $s = 2$), in each case $\pi', \pi \in \{\pi_1, \pi_2\}$. If $\pi' \neq \pi$, then we have the lemma. If for both relations (say) $\pi = \pi' = \pi_1$, then the terms $c_1 u(\rho_1 \subset \pi_1) + c_2 u(\rho_2 \subset \pi_1)$ and $c'_1 u(\rho'_1 \subset \pi_1) + c'_2 u(\rho'_2 \subset \pi_1)$ are proportional by a factor $\lambda \neq 0$. Then the (nontrivial) relation $R - \lambda R'$ written in the form (9.3) should give the only possible $\pi = \pi_2$, and the lemma holds.

Assume that we have the case c). Then two linearly independent relations (9.2) and (9.4) can be written in the form (9.3)

$$\begin{aligned} c_1 u(\rho_1 \subset \pi) + c_2 u(\rho_2 \subset \pi) + c_3 u(\rho_3 \subset \pi) + \sum_{\pi \prec \pi_i} c_i u(\rho_i \subset \pi_i) &= 0, \\ c'_1 u(\rho_1 \subset \pi) + c'_2 u(\rho_2 \subset \pi) + c'_3 u(\rho_3 \subset \pi) + \sum_{\pi \prec \pi'_i} c'_i u(\rho'_i \subset \pi'_i) &= 0. \end{aligned}$$

The matrix $\begin{pmatrix} c_1 & c_2 & c_3 \\ c'_1 & c'_2 & c'_3 \end{pmatrix}$ must have rank 2 (since otherwise the nontrivial relation of the form $R'' = R - \lambda R'$ would give another $\pi'' \neq \pi$ allowing more than one embedding). Hence we can find two linear combinations R_1 and R'_1 of R and R' such that the corresponding matrix would be (after a suitable permutation of columns) of the form

$$\begin{pmatrix} 1 & 0 & c \\ 0 & 1 & c' \end{pmatrix}.$$

Hence R_1 and R'_1 give relations stated in lemma. $\square$

Lemma 9.3. *Let $\rho_1, \rho_2 \in \ell t(\bar{R})$, $\pi = \rho_1 \rho_2$. Then*

$$u(\rho_1 \subset \pi) \in \mathbb{F}^\times u(\rho_2 \subset \pi) + \mathcal{R}_{(\pi)}.$$

Proof. The element $r(\rho_1)r(\rho_2)$ can be “expanded” in two ways: by writing either $r(\rho_1)$ or $r(\rho_2)$ as an (infinite) sum of monomials. Hence

$$r(\rho_1)r(\rho_2) = c_1 u(\rho_1)r(\rho_2) + \sum_{\rho_1 \prec \kappa} c_\kappa^1 u(\kappa)r(\rho_2),$$

$$r(\rho_1)r(\rho_2) = c_2 r(\rho_1)u(\rho_2) + \sum_{\rho_2 \prec \kappa} c_\kappa^2 r(\rho_1)u(\kappa),$$

$c_1, c_2 \in \mathbb{F}^\times$, $c_\kappa^1, c_\kappa^2 \in \mathbb{F}$, and the lemma follows. $\square$

Remark. In Section 8 we have constructed some relations among relations, but for the proof of Theorem 6.5.2 we need the relations (9.1) stated in terms of colored partitions. The first step is accomplished by Lemma 9.2 for “adjacent” embeddings like the one we had in the case of $\pi = x(-3)x(-2)x(-1)$, $\rho_1 = x(-3)x(-2)$ and $\rho_2 = x(-2)x(-1)$. On the other hand, Lemma 9.3 settles the problem of “disjoint” embeddings like for $\pi = x(-5)x(-4)x(-2)x(-1)$, $\rho_1 = x(-5)x(-4)$ and $\rho_2 = x(-2)x(-1)$. In the case of level 1 modules this is all that can happen, and, apart for the initial conditions discussed in Section 10, our proof of Theorem 6.5.2 would be finished.

However, in the case when level $k \geq 2$, say $k = 2$, we may have embeddings which are neither “adjacent” nor “disjoint”. For example, if $\pi = x(-3)^2x(-2)^2x(-1)$, $\rho_1 = x(-3)^2x(-2)$ and $\rho_2 = x(-2)^2x(-1)$, we can not apply Lemmas 9.2 and 9.3 directly. Of course, we can still apply Lemma 9.2 in two steps by considering an “intermediate” embedding $\rho_3 \subset \pi$ with $\rho_3 = x(-3)x(-2)^2$, and $\rho_1 \subset \pi$ and $\rho_2 \subset \pi$ is just the example of what we call linked embeddings. But this kind of argument will not work for $\pi = x(-3)^2x(-2)x(-1)^2$, $\rho_1 = x(-3)^2x(-2)$ and $\rho_2 = x(-2)x(-1)^2$, an example of what we call the exceptional case, listed in Lemma 7.3.1 within the “series” (12) of exceptional cases.

Similar situation appears in the principal picture for modules of level $k \geq 4$, cf. the proof of Lemma 9.18 in [MP], where both the linked embeddings and the exceptional cases are essentially of the form described above. Here the presence of three colors x , h and y leads to quite a few exceptional cases.

We shall prove Theorem 9.1 by induction on length $\ell(\pi)$. If $\ell(\pi) = k + 2$, then (9.1) holds by Lemma 9.2.

It is clear that if (9.1) holds for $\rho_1, \rho_2 \subset \pi$, then it holds as well for $\rho_1, \rho_2 \subset \kappa\pi$ for any $\kappa \in \mathcal{P}(\bar{B})$. In particular it is enough to prove (9.1) for any two embeddings $\rho_1, \rho_2 \subset \rho_1 \cup \rho_2$. The case when $\rho_1 \cap \rho_2 = \emptyset$ is proved by Lemma 9.3.

If (9.1) holds for every colored partition of length less than s , then (almost by definition) (9.1) holds for two linked embeddings $\rho_1 \subset \pi$, $\rho_2 \subset \pi$, $\ell(\pi) = s$. By the above remarks it is enough to prove (9.1) in the case when $\pi = \rho_1 \cup \rho_2$, $\rho_1 \cap \rho_2 \neq \emptyset$, $k + 3 \leq \ell(\pi) \leq s$, $\rho_1 \subset \pi$ and $\rho_2 \subset \pi$ are not linked, i.e. in the exceptional cases (1)–(14) listed in Lemma 7.3.1. In the proof of (9.1) for (some of) these exceptional cases we shall use the following two lemmas:

Lemma 9.4. *Let*


$$u(\rho \subset \pi) = \sum C_{\rho', \pi'} u(\rho' \subset \pi'),$$

where for each $\pi' \prec \pi$ the term $u(\rho' \subset \pi')$ appears for only one embedding $\rho' \subset \pi'$. Then $C_{\pi'} = C_{\rho', \pi'} = 0$ for $\pi' \prec \pi$. In particular


$$u(\rho \subset \pi) \in \sum C_{\rho', \pi} u(\rho' \subset \pi) + \mathcal{R}_{(\pi)}.$$

Proof. The statement follows from the fact that the right hand side has the unique leading term π (Proposition 6.4.3). $\square$

Example. As an example of the proof of (9.1) in the exceptional case let us consider $\pi = x(-3)^2x(-2)x(-1)^2$, $\rho_1 = x(-3)^2x(-2)$ and $\rho_2 = x(-2)x(-1)^2$. Here we have only one color x and $\text{sh } \pi = (-3)^2(-2)(-1)^2$. Note that $(-2)^5 \prec (-3)(-2)^3(-1) \prec (-3)^2(-2)(-1)^2$ are all plain partitions of degree -10 and $\preceq \text{sh } \pi$ and we can visualize them by Young diagrams



Since we want to prove (9.1), it will be enough to consider terms $u(\rho' \subset \pi')$ with $\pi' \preceq \pi$, which will amount to $\text{sh } \pi' \preceq \text{sh } \pi$, and all other terms will be denoted as $\dots$. We can also visualize the shapes of the leading terms of $r_{3\alpha}(-4) = r(x(-2)x(-1)^2)$, $r_{3\alpha}(-5) = r(x(-2)^2x(-1))$, $r_{3\alpha}(-6) = r(x(-2)^3)$, $r_{3\alpha}(-7) = r(x(-3)x(-2)^2)$ and $r_{3\alpha}(-8) = r(x(-3)^2x(-2))$ respectively as



It is clear that these are all possible $\text{sh } \rho'$ for $\rho' \subset \pi'$ with $\text{sh } \pi' \preceq \text{sh } \pi$ as above.

Now we start with $u(\rho_2 \subset \pi) = x(-3)^2r_{3\alpha}(-4)$. First we use the relation (8.1) with $k = 2$ and $n = -7$, i.e.

$$\sum_{j \in \mathbb{Z}} (4j + 7) x(j) r_{3\alpha}(-7 - j) = 0,$$

which we write as

$$x(-3)r_{3\alpha}(-4) = -\frac{1}{5}x(-2)r_{3\alpha}(-5) + \frac{3}{5}x(-1)r_{3\alpha}(-6) + \dots,$$

and we get

$$x(-3)^2r_{3\alpha}(-4) = -\frac{1}{5}x(-3)x(-2)r_{3\alpha}(-5) + \frac{3}{5}x(-3)x(-1)r_{3\alpha}(-6) + \dots$$

Now we apply the relation (8.1) again:

$$\begin{aligned} x(-3)r_{3\alpha}(-5) &= 0 \cdot x(-2)r_{3\alpha}(-6) + \frac{4}{4}x(-1)r_{3\alpha}(-7) + \dots, \\ x(-3)r_{3\alpha}(-6) &= \frac{1}{3}x(-2)r_{3\alpha}(-7) + \frac{5}{3}x(-1)r_{3\alpha}(-8) + \dots, \end{aligned}$$

and we get

$$\begin{aligned} x(-3)^2r_{3\alpha}(-4) &= -\frac{1}{5} \cdot 0 \cdot x(-2)^2r_{3\alpha}(-6) \\ &\quad + \left(\frac{3}{5} \cdot \frac{1}{3} - \frac{1}{5} \cdot \frac{4}{4}\right) \cdot x(-2)x(-1)r_{3\alpha}(-7) \\ &\quad + \frac{3}{5} \cdot \frac{5}{3} \cdot x(-1)^2r_{3\alpha}(-8) + \dots \\ &= x(-1)^2r_{3\alpha}(-8) + \dots \end{aligned}$$

Hence (9.1) holds.

In the above proof we can avoid the explicit calculation of coefficients: it is sufficient to know that, after applying the relation (8.1) in two steps on terms of the form $x(-3)r_{3\alpha}(n)$, we get a relation of the form

$$\begin{aligned}
u(\rho_2 \subset \pi) &= x(-3)^2 r_{3\alpha}(-4) \\
&= C_1 x(-2)^2 r_{3\alpha}(-6) \\
&\quad + C_2 x(-2)x(-1)r_{3\alpha}(-7) \\
&\quad + C_3 x(-1)^2 r_{3\alpha}(-8) + \dots \\
&= C_1 u(x(-2)^3 \subset x(-2)^5) \\
&\quad + C_2 u(x(-3)x(-2)^2 \subset x(-3)x(-2)^3 x(-1)) \\
&\quad + C_3 u(\rho_1 \subset \pi) + \dots
\end{aligned}$$

Then Lemma 9.4 implies $C_1 = C_2 = 0$. After that our choice of $r(\pi_{3\alpha}(n)) = r_{3\alpha}(n)$ also implies $C_3 = 1$. A general case of this type is treated in the proof of exceptional case (10).

As it was mentioned in the Introduction, and illustrated by the previous example, in some cases Lemma 9.4 enables us to prove (9.1) by first expressing $u(\rho_2 \subset \pi) - u(\rho_1 \subset \pi)$ as a linear combination of terms $u(\rho' \subset \pi')$, making sure afterwards that only terms for $\pi' \succ \pi$ appear nontrivially. In doing this we shall have to keep track of all the terms that may appear, and the following lemma will be used to check that the expression we obtain contains for each $\pi' \prec \pi$ at most one term of the form $u(\rho' \subset \pi')$.

Lemma 9.5. *Let π be a colored partition. Let π_1 be another colored partition such that $\text{sh } \pi_1 = \text{sh } \pi$, $\text{wt } \pi_1 = \text{wt } \pi$. Let $\rho \subset \pi_1$ be an embedding.*

- (2) *If $\pi = x(j-1)^{k+1-a-b}y(j)^a h(j)^b x(j)^a$, $a \geq 2$, $a+b \leq k$, and if $\pi_1 = y(j-1)^f h(j-1)^e y(j)^d h(j)^c \rho$, then $y(j)$ is not a part of ρ .*
- (3) *If $\pi = h(j-1)^{k+1-a-b}x(j-1)^a y(j)^b x(j)^{k+1-a}$, $a \geq 1$, and if $\pi_1 = y(j-1)^c h(j-1)^d x(j-1)^e y(j)^f h(j)^g \rho$, then $y(j)$ is not a part of ρ .*
- (4) *If $\pi = h(j-1)^{k+1-a-b}x(j-1)^a y(j)^b h(j)^{k+1-a-b}$, $1 \leq a+b \leq k$, and if $x(j)$ is not a part of π_1 , then $y(j-1)$ is not a part of ρ .*
- (5) *If $\pi = y(j-1)^{k+1-a}x(j-1)^b y(j)^a h(j)^{k+1-a-b}$, $a \geq 1$, and if $\pi_1 = h(j-1)^g x(j-1)^f y(j)^e h(j)^d x(j)^c \rho$, then $x(j-1)$ is not a part of ρ .*
- (6) *If $\pi = y(j-1)^a h(j-1)^b x(j-1)^a y(j)^{k+1-a-b}$, $a \geq 2$, $a+b \leq k$, and if $\pi_1 = h(j-1)^c x(j-1)^d h(j)^e x(j)^f \rho$, then $x(j-1)$ is not a part of ρ .*
- (11) *If $\pi = x(j-2)^{k+1-a-b}h(j-1)^a x(j-1)^b y(j)^c h(j)^{k+1-b-c}$, $a, b \geq 1$, $a+b+c \leq k$, and if $\pi_1 \prec \pi$, then $y(j-1)$ is not a part of ρ .*
- (12) *If $\pi = x(j-2)^{k+1-a-b}h(j-1)^a x(j-1)^b h(j)^c x(j)^{k+1-b-c}$, $a, b \geq 1$, $a+b \leq k$, and if $\pi_1 \prec \pi$, then $y(j-1)$ is not a part of ρ .*

Remark. Note that (2) and (3) have dual statements (6) and (5) (respectively) and that the dual proofs hold. Also note that the statement (4) is not self dual.

Proof. (2) The statement is clear since $\rho = \dots y(j)^p h(j)^q$ for $p \geq 1$, $q \geq 0$, $\text{sh } \rho = (j-1)^{k+1-p-q}j^{p+q}$, would imply $\text{wt}(\pi_1) < (k+1-a-b)\alpha = \text{wt}(\pi)$.

(3) The statement is clear since $\rho = \dots y(j)^p h(j)^q$ for $p \geq 1$, $q \geq 0$, $\text{sh } \rho = (j-1)^{k+1-p-q}j^{p+q}$, would imply $\text{wt}(\pi_1) < (k+1-b)\alpha = \text{wt}(\pi)$.

(4) By assumptions π_1 must be of the form $y(j-1)^c h(j-1)^e x(j-1)^f y(j)^d h(j)^g$, where $c + e + f = k + 1 - b$, $\text{wt}(\pi) = (a - b)\alpha = (-c - d + f)\alpha$. Let

$$\rho = y(j-1)^{c'} h(j-1)^{k+1-c'-d'} y(j)^{d'} \subset \pi_1$$

for some $c' \geq 1$. Then $c \geq c' \geq 1$, $e \geq k + 1 - c' - d'$, $d \geq d'$, and in particular $c + e \geq k + 1 - d'$. But then $d < d + a + c = f + b = k + 1 - (c + e) \leq d' \leq d$, a contradiction.

(6) The statement is clear since $\rho = h(j-1)^q x(j-1)^p \cdots$ for $p \geq 1$, $q \geq 0$, $\text{sh } \rho = (j-1)^{p+q} j^{k+1-p-q}$, would imply $\text{wt}(\pi_1) > -(k+1-a-b)\alpha = \text{wt}(\pi)$.

(11) Assume that $y(j-1)^d \subset \pi_1$ for some $d \geq 1$. Since by assumption $\pi_1 \prec \pi$, $\text{wt } \pi_1 = \text{wt } \pi$, we have that π_1 contains $y(j)^{c+i} h(j)^{k+1-b-c-i}$ for some $i \geq 1$. Because of $\text{wt } \pi_1 = \text{wt } \pi$ we also have that π_1 contains either

$$x(j-2)^{k+1-a-b} y(j-1)^s h(j-1)^{a-i-2s} x(j-1)^{b+i+s}, \quad s \geq d, \quad \text{or}$$

$$y(j-2)^r h(j-2)^p x(j-2)^{k+1-a-b-r-p} \cdot y(j-1)^{s-r} h(j-1)^{a-i-2s-p} x(j-1)^{b+i+s+r+p}, \quad s-r \geq d.$$

Now the assumption $\rho \subset \pi_1$, ρ of the form $\rho = y(j-1)^{d'} h(j-1)^{e'} y(j)^{k+1-e'-d'}$ would imply $k+1-e'-d' \leq c+i$ and

$$d' + e' \leq s + (a - i - 2s) \quad \text{or} \quad d' + e' \leq (s - r) + (a - i - 2s - p),$$

which in turn would imply $k+1 \leq c+a \leq c+a+b$, contrary to our assumption.

The possible form for π_1 shows that $x(j-1) \subset \pi_1$, so there is no embedding of the form $\rho \subset \pi_1$, $y(j-1) \subset \rho$, $\text{sh } \rho = (j-2)^{k+1-a-b} (j-1)^{a+b}$.

(12) Assume that $y(j-1)^d \subset \pi_1$ for some $d \geq 1$. Since by assumption $\pi_1 \prec \pi$, $\text{wt } \pi_1 = \text{wt } \pi$, we have that π_1 contains $y(j)^i h(j)^{c-i+m} x(j)^{k+1-b-c-m}$ for some $i, m \geq 0$, $i+m \geq 1$. Because of $\text{wt } \pi_1 = \text{wt } \pi$ we also have that π_1 contains either

$$x(j-2)^{k+1-a-b} y^s h(j-1)^{a-2s-m-i} x(j-1)^{b+m+i+s}, \quad s \geq d, \quad \text{or}$$

$$y(j-2)^r h(j-2)^p x(j-2)^{k+1-a-b-r-p} \cdot y(j-1)^{s-r} h(j-1)^{a-2s-m-i-p} x(j-1)^{b+m+i+s+r+p}, \quad s-r \geq d.$$

Now the assumption $\rho \subset \pi_1$, ρ of the form $\rho = y(j-1)^{d'} h(j-1)^{e'} y(j)^{k+1-e'-d'}$ would imply $k+1-e'-d' \leq i$ and

$$d' + e' \leq s + (a - 2s - m - i) \quad \text{or} \quad d' + e' \leq (s - r) + (a - 2s - m - i - p),$$

which in turn would imply $k+1 \leq a \leq a+b$, contrary to our assumption.

Since $\text{wt } \pi_1 = \text{wt } \pi$ and $x(j)^f \subset \pi_1$ for $f \leq k+1-b-c$, it must be that $x(j-1)^b \subset \pi_1$. Hence there is no embedding of the form

$$\rho \subset \pi_1, \quad y(j-1) \subset \rho, \quad \text{sh } \rho = (j-2)^{k+1-a-b} (j-1)^{a+b}. \quad \square$$

By using relations among relations constructed in Section 8 we shall prove (9.1) by induction on length $\ell(\pi)$ for all exceptional cases. Note that in the exceptional

cases we have only two embeddings $\rho_1 \subset \pi$, $\rho_2 \subset \pi$. At some places in the proof it will be convenient to write

$$a \underset{\text{sh}}{\sim} b \quad \text{or} \quad a - b \underset{\text{sh}}{\sim} 0$$

if $\text{sh } \ell(a) = \text{sh } \ell(b)$ and

$$a - b \in \overline{\mathbb{F}\text{-span}\{u(\rho \subset \pi) \mid \rho \in \ell(\bar{R}), \pi \in (\ell(\bar{R})), \rho \subset \pi, \text{sh } \pi \succ \text{sh } \ell(a)\}}.$$

At some places in the proof we shall use relations (8.5), $n = (k+2)j$, $|i| \leq k+1$, written in the form

$$(9.5) \quad \begin{aligned} (k+2-i)x(j)r_{(i-1)\alpha}((k+1)j) + ih(j)r_{i\alpha}((k+1)j) \\ + (k+2+i)y(j)r_{(i+1)\alpha}((k+1)j) \underset{\text{sh}}{\sim} 0. \end{aligned}$$

Now we consider all exceptional cases:

$$(1) \quad \begin{aligned} \pi &= y(j)^a h(j)^{k+1-a} x(j)^a, \quad 2 \leq a \leq k, \\ \rho_1 &= y(j)^a h(j)^{k+1-a}. \end{aligned}$$

By using relations (8.5), $n = (k+2)j$, $i \in [-(a-1), a-1]$, of the form (9.5) we get

$$x(j)^a r(\rho_1) \underset{\text{sh}}{\sim} \sum_{\substack{b, c \geq 0 \\ b+c=a}} C_{bc} y(j)^b h(j)^c r_{b\alpha}((k+1)j).$$

Now (9.1) follows by applying Lemma 9.4.

$$(2) \quad \begin{aligned} \pi &= x(j-1)^{k+1-a-b} y(j)^a h(j)^b x(j)^a, \quad a \geq 2, a+b \leq k, \\ \rho_1 &= x(j-1)^{k+1-a-b} y(j)^a h(j)^b. \end{aligned}$$

By using in a steps relations (8.6) of the form $C_x = 1, C^x = 0$, or (9.5), we get

$$(9.6) \quad x(j)^a r(\rho_1) \underset{\text{sh}}{\sim} \sum C_{cdef} y(j-1)^f h(j-1)^e y(j)^d h(j)^c r(\rho),$$

where $c+d+e+f = a$, $\rho \in \ell(\bar{R})$. (Note that the leading term ρ of $r(\rho)$ is completely determined by c, d, e, f .)

Note that each time when we apply one of the relations (8.6) or (9.5) we replace $x(j)r(\rho)$ by a combination of terms of the form $Xr(\rho')$, where X is either $y(j), h(j), y(j-1)$ or $h(j-1)$ and $\text{wt } \rho' \leq \text{wt } \rho + 2\alpha$. Hence after applying relations i times, $i < a$, all $r(\rho)$'s that appear have weights bounded by $(k+1-2a-b)\alpha + 2i\alpha < (k+1-b)\alpha$ and we see that there exists a relation of the form (8.6) or (9.5) which could be applied again. (In the subsequent cases we shall state the type of relations we use and we shall omit the proof that this can be done.)

By Lemma 9.5(2) $y(j)$ is not a part of ρ , i.e.

$$r(\rho) = r(x(j-1)^p h(j)^q x(j)^{k+1-p-q}).$$

Now we can apply Lemma 9.4 and (9.1) follows since $r(\rho_1)$ does not appear on the right hand side of (9.6).

$$(3) \quad \begin{aligned} \pi &= h(j-1)^{k+1-a-b} x(j-1)^a y(j)^b x(j)^{k+1-a}, \quad 1 \leq a \leq k-1, \\ \rho_1 &= h(j-1)^{k+1-a-b} x(j-1)^a y(j)^b. \end{aligned}$$

By using in $k+1-a$ steps relations: a) relations (8.6) of the form $C_x = 1, C_h = 0$ on the terms $x(j)r(\rho)$, $\text{sh } \rho = (j-1)^p j^{k+1-p}$ for $p \geq 1$, b) relations (9.5) on the terms $x(j)r(\rho)$, $\text{sh } \rho = j^{k+1}$, $\text{wt } \rho < k+1$, c) no relation otherwise, we get

$$(9.7) \quad \begin{aligned} x(j)^{k+1-a} r(\rho_1) &\underset{\text{sh}}{\sim} \sum C_{cdef} y(j-1)^c h(j-1)^d x(j-1)^e y(j)^f r(\rho) \\ &+ \sum C_{cdefgh} y(j-1)^c h(j-1)^d x(j-1)^e y(j)^f h(j)^g x(j)^h r(\rho). \end{aligned}$$

In the first sum in (9.7) we have $\text{sh } \rho = (j-1)^p j^{k+1-p}$, $p \geq 1$, and Lemma 9.5(3) implies that

$$r(\rho) = r(x(j-1)^p h(j)^q x(j)^{k+1-p-q}).$$

In the second sum in (9.7) we have $\text{sh } \rho = j^{k+1}$. If $h \geq 1$ then $\rho = x(j)^{k+1}$ since in last steps we used c), i.e. we didn't use any relation. If $h = 0$ then Lemma 9.5(3) implies that

$$r(\rho) = r(h(j)^q x(j)^{k+1-q}).$$

Since the number of $y(j)$'s in terms of the first sum is greater than in terms of the second sum, we see that we can apply Lemma 9.4. Since $r(\rho_1)$ does not appear on the right hand side of (9.7), we get (9.1).

$$(4) \quad \begin{aligned} \pi &= h(j-1)^{k+1-a-b} x(j-1)^a y(j)^b h(j)^{k+1-a-b}, \quad 1 \leq a+b \leq k-1, \\ \rho_1 &= x(j-1)^a y(j)^b h(j)^{k+1-a-b}. \end{aligned}$$

By using in $k+1-a-b$ steps relations (8.6) of the form $C^h = 1, C^x = 0$ we get

$$(9.8) \quad h(j-1)^{k+1-a-b} r(\rho_1) \underset{\text{sh}}{\sim} \sum C_{cdef} y(j-1)^c y(j)^d h(j)^e x(j)^f r(\rho).$$

We need not consider terms with $f > 0$ (since for those the leading terms are $\succ \pi$). Consider two classes of terms with a leading term π_1 : (A) $y(j-1)^c \subset \pi_1$ for some $c > 0$ and (B) terms of the form $r(\rho)y(j)^d h(j)^e$, $x(j)$ is not a part of ρ (since otherwise $\pi_1 \succ \pi$). By Lemma 9.5(4) and $\text{wt}(\pi) = \text{wt}(\pi_1)$, in the case (B) we have either

$$\begin{aligned} \rho &= h(j-1)^{k+1-b-p} x(j-1)^p y(j)^b, \quad 0 < p \leq k-b, \quad \text{or} \\ \rho &= x(j-1)^{k+1-b} y(j)^b, \quad \pi_1 = x(j-1)^{k+1-b} y(j)^{k+1-a}. \end{aligned}$$

In either case for given π_1 there is only one embedding. In the case (A) we may apply the induction hypothesis on $\pi_1/y(j-1)^c$. Hence we may apply Lemma 9.4 and (9.1) follows.

Remark. The reader may find a “dual” proof of case (4) a bit shorter, but some changes are needed. For example, the argument dual to: “We need not consider terms with $f > 0$ (since for those the leading terms are $\succ \pi$)” is false. However, in some cases (like (5) and (6)) dual arguments hold all the way through the proof.

(5) The proof is dual to the proof of (3).

(6) The proof is dual to the proof of (2).

(7) Let $\pi = y(j-1)^{k+1-a-c}h(j-1)^c x(j-1)^b y(j)^a h(j)^{k+1-a-b}$, $a \geq 1$, $a+b+c \leq k$. Note that we have already considered the case $c = 0$ in (5). We keep the shape

$$(j-1)^{k+1+b-a} j^{k+1-b}$$

fixed and prove by induction on c , $0 \leq c \leq k-a$, that embedding of the form (for $c \geq 1$)

$$(9.9) \quad y(j-1)^{k+1-a-c} h(j-1)^c r(x(j-1)^b y(j)^a h(j)^{k+1-a-b})$$

is, modulo terms with the leading term $\succ \pi$, proportional to

$$(9.10) \quad x(j-1)^b h(j)^{k+1-a-b} r(y(j-1)^{k+1-a-c} h(j-1)^c y(j)^a).$$

By the induction hypothesis the embedding of the form

$$(9.11) \quad y(j-1)^{k+2-a-c} h(j-1)^{c-1} r(x(j-1)^b y(j)^a h(j)^{k+1-a-b})$$

with the leading term π' is proportional to

$$(9.12) \quad x(j-1)^b h(j)^{k+1-a-b} r(y(j-1)^{k+2-a-c} h(j-1)^{c-1} y(j)^a)$$

modulo terms with the leading term $\succ \pi'$.

Now we apply the adjoint action of x on the relation involving (9.11) and (9.12).

The action of x on (9.11) gives three terms, one is (proportional to) (9.9), the other two have the leading terms bigger than π . The action of x on (9.12) gives (9.10) and another term $\succ \pi$.

What remains to show is that the action of x on any term (in the relation involving (9.11) and (9.12)) with the leading term $\pi'' \succ \pi'$ will give terms $\succ \pi$.

If π'' contains $x(j)$, then all terms obtained by the action of x contain $x(j)$, the result being $\succ \pi$.

If π'' contains $y(j)^{a-d} h(j)^{k+1-a-b+d}$, $d \geq 1$, then the action of x leaves this term, gives one $x(j)$ more and one $h(j)$ less, or gives one $h(j)$ more and one $y(j)$ less. In either case the result is $\succ \pi$.

If π'' contains $x(j-1)^{b+d} y(j)^a h(j)^{k+1-a-b}$, $d \geq 0$, then the action of x leaves this term or gives one $x(j)$, $h(j)$ or $x(j-1)$ more. In either case the result is $\succ \pi$ when $d \geq 1$. In the case $d = 0$ we have $\pi'' = \pi'$.

(8) Let $\pi = h(j-1)^{k+1-a-b}x(j-1)^ay(j)^bh(j)^cx(j)^{k+1-a-c}$, $a \geq 1$, $a+b+c \leq k$. Note that we have already considered the case $c = 0$ in (3). We keep the shape

$$(j-1)^{k+1-b}j^{k+1+b-a}$$

fixed and prove by induction on c , $0 \leq c \leq k-a-b$, that embedding of the form (for $c \geq 1$)

$$(9.13) \quad h(j)^cx(j)^{k+1-a-c}r(h(j-1)^{k+1-a-b}x(j-1)^ay(j)^b)$$

is, modulo terms with the leading term $\succ \pi$, proportional to

$$(9.14) \quad h(j-1)^{k+1-a-b}y(j)^br(x(j-1)^ah(j)^cx(j)^{k+1-a-c}).$$

By induction hypothesis the embedding of the form

$$(9.15) \quad h(j)^{c-1}x(j)^{k+2-a-c}r(h(j-1)^{k+1-a-b}x(j-1)^ay(j)^b)$$

with the leading term π' is proportional to

$$(9.16) \quad h(j-1)^{k+1-a-b}y(j)^br(x(j-1)^ah(j)^{c-1}x(j)^{k+2-a-c})$$

modulo terms with the leading term $\succ \pi'$. Now we apply the adjoint action of y on the relation involving (9.15) and (9.16).

The action of y on (9.15) gives three terms, one is (proportional to) (9.13), the other two (with the leading terms) $\succ \pi$. The action of y on (9.16) gives (9.14) and another term $\succ \pi$.

What remains to show is that the action of y on any term (in the relation involving (9.15) and (9.16)) with the leading term $\pi'' \succ \pi'$ will give terms $\succ \pi$.

First let $\pi'' \supset y(j)^bh(j)^{c-1}x(j)^{k+2-a-c}$, i.e.

$$\pi'' = y(j-1)^dh(j-1)^{k+1-a-b-2d}x(j-1)^{a+d}y(j)^bh(j)^{c-1}x(j)^{k+2-a-c},$$

where $d \geq 1$ and $\rho \subset \pi''$. If $x(j) \notin \rho$, the partition (with the obvious notation) $(y \cdot \rho)(\pi''/\rho)$ contains $x(j)^{k+2-a-c}$, so is $\succ \pi$. If y does not act on any $x(j) \in \pi''/\rho$, we again obtain a term $\succ \pi$. If y acts on $x(j)$, we obtain the term (with the leading term):

$$(9.17) \quad y(j-1)^dh(j-1)^{k+1-a-b-2d}x(j-1)^{a+d}y(j)^bh(j)^cx(j)^{k+1-a-c} \succ \pi.$$

If $x(j) \in \rho$, then we see that $(y \cdot \rho)(\pi''/\rho)$ is of the form (9.17). Also we see that the action of y on π''/ρ will give terms $\succ \pi$.

Now let $\pi'' \supset y(j)^{b-d}h(j)^{c-1+d}x(j)^{k+2-a-c}$, where $d \geq 1$ and $\rho \subset \pi''$. If $x(j) \notin \rho$, then $(y \cdot \rho)(\pi''/\rho)$ contains $x(j)^{k+2-a-c}$, so is $\succ \pi$. If y does not act on $x(j) \in \pi''/\rho$, we again obtain a term $\succ \pi$. If y acts on $x(j)$, we obtain the term (which contains)

$$(9.18) \quad \dots h(j)^{c-1+d}h(j)x(j)^{k+1-a-c} \succ \pi.$$

If $x(j) \in \rho$, then $y \cdot \rho$ has one less $x(j)$ and one more $h(j)$, so $(y \cdot \rho)(\pi''/\rho)$ is of the form (9.18). Also we see that the action of y on π''/ρ will give terms $\succ \pi$.

Finally, if $\pi'' \supset x(j)^{k+3-a-c}$, then the action of y will give terms that contain $x(j)^{k+2-a-c}$ and hence $\succ \pi$.

(9) and (10). We fix a shape $(j-2)^a(j-1)^{k+1-a}j^a$, $2 \leq a \leq k$. Let

$$\begin{aligned}\pi &= y(j-2)^a y(j-1)^{k+1-a} y(j)^a, \\ \rho_1 &= y(j-1)^{k+1-a} y(j)^a.\end{aligned}$$

By using relations (8.4), $i = -k-2$, of the form

$$y(j-2)r_{-(k+1)\alpha}(m) + c_1 y(j-1)r_{-(k+1)\alpha}(m-1) + c_2 y(j)r_{-(k+1)\alpha}(m-2) + \dots = 0,$$

where $m = (k+1)(j-1) \pm r$, $0 \leq r < k+1$, c_1, c_2 are some rational constants and $\dots$ denotes the sum of shorter elements (i.e. elements of length $k+1$) and terms which contain a factor $y(p)$, $p > j$ or $p < j-2$, we get

$$(9.19) \quad y(j-2)^a r(\rho_1) = \sum C_{bc} y(j)^b y(j-1)^c r(\rho) + A + B.$$

Here $b+c = a$; the terms in A contain a factor $y(p)$, $p > j$ (and their leading terms are greater than π); the terms B contain a factor $y(p)$, $p < j-2$.

If π_1 is a leading term of a summand in B , $\pi_1 \prec \pi$, then for any embedding $\rho \subset \pi_1$ the term $y(p)$, $p < j-2$, is not a part of ρ . By induction hypothesis (on length) for two embeddings in such π_1 (i.e. in $\pi_1/y(p)$) relation (9.1) holds, so we may rewrite B so that for each $\pi_1 \prec \pi$ that appears there is only one embedding.

In (9.19) we have that $r(\rho) = r(y(j-2)^b y(j-1)^{k+1-b})$, so each leading term appears with only one embedding. Hence we can apply Lemma 9.4 and we get

$$y(j-2)^a r(\rho_1) \in \mathbb{F}^\times y(j)^a r(y(j-2)^a y(j-1)^{k+1-a}) + \mathcal{R}_{(\pi)}.$$

But $\pi' \succ \pi$ implies $\text{sh}(\pi') \succ (j-2)^a(j-1)^{k+1-a}j^a$, and hence

$$(9.20) \quad y(j-2)^a r(y(j-1)^{k+1-a} y(j)^a) \underset{\text{sh}}{\sim} y(j)^a r(y(j-2)^a y(j-1)^{k+1-a}).$$

Since the adjoint action of x does not change the shape, we get (9.1) in the case (9) and (10) by adjoint action of x .

(11) and (12). Let $a \geq 0$, $b \geq 1$ and

$$\begin{aligned}\pi &= x(j-2)^{k+1-a-b} h(j-1)^a x(j-1)^b P, \\ \rho_1 &= x(j-1)^b P, \quad \text{where} \\ P &= y(j)^c h(j)^{k+1-b-c}, \quad a+b+c \leq k, \quad \text{or} \\ P &= h(j)^c x(j)^{k+1-b-c}, \quad a+b \leq k.\end{aligned}$$

Note that the case $a = 0$ follows from (9.20) by adjoint action of x . By using in a steps relations (8.6) of the form $C^h = 1$, $C^x = 0$ we get

$$(9.21) \quad \sum C_{defg} x(j-2)^{k+1-a-b} h(j-1)^a r(\rho_1) \underset{\text{sh}}{\sim} x(j)^d y(j)^e h(j)^f x(j)^g r(\rho),$$

where $0 \leq d \leq a$. It will be enough to consider only the terms with the leading term $\pi_1 \prec \pi$.

Consider three disjoint classes of $\pi_1 \prec \pi$: (A) $y(j-1)^d \subset \pi_1$ for some $d \geq 1$, (B) π_1 is not in A and $y(j-2)^p h(j-2)^q x(j-2)^{k+1-a-b-p-q} \subset \pi_1$ for some $p+q \geq 1$ and (C) π_1 is not in A and $x(j-2)^{k+1-a-b} \subset \pi_1$.

Let π_1 be in the class A.

For two embeddings $\rho, \rho' \subset \pi_1$ (i.e. $\rho, \rho' \subset \pi_1/y(j-1)^d$ by Lemma 9.5) by induction hypothesis on length terms $u(\rho \subset \pi_1)$ and $u(\rho' \subset \pi_1)$ are proportional modulo higher terms (with leading terms $\succ \pi_1$). These higher terms are again in class A.

Let π_1 be in the class B.

For two embeddings $\rho, \rho' \subset \pi_1$ ($\rho, \rho' \subset \pi_1/y(j-2)^p h(j-2)^q x(j-2)^{k+1-a-b-p-q}$ because of Lemma 9.5), by induction hypothesis on length, terms $u(\rho \subset \pi_1)$ and $u(\rho' \subset \pi_1)$ are proportional modulo higher terms (with leading terms $\succ \pi_1$). These higher terms are again in class B.

Now consider a term in (9.21) which has the leading term π_1 in the class C. Then $d = 0$. For $\rho \subset \pi_1$ (i.e. $\rho \subset \pi_1/y(j)^e h(j)^f x(j)^g$) the term $u(\rho \subset \pi_1)$ is (by using relations of the form (10), (11) or (12)) proportional to

$$y(j)^{e'} h(j)^{f'} x(j)^{g'} r(x(j-2)^{k+1-a-b} h(j-1)^{a'} x(j-1)^{b'})$$

modulo terms with leading terms $\pi' \succ \pi_1$.

Hence in a finite number of steps we can rewrite (first the terms in class C, and then the terms in class A and B) the right hand side of (9.21), modulo $\mathcal{R}_{(\pi)}$, as a linear combination of terms $u(\rho \subset \pi_1)$, where for each $\pi_1 \prec \pi$ there is only one embedding. By applying Lemma 9.4 we get (9.1).

(13) and (14). Let $a \geq 0, b \geq 1$ and

$$\begin{aligned} \pi &= P y(j-1)^b h(j-1)^a y(j)^{k+1-a-b}, \\ \rho_1 &= P y(j-1)^b, \quad \text{where} \\ P &= h(j-2)^{k+1-b-c} x(j-2)^c, \quad a+b+c \leq k, \quad \text{or} \\ P &= y(j-2)^{k+1-b-c} h(j-2)^c, \quad a+b \leq k. \end{aligned}$$

Note that the case $a = 0$ follows from (9.20) by adjoint action of x . Let us also remark that $\text{sh } \pi_1 = \text{sh } \pi$, $\pi_1 \prec \pi$, implies that for some $i \geq 0$

$$\pi_1 \supset y(j-1)^{b+i} h(j-1)^{a-i} y(j)^{k+1-a-b} = \rho'.$$

By using in a steps the relation (8.6) of the form $C_h = 1, C_y = 0$ we get

$$(9.22) \quad \begin{aligned} &h(j-1)^a y(j)^{k+1-a-b} r(\rho_1) \underset{\text{sh}}{\sim} \\ &\sum C_{defg} y(j-2)^d h(j-2)^e x(j-2)^f x(j-1)^g y(j)^{k+1-a-b} r(\rho), \end{aligned}$$

where $0 \leq g \leq a$. For $g > 0$ the leading term π_1 is greater than π . By the previous remark in the case $g = 0$ we may use the induction hypothesis and express terms in (9.22), modulo $\mathcal{R}_{(\pi)}$, as a linear combination of elements of the form

$$y(j-2)^{d'} h(j-2)^{e'} x(j-2)^{f'} r(\rho'),$$

where $d' + e' + f' = k + 1 - b$. Then each leading term $\pi_1 \prec \pi$ appears with only one embedding and we may apply Lemma 9.4.

10. LINEAR INDEPENDENCE OF BASES OF STANDARD MODULES

Let $\Lambda = k_0\Lambda_0 + k_1\Lambda_1$, $k_0, k_1 \in \mathbb{N}$, and $k = k_0 + k_1$. Let v_Λ be a highest weight vector of Verma module $M(\Lambda)$. Let $M^1(\Lambda)$ be the maximal submodule of $M(\Lambda)$.

Recall that we may choose a map

$$\ell(\bar{R}v_\Lambda) \rightarrow \bar{R}, \quad \rho \mapsto r(\rho)$$

such that $\ell(r(\rho)v_\Lambda) = \rho$. For $\rho \in \ell(\bar{R}v_\Lambda)$ and $\pi \in (\ell(\bar{R}v_\Lambda))$, $\rho \subset \pi$, we shall say that $\rho \subset \pi$ is an *embedding*. For an embedding $\rho \subset \pi$ we set

$$u(\rho \subset \pi) = u(\pi/\rho)r(\rho).$$

Define a filtration $(M_{(\pi)}^1 \mid \pi \in (\ell(\bar{R}v_\Lambda)))$ on $M^1 = M^1(\Lambda) = U(\tilde{\mathfrak{g}})\bar{R}v_\Lambda = U(\tilde{\mathfrak{n}}_-)\bar{R}v_\Lambda$ by

$$M_{[\pi]}^1 = \mathbb{F}\text{-span}\{urv_\Lambda \mid u \in U(\tilde{\mathfrak{n}}_-), r \in \bar{R}, \ell(urv_\Lambda) \not\supseteq \pi\},$$

$$M_{(\pi)}^1 = \bigcup_{\pi' \supseteq \pi} M_{[\pi']}^1.$$

(At this point this filtration should not be confused with the filtration $(M^1 \cap M(\Lambda)_{[\pi]} \mid \pi \in (\ell(\bar{R})))$.)

The main result of this section is an analogue of Theorem 9.1:

Theorem 10.1. *Let $\kappa \in (\ell(\bar{R}v_\Lambda))$, $\rho_1, \rho_2 \in \ell(\bar{R}v_\Lambda)$, $\rho_1, \rho_2 \subset \kappa$. Then*

$$(10.1) \quad u(\rho_1 \subset \kappa)v_\Lambda \in \mathbb{F}^\times u(\rho_2 \subset \kappa)v_\Lambda + M_{(\kappa)}^1.$$

Remark. Note that Theorem 6.5.2 is a simple consequence of Theorem 10.1: Since $M^1(\Lambda) = U(\tilde{\mathfrak{n}}_-)\bar{R}v_\Lambda$, the set of vectors

$$u(\rho \subset \pi)v_\Lambda, \quad \rho \in \ell(\bar{R}v_\Lambda), \pi \in (\ell(\bar{R}v_\Lambda)), \rho \subset \pi,$$

is a spanning set of $M^1(\Lambda)$. Using Theorem 10.1 and induction we see that (6.5.1) is a spanning set as well. Clearly this set is linearly independent, and Theorem 6.5.2 follows.

The rest of this section is devoted to the proof of Theorem 10.1.

We may assume that the map $\rho \mapsto r(\rho)$ is such that $r(\rho)$ is homogeneous of weight $\text{wt}(\rho)$ and degree $|\rho|$. We will construct such a map $r: \ell(\bar{R}v_\Lambda) \rightarrow \bar{R}$ by starting with a map (denoted by the same r)

$$r: \ell(\bar{R}) \rightarrow \bar{R}, \quad \rho \mapsto r(\rho)$$

defined by $r(\pi_{m\alpha}(n)) = r_{m\alpha}(n)$ for $n \in \mathbb{Z}$, $|m| \leq k+1$ (cf. Section 6.6).

Proposition 10.2. *Let $a, b \in [0, k+1]$, $a+b \leq k+1$. Then*

- (a) $r(x(-1)^a h^{k+1-a-b} x^b)v_\Lambda = 0$ if $b > 0$,
- (b) $r(x(-1)^a y^b h^{k+1-a-b})v_\Lambda = 0$ if $a \leq k_0$ and $b \leq k_1$,
- (c) $\ell(r(x(-1)^a y^b h^{k+1-a-b})v_\Lambda) = x(-1)^a y^b$ if $a > k_0$ or $b > k_1$,
- (d) $\ell(r(h(-1)^a x(-1)^b y^{k+1-a-b})v_\Lambda) = h(-1)^a x(-1)^b y^{k+1-a-b}$,
 $\ell(r(y(-1)^a h(-1)^b y^{k+1-a-b})v_\Lambda) = y(-1)^a h(-1)^b y^{k+1-a-b}$.

Proof. First recall that

$$x \cdot y^p = [x, y^p] = py^{p-1}(h - (p-1)).$$

For $r = r(y(-1)^a y^{k+1-a})$ we have

$$\begin{aligned}
(10.2) \quad & (x^q \cdot r)v_\Lambda = x^q(rv_\Lambda) = \\
& = \sum_{p=0}^q \binom{q}{p} (x^{q-p} \cdot y(-1)^a)(x^p \cdot y^{k+1-a})v_\Lambda \\
& = \sum_{p=0}^q \binom{q}{p} (x^{q-p} \cdot y(-1)^a) p! \binom{k+1-a}{p} \\
& \quad \cdot y^{k+1-a-p}(k_1 - (k-a))(k_1 - (k-a) + 1) \cdots \\
& \quad \cdots (k_1 - k - 1 + a + p)v_\Lambda,
\end{aligned}$$

where $y^{k+1-a-p}v_\Lambda = 0$ for $k+1-a-p < 0$ and $x^{q-p} \cdot y(-1)^a = 0$ for $q-p > 2a$.

(a) Let $q = k+1+a+b = 2a+k+1-a+b$, $b > 0$. Then $k+1-a > 0$ and $x^p y^{k+1-a}v_\Lambda = 0$ for $p \geq k+1-a+b$, and from (10.2) we get $(x^q \cdot r)v_\Lambda = 0$.

(b) and (c). Let $q = k+1+a-b$. The leading term of $(x^q \cdot r)v_\Lambda$ will be a nontrivial summand in (10.2) for $k+1-a-p$ maximal possible, i.e. minimal possible $p \geq q-2a$. For $p = q-2a = k+1-a-b$ the interval in $\mathbb{Z}$

$$[k_1 - (k-a), k_1 - k - 1 + a + p] = [a - k_0, k_1 - b]$$

does not contain zero for $a - k_0 > 0$ or $k_1 - b < 0$, and (c) follows. Similarly, for $p \geq q-2a$ we have $k_1 - k - 1 + a + p \geq k_1 - b$. Hence $(x^q \cdot r)v_\Lambda = 0$ if $0 \in [a - k_0, k_1 - b]$ and (b) follows.

(d) is obvious. $\square$

Clearly $\ell(\bar{R}v_\Lambda)$ consists of elements $\pi_{m\alpha}(n) = \ell(r_{m\alpha}(n))$ for $n \leq -k-1$, $|m| \leq k+1$, and the elements listed in (c) and (d) of Proposition 10.2. We define a map $r: \ell(\bar{R}v_\Lambda) \rightarrow \bar{R}$, $\rho \mapsto r(\rho)$, by

$$(10.3) \quad r(x(-1)^a y^b) = r(x(-1)^a y^b h^{k+1-a-b})$$

for $a+b \leq k$, $a > k_0$ or $b > k_1$, and as previously defined r on the set $\ell(\bar{R}v_\Lambda) \cap \ell(\bar{R})$. We shall say that the leading term $\rho = x(-1)^a y^b \in \ell(\bar{R}v_\Lambda)$, $a+b \leq k$, is *short*. We shall also say that $\rho \in \ell(\bar{R}v_\Lambda)$, $\ell(\rho) = k+1$, is *long*. Obviously the long $\rho \in \ell(\bar{R}v_\Lambda)$ are characterized by equality $\ell(r(\rho)) = \ell(r(\rho)v_\Lambda)$.

By Theorem 9.1 we have the following:

Lemma 10.3. *For two embeddings $\rho_1, \rho_2 \subset \pi$, both ρ_1 and ρ_2 long, (10.1) holds.*

Relations (8.4) and (8.5) constructed in Section 8 can be written in the form

$$\begin{aligned}
& \sum_{j < 0} ((k+2)j-n) \{x(j)r_{(i-1)\alpha}(n-j)v_\Lambda - h(j)r_{i\alpha}(n-j)v_\Lambda - y(j)r_{(i+1)\alpha}(n-j)v_\Lambda\} \\
& - n(k_0 + 1 - i)r_{i\alpha}(n)v_\Lambda + nyr_{(i+1)\alpha}(n)v_\Lambda = 0,
\end{aligned}$$

$$\sum_{j < 0} \{ (k+2-i)x(j)r_{(i-1)\alpha}(n-j) + ih(j)r_{i\alpha}(n-j) + (k+2+i)y(j)r_{(i+1)\alpha}(n-j) \} v_{\Lambda} \\ + [(k+2)n + i(k_1 + 1 + i)]r_{i\alpha}(n)v_{\Lambda} + (k+2+i)yr_{(i+1)\alpha}(n)v_{\Lambda} = 0.$$

We shall also use linear combinations of (8.4) and (8.5) for $-k-1 \leq n \leq 0$ of the form

$$(10.4) \quad C_{\Lambda}r_{i\alpha}(n)v_{\Lambda} + C_y yr_{(i+1)\alpha}(n)v_{\Lambda} \\ + C^x x(-1)r_{(i-1)\alpha}(n+1)v_{\Lambda} + C^h h(-1)r_{i\alpha}(n+1)v_{\Lambda} + C^y y(-1)r_{(i+1)\alpha}(n+1)v_{\Lambda} \\ + \sum_{j \leq -2} a_j x(j)r_{(i-1)\alpha}(n-j)v_{\Lambda} + b_j h(j)r_{i\alpha}(n-j)v_{\Lambda} + c_j y(j)r_{(i+1)\alpha}(n-j)v_{\Lambda} = 0.$$

Lemma 10.4. *For two embeddings $\rho_1, \rho_2 \subset \pi$, both ρ_1 and ρ_2 short, (10.1) holds.*

Proof. Case 1. Let $\rho_1 = x(-1)^a y^b$, $a, b \geq 0$, $a + b \leq k$, $\rho_2 = x(-1)\rho_1$.

Then $r(\rho_1) = r_{(i-1)\alpha}(n+1)$ for $n+1 = -a$, $i-1 = a-b$. We can write (10.4) in the form

$$Cr(x(-1)^{a+1}y^b)v_{\Lambda} + C_y yr(x(-1)^{a+1}y^{b-1})v_{\Lambda} \\ + C^x x(-1)r(x(-1)^a y^b)v_{\Lambda} + C^h h(-1)r(x(-1)^a y^{b-1})v_{\Lambda} \\ + C^y y(-1)r(x(-1)^a y^{b-2})v_{\Lambda} + A = 0,$$

where A denotes a sum of terms with leading terms π_1 , $\ell(\pi_1) \leq a+b$. Now we choose $C^x = 1$, $C_y = 0$ and we get

$$(10.5) \quad x(-1)r(x(-1)^a y^b)v_{\Lambda} \in Cr(x(-1)^{a+1}y^b)v_{\Lambda} + M_{(x(-1)^{a+1}y^b)}^1.$$

Clearly (10.5) implies $C \neq 0$, so (10.1) holds.

Case 2. Let $\rho_1 = x(-1)^a y^b$, $a, b \geq 0$, $a + b \leq k$, $\rho_2 = y\rho_1$.

Then $r(\rho_1) = r_{(i+1)\alpha}(n)$ for $n = -a$, $i+1 = a-b$. We can write (10.4) in the form

$$Cr(x(-1)^a y^{b+1})v_{\Lambda} + C_y yr(x(-1)^a y^b)v_{\Lambda} \\ + C^x x(-1)r(x(-1)^{a-1}y^{b+1})v_{\Lambda} + C^h h(-1)r(x(-1)^{a-1}y^b)v_{\Lambda} \\ + C^y y(-1)r(x(-1)^{a-1}y^{b-1})v_{\Lambda} + A = 0,$$

where A denotes a sum of terms with leading terms π_1 , $\ell(\pi_1) \leq a+b$. Now we choose $C_y = 1$, $C^x = 0$ and we get

$$(10.6) \quad yr(x(-1)^a y^b)v_{\Lambda} \in Cr(x(-1)^a y^{b+1})v_{\Lambda} + M_{(x(-1)^a y^{b+1})}^1.$$

Clearly (10.6) implies $C \neq 0$, so (10.1) holds.

In the general case when $\rho_2 = x(-1)^{a'} y^{b'}$, $a', b' \geq 0$, $a' + b' \leq k$ we can find a sequence $\rho_p = x(-1)^{a_p} y^{b_p}$, $p = 1, \dots, s$, so that $(a, b) = (a_1, b_1)$, $(a', b') = (a_s, b_s)$, and that for ρ_p, ρ_{p+1} we can apply either Case 1, Case 2 or Lemma 10.3. $\square$

It remains to prove (10.1) for two embeddings $\rho_1, \rho_2 \subset \pi$, where one ρ_i is short, the other long. First we have

Lemma 10.5. *Let $\rho_1, \rho_2 \in \ell(\bar{R}v_\Lambda)$, $\pi = \rho_1 \rho_2$. Then*

$$u(\rho_1 \subset \pi)v_\Lambda \in \mathbb{F}^\times u(\rho_2 \subset \pi)v_\Lambda + M_{(\pi)}^1.$$

Proof. Because of Lemmas 10.3 and 10.4 it is enough to consider the case when ρ_1 is long, ρ_2 is short. In this case the proof is similar to the proof of Lemma 9.3. $\square$

We shall say that $\rho_1, \rho_2 \in \ell(\bar{R}v_\Lambda)$ are $\{y, x(-1)\}$ -linked if there exists a sequence of embeddings $\sigma_1, \dots, \sigma_p \subset \rho_1 \cup \rho_2$ such that $\sigma_1 = \rho_1$, $\sigma_p = \rho_2$ and

$$(10.7) \quad \text{either } \ell(\sigma_{i-1} \cup \sigma_i) < \ell(\rho_1 \cup \rho_2) \text{ for each } i = 2, \dots, p$$

$$(10.8) \quad \text{or } (\sigma_r \cup \sigma_{r+1})/\sigma_r \subset \{y, x(-1)\}, (\sigma_r \cup \sigma_{r+1})/\sigma_{r+1} \subset \{y, x(-1)\}.$$

Note that the assumption (10.8) together with Lemma 10.3, (10.5) or (10.6) imply for $\pi = \sigma_r \cup \sigma_{r+1}$ that

$$(10.9) \quad r(\sigma_r \subset \pi)v_\Lambda \in \mathbb{F}^\times r(\sigma_{r+1} \subset \pi)v_\Lambda + M_{(\pi)}^1.$$

By modifying the proof of Lemma 7.3.1 we get the following:

Lemma 10.6. *Let $\rho_1, \rho_2 \in \ell(\bar{R}v_\Lambda)$, ρ_1 short, ρ_2 long, $\rho_1 \cap \rho_2 \neq \emptyset$. Then ρ_1 and ρ_2 are not $\{y, x(-1)\}$ -linked only in the following cases*

$$(4) \quad \begin{aligned} \rho_1 &= x(-1)^a y^b, & \rho_2 &= h(-1)^{k+1-a-b} x(-1)^a y^b, \\ & & a+b &\leq k, \end{aligned}$$

$$(5) \quad \begin{aligned} \rho_1 &= x(-1)^b y^a, & \rho_2 &= y(-1)^{k+1-a} y^a, \\ & & a &\geq 1, a+b \leq k, \end{aligned}$$

$$(7) \quad \begin{aligned} \rho_1 &= x(-1)^b y^a, & \rho_2 &= y(-1)^{k+1-a-c} h(-1)^c y^a, \\ & & a &\geq 1, c \geq 1, a+b \leq k, \end{aligned}$$

$$(11) \quad \begin{aligned} \rho_1 &= x(-1)^b y^c, & \rho_2 &= x(-2)^{k+1-a-b} h(-1)^a x(-1)^b, \\ & & b &\geq 1, a+b \leq k, b+c \leq k. \end{aligned}$$

We shall say that the cases listed in Lemma 10.6 are *exceptional cases*.

It is clear that if (10.1) holds for $\rho_1, \rho_2 \subset \pi$, then it holds as well for $\rho_1, \rho_2 \subset \kappa\pi$ for any $\kappa \in \mathcal{P}(\bar{B}_-)$. In particular it is enough to prove (10.1) for any two embeddings $\rho_1, \rho_2 \subset \rho_1 \cup \rho_2$. By Lemmas 10.3 and 10.4 it is enough to show (10.1) for $\rho_1, \rho_2 \subset \rho_1 \cup \rho_2$, where one ρ_i is short, the other one is long. The case $\rho_1 \cap \rho_2 = \emptyset$ is proved by Lemma 10.5. From (10.9) we see that (10.1) holds in the case that a short ρ_1 and a long ρ_2 satisfy (10.8). Moreover, if (10.1) holds for all embeddings of length less than $\ell(\rho_1 \cup \rho_2)$, then (10.1) holds when ρ_1 and ρ_2 are $\{y, x(-1)\}$ -linked. What is left is to prove (10.1) in the exceptional cases. We shall prove this by double induction on degree $0, -1, -2, \dots$ of $\rho_1 \cup \rho_2$ and, for a fixed degree, on length $\ell(\rho_1 \cup \rho_2)$.

We shall use the following modification of Lemma 9.4:

Lemma 10.7. *Let*

$$u(\rho \subset \pi)v_\Lambda = \sum C_{\rho', \pi'} u(\rho' \subset \pi')v_\Lambda,$$

where for each $\pi' \prec \pi$ the term $u(\rho' \subset \pi')$ appears for only one embedding $\rho' \subset \pi'$. Then $C_{\pi'} = C_{\rho', \pi'} = 0$ for $\pi' \prec \pi$. In particular

$$u(\rho \subset \pi)v_\Lambda \in \sum C_{\rho', \pi} u(\rho' \subset \pi)v_\Lambda + M_{(\pi)}^1.$$

At some places in the proof it will be convenient to write $a \underset{-p}{\sim} b$ if parts of $\ell(a)$ and $\ell(b)$ are of degree $0, -1, \dots, -p$, and $a - b$ is a linear combination of terms of the form $u(\rho \subset \pi)v_\Lambda$, where π has at least one part of degree $\leq -p - 1$.

Now we consider all exceptional cases:

(4) Let $\pi = h(-1)^{k+1-a-b}x(-1)^ay^bh^{k+1-a-b}$, $a + b \leq k$, and note that

$$r(\rho_1) = r(x(-1)^ay^bh^{k+1-a-b}).$$

By using in $k + 1 - a - b$ steps the relations (10.4) of the form $C^h = 1$, $C^x = 0$ we get

$$h(-1)^{k+1-a-b}r(\rho_1)v_\Lambda \underset{-1}{\sim} \sum C_{cd} y(-1)^cy^dr(\rho)v_\Lambda.$$

Consider two classes of terms with a leading term π_1 : (A) $y(-1)^c \subset \pi_1$ for some $c > 0$ and (B) terms of the form $y^dr(\rho)v_\Lambda$. Recall that by Proposition 10.2(a) for the term $\dots r(\rho)v_\Lambda$ we may assume that x is not a part of ρ . By Lemma 9.5(4) in the case (B) we have either

$$\begin{aligned} \rho &= h(-1)^{k+1-b-p}x(-1)^py^b, \quad p > 0, \quad \text{or} \\ \rho &= x(-1)^{k+1-b}y^b, \quad \pi_1 = x(-1)^{k+1-b}y^{k+1-a}. \end{aligned}$$

In either case there is only one embedding.

In the case (A) we may apply the induction hypothesis on $\pi_1/y(-1)^c$ and rewrite the terms in the class (A) so that each π_1 appears with only one embedding.

Finally we should consider the terms $u(\rho \subset \pi_1)v_\Lambda$, where π_1 contains a part of degree ≤ -2 , say $b(j)$. Since $|\pi_1| \geq -k - 1$, the part $b(j)$ cannot be a part of any embedding $\rho' \subset \pi_1$. Hence we may apply induction hypothesis on $\pi_1/b(j)$ and rewrite a combination of such terms so that each π_1 appears with only one embedding.

Now we can apply Lemma 10.7 and (10.1) follows.

Remark. Note that the proof for the case (4) is only a slight modification of the argument given in the exceptional case (4) in Section 9. With some other exceptional cases we have a similar situation as well.

(5) Set $\pi = y(-1)^{k+1-a}x(-1)^by^ah^{k+1-a-b}$, $a \geq 1$, $a + b \leq k$, and note that

$$r(\rho_1) = r(x(-1)^by^ah^{k+1-a-b}).$$

For $\text{sh } \rho = (-1)^{k+1-p} 0^p$, $p \geq 1$, we shall express the term $y(-1)r(\rho)$ by using the relation (10.4) of the form $C^y = 1$, $C_y = 0$. Note that

$$y(-1)r(\rho)v_\Lambda \underset{-2}{\simeq} Cr(\rho')v_\Lambda + \sum C_{j,X} X(j)r(\rho_{j,X})v_\Lambda,$$

where $X \in \{x, h, y\}$, $j \in \{-1, -2\}$, and that $\text{sh } \rho' = (-1)^{k+1-p+1} 0^{p-1}$, $\text{sh } \rho_{j,X} = (-1)^{k+1-q} 0^q$, $q \geq p$. We can use this type of relations to express $y(-1)^{k+1-a}r(\rho_1)$ by applying the relations on terms $u(\rho \subset \pi_1)$ with $\text{sh } \rho = (-1)^{k+1-p} 0^p$, $p \geq 1$, and no parts of π_1 of degree ≤ -2 (note that then $p \leq k+1-b$):

$$\begin{aligned} & y(-1)^{k+1-a}r(\rho_1)v_\Lambda \\ & \underset{-2}{\simeq} \sum C_{c\dots h} y(-2)^c h(-2)^d x(-2)^e y(-1)^f h(-1)^g x(-1)^h r(\rho)v_\Lambda. \end{aligned}$$

For terms with $c+d+e > 0$ we may apply the induction hypothesis and rewrite their sum in such a way that for each leading term π' there is only one embedding $\rho \subset \pi'$ which contributes with a term $u(\rho \subset \pi')v_\Lambda$.

Now note that for the term uv_Λ for which $\text{sh } \rho = (-1)^{k+1}$ we have (by the way it was obtained) $\ell(\ell(uv_\Lambda)) \leq k+1-a+b < k+1+b$, and hence $\ell(uv_\Lambda) \succ \ell(y(-1)^{k+1-a}r(\rho_1)v_\Lambda)$. Since we want to apply Lemma 10.7, we need not consider further such terms. What is left to consider is terms with $\text{sh } \rho = (-1)^{k+1-p} 0^p$, $p \geq 1$. Then either $f = 0$ or $f > 0$ and $\rho = y(-1)^{k+1-p} y^p$.

For a term uv_Λ , $u = h(-1)^g x(-1)^h r(\rho)$, $\text{wt } u = \text{wt } \pi = -(k+1)+b$, we must have (since $p \leq k+1-b$)

$$\rho = y(-1)^q h(-1)^{k+1-p-q} y^p, \quad q \geq 0.$$

Finally note that for terms omitted when writing $\underset{-2}{\simeq}$, i.e. terms of the form $u(\rho \subset \pi')v_\Lambda$, where π' has at least one part of the form $X(j)$, $j \leq -3$, $X \in \{x, h, y\}$, we may apply the induction hypothesis (since $X(j)$ cannot be a part of ρ) and rewrite their sum in such a way that for each leading term π' there is only one embedding $\rho \subset \pi'$ which contributes with a term $u(\rho \subset \pi')v_\Lambda$.

Hence we can apply Lemma 10.7.

$$\begin{aligned} (7) \quad & \rho_1 = x(-1)^b y^a, \quad a \geq 1, a+b \leq k, \\ & \rho_2 = y(-1)^{k+1-a-c} h(-1)^c y^a, \quad c \geq 0. \end{aligned}$$

Note that we have already considered the case $c = 0$ in (5). We keep the shape

$$(-1)^{k+1+b-a} 0^a$$

fixed and prove (10.1) by induction on c , $0 \leq c \leq k+1-a$. By the induction hypothesis we have

$$\begin{aligned} & y(-1)^{k+2-a-c} h(-1)^{c-1} r(x(-1)^b y^a)v_\Lambda \in \\ & \in \mathbb{F}^\times x(-1)^b r(y(-1)^{k+2-a-c} h(-1)^{c-1} y^a)v_\Lambda \end{aligned}$$

$$+M_{(y(-1)^{k+2-a-c}h(-1)^{c-1}x(-1)^by^a)}^1.$$

We get a relation (10.1) for $\pi = \rho_1 \cup \rho_2$ by adjoint action of x .

(11) Let $a \geq 0, b \geq 1, c \geq 0, a + b \leq k, b + c \leq k$,

$$\begin{aligned}\pi &= x(-2)^{k+1-a-b}h(-1)^ax(-1)^by^ch^{k+1-b-c}, \\ \rho_2 &= x(-2)^{k+1-a-b}h(-1)^ax(-1)^b,\end{aligned}$$

and note that $r(\rho_1) = r(x(-1)^by^ch^{k+1-b-c})$.

First we consider the case $a = 0$: the relation (9.20) gives

$$\begin{aligned}(10.10) \quad & y(-2)^{k+1-b}r(y(-1)^by^{k+1-b})v_\Lambda \simeq_2 y^{k+1-b}r(y(-2)^{k+1-b}y(-1)^b)v_\Lambda \\ & + \sum_{\substack{d+e+f+k+1=\ell(\pi) \\ d+e < k+1}} C_{def} y(-2)^dy(-1)^ey^fr(y(-1)^py^{k+1-p})v_\Lambda.\end{aligned}$$

By adjoint action of x on (10.10) we get

$$\begin{aligned}(10.11) \quad & x(-2)^{k+1-b}r(h(-1)^px(-1)^{b-p}y^{k+1-b})v_\Lambda \\ & \simeq_2 y^{k+1-b}r(x(-2)^{k+1-b}h(-1)^px(-1)^{b-p})v_\Lambda \\ & + \sum_{d+\dots+i < k+1} C_{d\dots j} y(-2)^dh(-2)^ex(-2)^fy(-1)^gh(-1)^hx(-1)^iy^jr(\rho)v_\Lambda\end{aligned}$$

for $0 \leq p \leq b$, and

$$\begin{aligned}(10.12) \quad & x(-2)^{k+1-b}r(x(-1)^by^c)v_\Lambda \simeq_2 y^cr(x(-2)^{k+1-b}x(-1)^b)v_\Lambda \\ & + \sum_{d+\dots+i < k+1} C_{d\dots j} y(-2)^dh(-2)^ex(-2)^fy(-1)^gh(-1)^hx(-1)^iy^jr(\rho)v_\Lambda,\end{aligned}$$

where for each summand $u(\rho \subset \pi_1)$ in (10.11) and (10.12) ρ has no parts of degree -2 , so we may apply the induction hypothesis on $\pi_1/y(-2)^dh(-2)^ex(-2)^f$ and then apply Lemma 10.7. In particular (10.1) holds in the case $a = 0$.

Now we prove the case $a \geq 1$: By using in a steps relations (10.4) of the form $C^h = 1, C^x = 0$ we get

$$(10.13) \quad h(-1)^ar(\rho_1) \simeq_1 \sum C_{de} y(-1)^dy^er(\rho),$$

where $0 \leq d \leq a$. Multiply both sides of (10.13) with $x(-2)^{k+1-a-b}$; in lack of better notation we will write this as

$$(10.14) \quad x(-2)^{k+1-a-b}(h(-1)^ar(\rho_1) \simeq_1 \sum C_{de} y(-1)^dy^er(\rho)).$$

Consider four disjoint classes of $\pi_1 \prec x(-2)^{k+1-a-b}h(-1)^ax(-1)^by^c$; in the case $\text{sh } \pi_1 \subset \pi$ three classes: (A) $y(-1)^d \subset \pi_1$ for some $d \geq 1$, (B) π_1 is not in A and $y(-2)^ph(-2)^qx(-2)^{k+1-a-b-p-q} \subset \pi_1$ for some $p+q \geq 1$ and (C) π_1 is not in A

and $x(-2)^{k+1-a-b} \in \pi_1$, and the fourth class (D) $\text{sh } \pi_1 \not\subseteq \pi$, i.e. π_1 has either a part of degree ≤ -3 or more than $k+1-a-b$ parts of degree -2 .

Let π_1 be in the class A.

For two embeddings $\rho, \rho' \subset \pi_1$ (i.e. $\rho, \rho' \subset \pi_1/y(-1)^d$ by Lemma 9.5), by induction hypothesis on degree, terms $u(\rho \subset \pi_1)$ and $u(\rho' \subset \pi_1)$ are proportional modulo higher terms (with leading terms $\succ \pi_1$). These higher terms are in class A or D.

Let π_1 be in the class B.

For two embeddings $\rho, \rho' \subset \pi_1$ ($\rho, \rho' \subset \pi_1/y(-2)^ph(-2)^qx(-2)^{k+1-a-b-p-q}$ because of Lemma 9.5), by induction hypothesis on degree, terms $u(\rho \subset \pi_1)$ and $u(\rho' \subset \pi_1)$ are proportional modulo higher terms (with leading terms $\succ \pi_1$). These higher terms are in class B or D.

Let π_1 be in the class D.

Then for $\rho \subset \pi_1$ we have that $\text{sh } \rho \subset (-1)^{k+1}0^{k+1}$. As above, for two embeddings $\rho, \rho' \subset \pi_1$, by induction hypothesis on degree, terms $u(\rho \subset \pi_1)$ and $u(\rho' \subset \pi_1)$ are proportional modulo higher terms (with leading terms $\succ \pi_1$). These higher terms are again in class D.

Now consider a term in (10.14) which has the leading term π_1 in the class C. Then $d = 0$. For $\rho \subset \pi_1$ (i.e. $\rho \subset \pi_1/y^e$) the term $u(\rho \subset \pi_1)$ is (by using the relation of the form (10.11)) proportional to

$$y^{e'} r(x(-2)^{k+1-a-b} h(-1)^{a'} x(-1)^{b'})$$

modulo terms with leading terms $\pi' \succ \pi_1$.

Hence in a finite number of steps we can rewrite (first the terms in class C, and then the terms in class A, B and D) the right hand side of (10.14), modulo $M_{(x(-2)^{k+1-a-b} h(-1)^a x(-1)^b y^e)}^1$, as a linear combination of terms $u(\rho \subset \pi_1)$, where for each $\pi_1 \prec \pi$ there is only one embedding. By applying Lemma 10.7 we get (10.1).

11. SOME COMBINATORIAL IDENTITIES OF ROGERS-RAMANUJAN TYPE

11.1. Lepowsky-Wakimoto product formulas.

Let $A = (a_{ij})$ be an $(\ell+1) \times (\ell+1)$ generalized Cartan matrix (GCM) so that the corresponding Kac-Moody Lie algebra $\mathfrak{g}(A)$ is an affine Lie algebra. Use the usual notation for roots $\{\alpha_0, \alpha_1, \dots, \alpha_\ell\} \subset \mathfrak{h}^*$ and coroots $\{\alpha_0^\vee, \alpha_1^\vee, \dots, \alpha_\ell^\vee\} \subset \mathfrak{h}$, $a_{ij} = \langle \alpha_i^\vee, \alpha_j \rangle$. Let $\mathbf{s} = (s_0, s_1, \dots, s_\ell) \in (\mathbb{Z}_{\geq 0})^{\ell+1}$ and let $\rho_{\mathbf{s}}^\vee \in \mathfrak{h}$ be an element such that $\langle \rho_{\mathbf{s}}^\vee, \alpha_i \rangle = s_i$ for $i = 0, 1, \dots, \ell$. Then $\{\alpha_0^\vee, \alpha_1^\vee, \dots, \alpha_\ell^\vee, \rho_{\mathbf{s}}^\vee\}$ is a basis of $\mathfrak{h}$. Write $\rho^\vee$ instead of $\rho_{\mathbf{s}}^\vee$ in the case $\mathbf{s} = (1, 1, \dots, 1)$.

Recall the Weyl-Kac character formula for the standard $\mathfrak{g}(A)$ -module $L(\Lambda)$ with highest weight Λ (cf. [K]):

$$(11.1.1) \quad e^{-\Lambda} \text{ch } L(\Lambda) = N(\Lambda + \rho; A)/D(A),$$

where

$$N(\Lambda + \rho; A) = \sum_{w \in W(A)} \varepsilon(w) e^{w(\Lambda + \rho) - (\Lambda + \rho)},$$

$$D(A) = \sum_{w \in W(A)} \varepsilon(w) e^{w\rho - \rho} = \prod_{\alpha > 0} (1 - e^{-\alpha})^{\dim \mathfrak{g}(A)_\alpha}.$$

Then after applying the homomorphism $e^{-\alpha} \mapsto q^{\langle \rho_s^\vee, \alpha \rangle}$ (i.e. by $\rho_s^\vee$ -specialization) we get power series in q :

$$(11.1.2) \quad q^{\langle \Lambda, \rho_s^\vee \rangle} \text{ch}^{\rho_s^\vee} L(\Lambda) = N^{\rho_s^\vee}(\Lambda + \rho; A) / D^{\rho_s^\vee}(A),$$

where

$$N^{\rho_s^\vee}(\Lambda + \rho; A) = \sum_{w \in W(A)} \varepsilon(w) q^{-\langle \rho_s^\vee, w(\Lambda + \rho) - (\Lambda + \rho) \rangle},$$

$$D^{\rho_s^\vee}(A) = \sum_{w \in W(A)} \varepsilon(w) q^{-\langle \rho_s^\vee, w\rho - \rho \rangle} = \prod_{\alpha > 0} (1 - q^{\langle \rho_s^\vee, \alpha \rangle})^{\dim \mathfrak{g}(A)_\alpha}.$$

Note that $\langle \rho^\vee, w(\Lambda + \rho) - (\Lambda + \rho) \rangle = \langle \Lambda + \rho, w\rho^\vee - \rho^\vee \rangle$ implies Lepowsky's numerator formula $N^{\rho^\vee}(\Lambda + \rho; A) = D^{\Lambda + \rho}(\tau A)$ (cf. [L]). M. Wakimoto generalized this formula in the following way (cf. [W]): Let $\tilde{A} = (\tilde{a}_{ij})$ be another GCM of affine type with roots $\{\tilde{\alpha}_0, \tilde{\alpha}_1, \dots, \tilde{\alpha}_\ell\} \subset \tilde{\mathfrak{h}}^*$ and coroots $\{\tilde{\alpha}_0^\vee, \tilde{\alpha}_1^\vee, \dots, \tilde{\alpha}_\ell^\vee\} \subset \tilde{\mathfrak{h}}$, $\tilde{a}_{ij} = \langle \tilde{\alpha}_i^\vee, \tilde{\alpha}_j \rangle$. Fix $\tilde{\rho}^\vee, \langle \tilde{\rho}^\vee, \tilde{\alpha}_i \rangle = 1$ for $i = 0, 1, \dots, \ell$. For fixed $\mathbf{s} = (s_0, s_1, \dots, s_\ell)$ define isomorphism $\Theta_s : \mathfrak{h}^* \rightarrow \tilde{\mathfrak{h}}^*$ so that $\langle \tilde{\alpha}_i^\vee, \Theta_s(\lambda) \rangle = s_i \langle \alpha_i^\vee, \lambda \rangle$, $\langle \tilde{\rho}^\vee, \Theta_s(\lambda) \rangle = \langle \rho_s^\vee, \lambda \rangle$. Now assume that $\tilde{a}_{ij} = s_i a_{ij} s_j^{-1}$. Then it is easy to check that $\tilde{r}_i = \Theta_s r_i \Theta_s^{-1}$, where $\tilde{r}_i$ and r_i are reflections with respect to simple roots $\tilde{\alpha}_i$ and α_i (respectively). Hence $w \mapsto \tilde{w} = \Theta_s w \Theta_s^{-1}$ defines an isomorphism between the Weyl groups $W(A)$ of $\mathfrak{g}(A)$ and $W(\tilde{A})$ of $\mathfrak{g}(\tilde{A})$ (or $\mathfrak{g}(\tau \tilde{A})$). We also have $\langle \rho_s^\vee, w\lambda \rangle = \langle \tilde{\rho}^\vee, \tilde{w}(\Theta_s \lambda) \rangle$. This implies the first statement in the following theorem:

Theorem 11.1.1. [W]. *If $\tilde{a}_{ij} = s_i a_{ij} s_j^{-1}$, $i, j = 0, 1, \dots, \ell$, then*

$$(a) \quad N^{\rho_s^\vee}(\Lambda + \rho; A) = D^{\Theta_s(\Lambda + \rho)}(\tau \tilde{A}),$$

$$(b) \quad N^{\rho_s^\vee}(\Lambda + \rho; A) = N^{\Lambda + \rho}(\rho_s^\vee; \tau A).$$

Formula (a) implies that for certain choices of $\mathbf{s}$ the $\rho_s^\vee$ -specialization of characters of all standard modules can be written as infinite products. The “duality” formula (b) implies that for certain (depending on $\rho_s^\vee$) standard modules all $(\Lambda + \rho)$ -specializations of characters can be written as infinite products.

Now let $\mathfrak{g}(A) = \tilde{\mathfrak{g}} = \mathfrak{sl}(2, \mathbb{F})^\sim$, i.e. $A = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$. For $\mathbf{s} = (n, n)$, $(n, 2n)$ or $(2n, n)$ Theorem 11.1.1 may be applied. In the case $\mathbf{s} = (n, n)$ we have $\tilde{A} = A$ and (a) is Lepowsky's numerator formula. In the case $(n, 2n)$ we have $\tilde{A} = \begin{pmatrix} 2 & -1 \\ -4 & 2 \end{pmatrix}$, $\mathfrak{g}(\tau \tilde{A})$ is of the type $A_2^{(2)}$, $\tilde{\alpha}_1^\vee$ is a long root, $\tilde{\alpha}_0^\vee$ is a short root. The case $(2n, n)$ is related to $(n, 2n)$ by an outer automorphism. For $\mathbf{s} = (s_0, s_1)$, $A = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$, and for $\mu \in \tilde{\mathfrak{h}}^*$, $\mu(\tilde{\alpha}_0^\vee) = m_0 \in \mathbb{Z}_{>0}$, $\mu(\tilde{\alpha}_1^\vee) = 2m_1 \in \mathbb{Z}_{>0}$, $\tilde{A} = \begin{pmatrix} 2 & -1 \\ -4 & 2 \end{pmatrix}$, write

$$P(s_0, s_1; q) = D^{\rho_s^\vee}(A), \quad Q(m_0, 2m_1; q) = D^\mu(\tau \tilde{A}).$$

Lemma 11.1.2. *Let $s_0, s_1, m_0 \in \{1, 2, 3, \dots\}$, $m_1 \in \{\frac{1}{2}, 1, \frac{3}{2}, \dots\}$, $s = s_0 + s_1$, $m = m_0 + m_1$. Then*

$$(a) \quad P(s_0, s_1; q) = \prod_{r \equiv 0, s_0 \pmod s} (1 - q^r) \prod_{r \equiv s_1 \pmod s} (1 - q^r),$$

$$(b) \quad Q(m_0, 2m_1; q) = \prod_{r \equiv 0, \pm m_0 \pmod{2m}} (1 - q^r) \prod_{r \equiv \pm 2m_1 \pmod{4m}} (1 - q^r).$$

(Here is assumed $r \in \mathbb{Z}_{>0}$.)

For $\mathbf{s} = (s_0, s_1)$ and $\Lambda = k_0\Lambda_0 + k_1\Lambda_1$ write $d_{k_0, k_1}^{s_0, s_1}(q) = q^{\langle \Lambda, \rho_{\mathbf{s}}^\vee \rangle} \text{ch}^{\rho_{\mathbf{s}}^\vee} L(\Lambda)$. Then Theorem 11.1.1 and Lemma 11.1.2 give Lepowsky-Wakimoto product formulas for specialized characters of standard modules for $\mathfrak{sl}(2, \mathbb{F})^\sim$:

$$(11.1.3) \quad d_{k_0, k_1}^{1,1}(q) = P(k_0+1, k_1+1; q)/P(1, 1; q),$$

$$(11.1.4) \quad d_{k_0, k_1}^{1,2}(q) = Q(k_0+1, 2(k_1+1); q)/P(1, 2; q),$$

$$(11.1.5) \quad d_{n-1, 2n-1}^{s_0, s_1}(q) = Q(ns_0, 2ns_1; q)/P(s_0, s_1; q),$$

$$(11.1.6) \quad d_{2n-1, n-1}^{s_0, s_1}(q) = Q(ns_1, 2ns_0; q)/P(s_0, s_1; q),$$

$$(11.1.7) \quad d_{n-1, n-1}^{s_0, s_1}(q) = P(ns_0, ns_1; q)/P(s_0, s_1; q).$$

It is easy to see the following:

Proposition 11.1.3. *Each of the products (11.1.3)–(11.1.7) is of the form*

$$(11.1.8) \quad \prod_i \prod_{r \in B_i} (1 + q^r) \prod_j \prod_{r \in C_j} (1 - q^r)^{-1}$$

for some congruence classes $B_1, \dots, B_b \subset \mathbb{Z}_{>0}$, $b \geq 0$, $C_1, \dots, C_c \subset \mathbb{Z}_{>0}$, $c \geq 1$. (These classes need not be disjoint, some may be empty.)

As particular examples we have:

$$(11.1.9) \quad d_{n-1, n-1}^{1,1}(q) = \prod_{r \text{ odd}, r \not\equiv b \pmod{2b}} (1 - q^r)^{-1} \prod_{r \not\equiv 0, n \pmod{2n}} (1 - q^r)^{-1} \\ \cdot \prod_{1 \leq j \leq a} \prod_{r \equiv 2^{a-j}b \pmod{2^{a-j+1}b}} (1 + q^r), \quad \text{for } n = 2^a b, a \geq 0, b \text{ odd},$$

$$(11.1.10) \quad d_{n-1, 2n-1}^{1,1}(q) = \prod_{r \text{ odd}} (1 - q^r)^{-1} \prod_{r \not\equiv 0, \pm n \pmod{3n}} (1 - q^r)^{-1},$$

$$(11.1.11) \quad d_{k_0, k_1}^{1,1}(q) = \prod_{r \text{ odd}} (1 - q^r)^{-1} \prod_{r \not\equiv 0, \pm(k_0+1) \pmod{(k+2)}} (1 - q^r)^{-1}$$

for $k = k_0 + k_1 \neq 2k_1$,

$$(11.1.12) \quad d_{k_0, k_1}^{1,2}(q) = \prod_{\substack{r \not\equiv 0, \pm(k_0+1) \pmod{2(k+2)} \\ r \not\equiv \pm 2(k_1+1) \pmod{4(k+2)}}} (1 - q^r)^{-1}$$

for $k = k_0 + k_1 \neq 3k_1 + 1$,

$$(11.1.13) \quad d_{2n-1, n-1}^{1,2}(q) = \prod_{r \equiv \pm n \pmod{6n}} (1 + q^r) \prod_{r \not\equiv 0, \pm n, \pm 2n \pmod{6n}} (1 - q^r)^{-1},$$

$$(11.1.14) \quad d_{n-1, n-1}^{1,2}(q) = \prod_{r \not\equiv 0 \pmod{n}} (1 - q^r)^{-1}.$$

For nonempty subsets $A_1, A_2, \dots, A_s \subset \mathbb{Z}_{>0}$, $s \geq 2$, set $A = A_1 \sqcup A_2 \sqcup \dots \sqcup A_s$, a disjoint union of sets. We may call elements of $\mathcal{P}(A)$ *colored partitions* with parts in A , where for $i \in \{1, \dots, s\}$ and $j \in A_i$ we say that j is of *color* i and of *weight* $|j| = j \in \mathbb{Z}_{>0}$. We define the *degree* $|\pi|$ of $\pi \in \mathcal{P}(A)$ as before: $|\pi| = \sum_{a \in A} \pi(a)|a|$. It is clear that infinite products of the form (11.1.8) may be interpreted as generating functions of partition functions for colored partitions defined by congruence conditions.

11.2. Difference conditions and partition ideals.

Let A be a nonempty set and $\mathcal{P}(A)$ the set of all partitions with parts in A . By following [A1] and [A2] we shall say that $\mathcal{C} \subset \mathcal{P}(A)$ is a *partition ideal* in $\mathcal{P}(A)$ if $\kappa \in \mathcal{C}$ and $\pi \in \mathcal{P}(A)$ implies $\kappa \cap \pi \in \mathcal{C}$, or equivalently, if $\mu \in \mathcal{C}$, $\nu \in \mathcal{P}(A)$, $\nu \subset \mu$, implies $\nu \in \mathcal{C}$. If $\mathcal{C}, \mathcal{I} \subset \mathcal{P}(A)$, $\mathcal{P}(A) = \mathcal{C} \cup \mathcal{I}$, $\mathcal{C} \cap \mathcal{I} = \emptyset$, then it is easy to see that $\mathcal{C}$ is a partition ideal in $\mathcal{P}(A)$ if and only if $\mathcal{I}$ is an ideal in the monoid $\mathcal{P}(A)$. In particular,

$$\mathcal{C}_\Lambda = \mathcal{P}(\bar{B}_-) \setminus (\ell t(\bar{R}v_\Lambda))$$

is a partition ideal in $\mathcal{P}(\bar{B}_-)$. By Theorem 6.5.5 the partition ideal $\mathcal{C}_\Lambda$ determines the character of standard $\mathfrak{g}(A)$ -module $L(\Lambda)$ with highest weight Λ :

$$(11.2.1) \quad e^{-\Lambda} \text{ch } L(\Lambda) = \sum_{\pi \in \mathcal{C}_\Lambda} e^{\text{wt } \pi + |\pi| \delta}.$$

In Section 6.6 we have described the set $\ell t(\bar{R}v_\Lambda)$, and as a consequence we may describe the partition ideal $\mathcal{C}_\Lambda$, $\Lambda = k_0 \Lambda_0 + k_1 \Lambda_1$, $k = k_0 + k_1$, as the set of all colored partitions π in $\mathcal{P}(\bar{B}_-)$ satisfying *difference conditions*:

$$(11.2.2) \quad \begin{aligned} \pi(y(j-1)) + \pi(h(j-1)) + \pi(y(j)) &\leq k, \\ \pi(h(j-1)) + \pi(x(j-1)) + \pi(y(j)) &\leq k, \\ \pi(x(j-1)) + \pi(y(j)) + \pi(h(j)) &\leq k, \\ \pi(x(j-1)) + \pi(h(j)) + \pi(x(j)) &\leq k, \end{aligned}$$

and *initial conditions*:

$$(11.2.3) \quad \pi(x(-1)) \leq k_0, \quad \pi(y(0)) \leq k_1.$$

For $\mathbb{B} \subset \mathbb{Z}_{>0}$ we shall write an ordinary partition $f: \mathbb{B} \rightarrow \mathbb{N}$ (with parts in $\mathbb{B}$) by its values $(f_r \mid r \in \mathbb{B})$. For $\mathbf{s} = (s_0, s_1)$, $s_0, s_1 \geq 1$, $s_0 \neq s_1$, $s = s_0 + s_1$, we have a bijection $\bar{B}_- \rightarrow \mathbb{B}^{s_0, s_1}$, where

$$(11.2.4) \quad \mathbb{B}^{s_0, s_1} = \{r \in \mathbb{Z}_{>0} \mid r \equiv 0, \pm s_1 \pmod{s}\},$$

defined by the specialization

$$(11.2.5) \quad \begin{aligned} y(-j) &\mapsto \deg y(-j) = s_1 + js, & j \in \mathbb{Z}_{\geq 0}, \\ h(-j) &\mapsto \deg h(-j) = js, & j \in \mathbb{Z}_{> 0}, \\ x(-j) &\mapsto \deg x(-j) = -s_1 + js, & j \in \mathbb{Z}_{> 0}. \end{aligned}$$

This bijection extends to an isomorphism of monoids $\mathcal{P}(\bar{B}_-) \rightarrow \mathcal{P}(\mathbb{B}^{s_0, s_1})$. Hence we have a bijection of partition ideals

$$\mathcal{C}_\Lambda \rightarrow \mathcal{C}_{k_0, k_1}^{s_0, s_1},$$

where $\mathcal{C}_{k_0, k_1}^{s_0, s_1} \subset \mathcal{P}(\mathbb{B}^{s_0, s_1})$ is the set of all partitions $f: \mathbb{B}^{s_0, s_1} \rightarrow \mathbb{N}$ which satisfy *difference conditions*:

$$(11.2.6) \quad \begin{aligned} f_{js+s_1} + f_{js} + f_{js-s_0} &\leq k, \\ f_{js} + f_{js-s_1} + f_{js-s_0} &\leq k, \\ f_{js+s_0} + f_{js+s_1} + f_{js} &\leq k, \\ f_{js+s_0} + f_{js} + f_{js-s_1} &\leq k, \end{aligned}$$

and *initial conditions*:

$$(11.2.7) \quad f_{s_0} \leq k_0, \quad f_{s_1} \leq k_1.$$

As a particular example in the case $\mathbf{s} = (1, 2)$ we have $\mathbb{B}^{1,2} = \mathbb{Z}_{>0}$ and the partition ideal $\mathcal{C}_{k_0, k_1}^{1,2}$ in $\mathcal{P}(\mathbb{Z}_{>0})$ is defined by

$$(11.2.8) \quad \begin{aligned} f_{3j+2} + f_{3j} + f_{3j-1} &\leq k, \\ f_{3j} + f_{3j-1} + f_{3j-2} &\leq k, \\ f_{3j+2} + f_{3j+1} + f_{3j} &\leq k, \\ f_{3j+1} + f_{3j} + f_{3j-2} &\leq k, \\ f_1 &\leq k_0, \quad f_2 \leq k_1. \end{aligned}$$

For a nonempty set A we shall write a partition $\pi: A \rightarrow \mathbb{N}$ (with parts in A) by its values $(\pi_a \mid a \in A)$. Now consider the case of principal specialization, i.e. $\mathbf{s} = (1, 1)$. Set

$$A_1 = \{1, 2, 3, \dots\} \cong \mathbb{Z}_{>0}, \quad A_2 = \{\underline{1}, \underline{2}, \underline{3}, \dots\} \cong \mathbb{Z}_{>0}, \quad A = A_1 \bigsqcup A_2.$$

We shall say that $j \in A_1$ is of degree $|j| = j \in \mathbb{Z}_{>0}$ and of plain color, and that $\underline{j} \in A_2$ is of degree $|\underline{j}| = j \in \mathbb{Z}_{>0}$ and of underline color. Set $\underline{B} = A_1 \bigsqcup \{\underline{j} \in A_2 \mid j \text{ odd}\}$. Then we have a bijection $\bar{B}_- \rightarrow \underline{B}$ defined by the specialization

$$(11.2.9) \quad \begin{aligned} y(-i) &\mapsto 2i + 1 \in A_1, \quad i \in \mathbb{Z}_{\geq 0}, \\ h(-i) &\mapsto 2i \in A_1, \quad i \in \mathbb{Z}_{>0}, \\ x(-i) &\mapsto \underline{2i - 1} \in A_2, \quad i \in \mathbb{Z}_{>0}. \end{aligned}$$

This bijection extends to an isomorphism of monoids $\mathcal{P}(\bar{B}_-) \rightarrow \mathcal{P}(\underline{B})$. Hence we have a bijection of partition ideals

$$\mathcal{C}_\Lambda \rightarrow \mathcal{C}_{k_0, k_1}^{1,1},$$

where $\mathcal{C}_{k_0, k_1}^{1,1} \subset \mathcal{P}(\underline{B})$ is the set of all colored partitions $\pi: \underline{B} \rightarrow \mathbb{N}$ which satisfy *difference conditions*:

$$(11.2.10) \quad \begin{aligned} \pi_{2i+1} + \pi_{2i} + \pi_{2i-1} &\leq k, \\ \pi_{2i} + \pi_{\underline{2i-1}} + \pi_{2i-1} &\leq k, \\ \pi_{\underline{2i+1}} + \pi_{2i+1} + \pi_{2i} &\leq k, \\ \pi_{\underline{2i+1}} + \pi_{2i} + \pi_{\underline{2i-1}} &\leq k, \end{aligned}$$

and *initial conditions*:

$$(11.2.11) \quad \pi_{\underline{1}} \leq k_0, \quad \pi_1 \leq k_1.$$

11.3. Combinatorial identities.

Formulas (11.1.1) and (11.2.1) express the character $\text{ch } L(\Lambda)$ in two different ways which may have different combinatorial interpretations. In particular, if we take a specialization (11.2.5) for $s_0 \neq s_1$, then the specialized character $d_{k_0, k_1}^{s_0, s_1}(q)$ is a generating function for $\mathcal{C}_{k_0, k_1}^{s_0, s_1}$ -partition function, where the partition ideal $\mathcal{C}_{k_0, k_1}^{s_0, s_1}$ is defined by difference conditions (11.2.6), initial conditions (11.2.7) and congruence conditions (11.2.4). On the other side, for certain choices of $\mathbf{s}$ and/or Λ we get Lepowsky-Wakimoto product formulas (11.1.4)–(11.1.7) for specialized characters $d_{k_0, k_1}^{s_0, s_1}(q)$, and these have a combinatorial interpretation in terms of congruence conditions. As a consequence we get a series of Rogers-Ramanujan type combinatorial identities. In the case of principal specialization (11.2.9) (i.e. for $\mathbf{s} = (1, 1)$) the product formula (11.1.3) on one side and the conditions (11.2.10)–(11.2.11) on the other side give a series of Rogers-Ramanujan type combinatorial identities for colored partitions.

For example, in the case $\mathbf{s} = (1, 2)$ and $k_0 = k_1 = n - 1$ we have a product formula (11.1.14) for $d_{n-1, n-1}^{1, 2}(q)$, so (11.2.8) for the partition ideal $\mathcal{C}_{n-1, n-1}^{1, 2}$ in $\mathcal{P}(\mathbb{Z}_{>0})$ gives that for every $m \in \mathbb{N}$ the number of partitions f such that $|f| = m$,

$$f_j > 0 \quad \text{implies} \quad j \not\equiv 0 \pmod{n},$$

equals the number of partitions f such that $|f| = m$,

$$\begin{aligned} f_{3j+2} + f_{3j+1} + f_{3j} &\leq 2n - 2, \\ f_{3j+2} + f_{3j} + f_{3j-1} &\leq 2n - 2, \\ f_{3j+1} + f_{3j} + f_{3j-2} &\leq 2n - 2, \\ f_{3j} + f_{3j-1} + f_{3j-2} &\leq 2n - 2, \\ f_1 &\leq n - 1, \quad f_2 \leq n - 1. \end{aligned}$$

Of course, this also equals the number of partitions f of m such that each part of f appears at most $n - 1$ times, and the example given in the introduction is for the $(1, 2)$ -specialization of the level 2 standard module $L(\Lambda_0 + \Lambda_1)$.

By some sort of coincidence, the combinatorial identities for the $(1, 2)$ -specialization of the fundamental $\mathfrak{sl}(2, \mathbb{C})$ -modules are identical with the combinatorial identities for the level 3 modules in the principal picture for the type $A_2^{(2)}$ affine Lie algebra (cf. [C1], [C2], [A3]). Note that in this case (i.e. the level 1 $\mathfrak{sl}(2, \mathbb{C})$ -modules) Lemma 9.2 is (essentially) all that is needed for the proof of linear independence.

Another example that we may take is the case $\mathbf{s} = (1, 1)$ and $k_0 = 1, k_1 = 2$. Then we have a product formula (11.1.11) for $d_{1, 2}^{1, 1}(q)$, so (11.2.10)–(11.2.11) for the partition ideal $\mathcal{C}_{1, 2}^{1, 1}$ in $\mathcal{P}(\underline{B})$ gives that for every $m \in \mathbb{N}$ the number of colored partitions $\pi \in \mathcal{P}(\underline{B})$ such that $|\pi| = m$,

$$\pi_a > 0 \quad \text{implies} \quad a \in \{\underline{i} \mid i \text{ odd}\} \bigsqcup \{i \mid i \equiv \pm 1 \pmod{5}\},$$

equals the number of colored partitions $\pi \in \mathcal{P}(\underline{B})$ such that $|\pi| = m$,

$$\begin{aligned} \pi_{\underline{2i+1}} + \pi_{2i+1} + \pi_{2i} &\leq 3, \\ \pi_{2i+1} + \pi_{2i} + \pi_{2i-1} &\leq 3, \\ \pi_{\underline{2i+1}} + \pi_{2i} + \pi_{\underline{2i-1}} &\leq 3, \\ \pi_{2i} + \pi_{\underline{2i-1}} + \pi_{2i-1} &\leq 3, \\ \pi_{\underline{1}} &\leq 1, \quad \pi_1 \leq 2. \end{aligned}$$

In a similar way we get that for every $m \in \mathbb{N}$ the number of colored partitions $\pi \in \mathcal{P}(\underline{B})$ such that $|\pi| = m$,

$$\pi_a > 0 \quad \text{implies} \quad a \in \{\underline{i} \mid i \text{ odd}\} \bigsqcup \{i \mid i \equiv \pm 2 \pmod{5}\},$$

equals the number of colored partitions $\pi \in \mathcal{P}(\underline{B})$ such that $|\pi| = m$,

$$\begin{aligned} \pi_{\underline{2i+1}} + \pi_{2i+1} + \pi_{2i} &\leq 3, \\ \pi_{2i+1} + \pi_{2i} + \pi_{2i-1} &\leq 3, \\ \pi_{\underline{2i+1}} + \pi_{2i} + \pi_{\underline{2i-1}} &\leq 3, \\ \pi_{2i} + \pi_{\underline{2i-1}} + \pi_{2i-1} &\leq 3, \\ \pi_{\underline{1}} &\leq 0, \quad \pi_1 \leq 3. \end{aligned}$$

REFERENCES

- [Ad] D. Adamović, *Some rational vertex algebras*, Glasnik Matematički **29** (1994), 25–40.
- [A1] G. E. Andrews, *A general theory of identities of the Rogers-Ramanujan type*, Bull. Amer. Math. Soc. **80** (1974), 1033–1052.
- [A2] G. E. Andrews, *The Theory of Partitions*, Encyclopedia of math. and appl., Addison-Wesley, Amsterdam, 1976.
- [A3] G. E. Andrews, *Schur's theorem, Capparelli's conjecture and the q -trinomial coefficients*, in Proc. Rademacher Centenary Conf. (1992), Contemporary Math. **167**, 1994, pp. 141–154.
- [B] R. E. Borcherds, *Vertex algebras, Kac-Moody algebras, and the Monster*, Proc. Natl. Acad. Sci. USA **83** (1986), 3068–3071.
- [C1] S. Capparelli, *On some representations of twisted affine Lie algebras and combinatorial identities*, J. Algebra **154** (1993), 335–355.
- [C2] S. Capparelli, *A combinatorial proof of a partition identity related to the level 3 representations of a twisted affine Lie algebra*, Comm. Algebra **23** (1995), 2959–2969.
- [CP] V. Chari, A. Pressley, *Integrable Representations of Kac-Moody Algebras: Results and Open Problems*, in Infinite Dimensional Lie algebras and groups, Advanced series in Math. Phys. Vol. 7 (V. G. Kac, ed.), World Scientific, Singapore, 1989.
- [DL] C. Dong, J. Lepowsky, *Generalized vertex algebras and relative vertex operators*, Progress in Mathematics **112**, Birkhäuser, Boston, 1993.
- [FF] B. Feigin, E. Frenkel, *Affine Kac-Moody algebras at the Critical Level and Gelfand-Dikii Algebras*, Intern. Jour. of Modern Physics A **7**, Suppl. 1A (1992), 197–215.
- [FHL] I. B. Frenkel, Y.-Z. Huang, J. Lepowsky, *On Axiomatic Approaches to Vertex Operator Algebras and Modules*, Memoirs of the Amer. Math. Soc. **104**, No. 494 (1993).
- [FK] I. B. Frenkel, V. G. Kac, *Basic representations of affine Lie algebras and dual resonance models*, Invent. Math. **62** (1980), 23–66.
- [FLM] I. B. Frenkel, J. Lepowsky, A. Meurman, *Vertex Operator Algebras and the Monster*, Pure and Applied Math., Academic Press, San Diego, 1988.
- [FNO] B. Feigin, T. Nakanishi, H. Ooguri, *The annihilating ideals of minimal models*, Intern. Jour. of Modern Physics A **7**, Suppl. 1A (1992), 217–238.
- [FZ] I. B. Frenkel, Y. Zhu, *Vertex operator algebras associated to representations of affine and Virasoro algebras*, Duke Math. J. **66** (1992), 123–168.
- [G] P. Goddard, *Meromorphic Conformal Field Theory*, in Infinite Dimensional Lie algebras and groups, Advanced series in Math. Phys. Vol. 7 (V. G. Kac, ed.), World Scientific, Singapore, 1989.
- [H] T. Hayashi, *Sugawara operators and Kac-Kazhdan conjecture*, Invent. Math. **94** (1988), 13–52.
- [K] V. G. Kac, *Infinite-dimensional Lie algebras* 3rd ed., Cambridge Univ. Press, Cambridge, 1990.
- [L] J. Lepowsky, *Lectures on Kac-Moody Lie algebras*, Université de Paris VI, 1978.

- [LW] J. Lepowsky, R. L. Wilson, *The structure of standard modules, I: Universal algebras and the Rogers-Ramanujan identities*, Invent. Math. **77** (1984), 199–290; *II: The case $A_1^{(1)}$, principal gradation*, Invent. Math. **79** (1985), 417–442.
- [Li] H.-S. Li, *Local systems of vertex operators, vertex superalgebras and modules*, J. of Pure and Appl. Alg. **109** (1996), 143–195.
- [MP] A. Meurman, M. Primc, *Annihilating Ideals of Standard Modules of $\mathfrak{sl}(2, \mathbb{C})^\sim$ and Combinatorial Identities*, Advances in Math. **64**, No. 3 (1987), 177–240.
- [W] M. Wakimoto, *Two formulae for specialized characters of Kac-Moody Lie algebras*, Hiroshima University, 1983 (preprint).
- [Xu] X. Xu, *Characteristics of Spinor Vertex Operator Algebras and Their Modules* (1992), Hong Kong Univ. of Science and Technology (Technical report 92-1-2).

UNIV. OF LUND, DEPT. OF MATHEMATICS, S-22100 LUND, SWEDEN
E-mail address: `arnem@maths.lth.se`

UNIV. OF ZAGREB, DEPT. OF MATHEMATICS, ZAGREB, CROATIA
E-mail address: `primc@cromath.math.hr`